







Weekly Report on Severe Acute Respiratory Infection (SARI), Week 11 2024 (week ending 17/03/2024)

This report includes data on SARI hospitalised cases, aged 15 years and older who were admitted to St. Vincent's University Hospital (SVUH), Dublin, up to week 11 2024. Please note that this report pertains to one hospital site only, data are not nationally representative. Therefore, caution is advised when interpreting rates and trends outlined in this report, as these may fluctuate due to the low case numbers.

Key points

Week 11 2024 (week ending 17/03/2024):

- **Number of cases:** 10 SARI cases admitted to the SARI hospital site, compared to 11 cases in week 10 2024 (9.1% decrease).
- Incidence rate per hospital catchment population: 3.2 per 100,000 population aged 15 years and older, compared to 3.5 per 100,000 in week 10 2024.
- Incidence rate per emergency hospitalisations: 31.2 per 1,000, compared to 35.6 per 1,000 in week 10 2024 (12.4% decrease).
- Age profile: Nine (90.0%) SARI cases were aged ≥65 years; Median age: 82 years; IQR: 76-86 years.
- Underlying medical conditions: Nine (90.0%) SARI cases reported having underlying medical conditions.
- **PCR testing:** Of those tested, one (10.0%) tested positive for SARS-CoV-2; two (20.0%) tested positive for influenza (two A(not subtyped)); and none tested positive for RSV.

Last four weeks (weeks 8 2024 - 11 2024)

- Number of cases: 59 SARI cases admitted to the SARI hospital site.
- Age profile: 47 (79.7%) SARI cases were aged ≥65 years; Median age: 78 years; IQR: 66-84 years.
- Underlying medical conditions: 53 (89.8%) SARI cases reported having underlying medical conditions.
- **PCR testing:** Of those tested, four (6.8%) tested positive for SARS-CoV-2; 12 (20.3%) tested positive for influenza (three A(not subtyped); seven A(H3); two A(H1)pdm09); and none tested positive for RSV.
- SARS-CoV-2 whole genome sequencing (WGS): There can be a lag-time before WGS results are available. Among those sequenced (n=1), one (100.0%) identified as variant BA.2 sub-lineage JN.1 and its sub-lineages.

Season 2023/2024 to date (weeks 40 2023 - 11 2024)

Collection of discharge data is a manual process, there is a significant lag time between discharge and data collection. Vaccination data is available approximately one week after cases are notified.

- Number of cases: 369 SARI cases admitted to the SARI hospital site.
- **PCR testing:** Of those tested, 43 (12.0%) tested positive for SARS-CoV-2; 54 (15.1%) tested positive for influenza (eight A(not subtyped); 30 A(H3); 16 A(H1)pdm09); and 19 (5.3%) tested positive for RSV.
- **Vaccination status:** Of those who tested positive for SARS-CoV-2 with known vaccination status (n=35), 18 (51.4%) had not received a COVID-19 vaccine dose within 180 days prior to their episode of illness. Of those who tested positive for influenza with known vaccination status (n=51), 23 (45.1%) had not received this season's influenza vaccine prior to their episode of illness.
- **ICU admissions**: Among those for whom admission to ICU and/or respiratory status is known (n=363), 194 (53.4%) reported admission to ICU and/or required respiratory support.
- Outcome: Of those discharged, with known outcome (n=272), 30 (11.0%) SARI cases died in hospital.

Table of contents

Table of contents	2
Background	3
Methods	3
Case definition	3
Denominator data	4
Laboratory testing	4
Data collection and reporting	4
The influenza season	4
Reference dates	5
Results	6
SARI cases and incidence rates	6
Demographics	7
Underlying medical conditions and risk factors	8
Symptoms	10
Severe clinical course during hospitalisation	11
Laboratory testing for SARS-CoV-2, Influenza and RSV	12
Outcome	16
Vaccination status	16
Acknowledgements	19
Technical notes	19
Appendix	20
Table A1	20
Table A2	20
Table A3	21

Background

Severe acute respiratory infection (SARI) is of major relevance to public health worldwide. Surveillance of SARI is essential to monitor the (co-) circulation of respiratory pathogens and to assess disease severity. Data collected as part of SARI surveillance can provide important early warning information in the context of respiratory disease outbreaks and pandemics. SARI data can also be used as a platform to measure vaccine and antiviral effectiveness and impact. The objectives of SARI surveillance are:

- To describe the number and incidence of SARI cases by aetiology, time, place and person
- To describe and monitor trends, intensity of activity and severity of SARI infections
- To identify groups at risk of severe disease
- · To detect unusual and unexpected events
- To assess the SARI burden of disease in the participating hospital
- To assess and monitor vaccine effectiveness

Methods

SARI surveillance was implemented in one tertiary care adult hospital; St. Vincent's University Hospital (SVUH), Dublin. Surveillance commenced on the 5th of July 2021. The SARI surveillance system includes people who are aged 15 years or older.

Case definition

SARI cases are identified from new admissions through the Emergency Department, based on clinical symptoms. Patients that develop SARI during their admission, or are admitted through alternate routes, are not included.

Clinical SARI case:

The European Centre for Disease Prevention and Control (ECDC) clinical SARI case definition is used for SARI surveillance in Ireland since week 34 2021:

• ECDC SARI definition: A hospitalised¹ person with acute respiratory infection, with at least one of the following symptoms: cough, fever, shortness of breath OR sudden onset of anosmia, ageusia or dysgeusia with onset of symptoms within 14 days prior to hospital admission.

_

¹ Hospitalised for at least 24 hours

Denominator data

Denominator data for the hospital catchment area are based on the Census of Population, 2022. The hospital catchment data were prepared and provided by the Health Intelligence Unit (HIU) of the Health Service Executive (HSE) and were extracted from Health Atlas Ireland on 15/12/2023.

Weekly denominator data on all-cause hospital admissions, through the Emergency Department, are provided by the SVUH Statistics Department.

Laboratory testing

SARS-CoV-2, influenza, and RSV PCR testing is carried out on admission.

SARI samples that are positive for SARS-CoV-2 and that have a cycle threshold (Ct) value <25 are referred for whole genome sequencing (WGS). All WGS testing was performed in the NVRL up to week 44 2022. The molecular laboratory in SVUH has been identified as a spoke WGS testing site as part of the national SARS-CoV-2 WGS surveillance programme. From week 45 2022, SARI WGS testing has been performed on-site at SVUH.

Samples that are PCR positive for influenza are sent to the National Virus Reference Laboratory (NVRL) for influenza typing/subtyping/genetic and antigenic characterisation.

Data collection and reporting

Clinical data are collected and managed using REDCap electronic data capture tools hosted at University College Dublin. Laboratory data are extracted from APEX, the laboratory information management system (LIMS), using IBM Cognos software hosted at SVUH.

Case-based data are reported by SVUH to the HSE Health Protection Surveillance Centre (HPSC) on a weekly basis. Data are also reported by HPSC to ECDC via The European Surveillance System (TESSy) on weekly basis as part of European level SARI surveillance.

COVID-19 vaccination data are obtained from the National COVID-19 Vaccination Management System (COVAX) and linked to SARI cases by the HSE-Integrated Information Service (IIS), where data are available.

The influenza season

The influenza surveillance season runs from week 40 (early October) to week 20 (end of May). During this time, seasonal viruses usually circulate at higher levels, compared to the summer period (weeks 21 to 39). The seasonal comparisons used in this report refer to the influenza surveillance season.

Reference dates

```
05/07/2021 (Week 27 2021) - commencement of SARI surveillance project
27/09/2021 (Week 39 2021) - rollout of the first COVID-19 booster vaccination campaign
22/04/2022 (Week 16 2022) - rollout of the second COVID-19 booster vaccination campaign
03/10/2022 (Week 40 2022) - rollout of the third COVID-19 booster vaccination campaign
28/04/2023 (Week 17 2023) - rollout of the fourth COVID-19 booster vaccination campaign
02/10/2023 (Week 40 2023) - rollout of the fifth COVID-19 booster vaccination campaign
04/10/2021 (Week 40 2021) - start of the 2021/2022 season
03/10/2022 (Week 40 2022) - start of the 2022/2023 season
02/10/2023 (Week 40 2023) - start of the 2023/2024 season
```

Week number refers to the week of hospital admission. Weeks are from Monday to Sunday, as per the international ISO week².

² Monday to Sunday (ISO week) used as per ECDC/WHO/International reporting protocol.

Results

Data were extracted from the HPSC SARI surveillance database on 20/03/2024. Data are provisional and subject to ongoing review, validation and update. As a result, figures presented in this report may differ from previously published figures.

SARI cases and incidence rates

In total, 369 SARI cases were admitted to St. Vincent's University Hospital (SVUH) during the current season (weeks 40 2023 - 11 2024), 437 SARI cases were admitted during the same period in the 2022/2023 season (weeks 40 2022 - 11 2023).

In week 11 2024:

- 10 SARI cases were reported, a 9.1% decrease compared to 113 SARI cases reported in week 10 2024 (Figure 1)
- The SARI incidence rate was 3.2 per 100,000 hospital catchment population aged 15 years and older, compared to the rate of 3.5 per 100,000 in week 10 2024.
- The incidence rate per emergency hospitalisations was 31.2 per 1,000 emergency admissions, a 12.4% decrease compared to the rate of 35.6 per 1,000 emergency admissions in week 10 2024.




Figure 1: Number and incidence of SARI hospitalised cases (emergency admissions) by week of hospital admission, week 40 2021 to week 11 2024 (n=1778)

³ On review, two cases notified in week 10 2024 and one case notified in week 7 2024, were denotified in week 11 2024

Demographics

In week 11 2024, of the 10 SARI cases reported:

- Males accounted for a higher proportion of SARI cases, n=6 (60.0%) (Table 1)
- Median age of SARI cases admitted was 82 years (interquartile range: 76-86 years)
- Age specific incidence rate amongst those aged 65 years and older was 14.3 per 100,000 compared to 14.3 per 100,000 in week 10 2024.

The incidence rate per 100,000 hospital catchment population by age group is shown in Figure 2.

Table 1: Number and proportion of SARI cases by sex and age, for the current week, last four weeks (weeks 8 - 11 2024), current 2023/2024 season (weeks 40 2023 - 11 2024) and the previous 2022/2023 season (weeks 40 2022 - 11 2023).

Season	Curren	t week	Last fou	r weeks	Current	season	Previous	season
Week/Year	W11	2024	W8 2024- W11 2024		W40 2023- W11 2024		W40 2022- W11 2023	
	n	(%)	n	(%)	n	(%)	n	(%)
All SARI cases	10		59		369		437	
Gender								
Male	6	60.0	33	55.9	188	50.9	206	47.1
Female	4	40.0	26	44.1	181	49.1	231	52.9
Age (years)								
Mean	80		75		72		71	
Median	82		78		77		74	
IQR	76-86		66-84		66-84		63-83	
Range	64-92		41-100		18-100		17-101	
Age groups (years)								
15-24	0	0.0	0	0.0	6	1.6	9	2.1
25-34	0	0.0	0	0.0	11	3.0	10	2.3
35-44	0	0.0	3	5.1	21	5.7	19	4.3
45-54	0	0.0	2	3.4	18	4.9	22	5.0
55-64	1	10.0	7	11.9	26	7.0	63	14.4
65-74	2	20.0	14	23.7	80	21.7	97	22.2
75-84	3	30.0	19	32.2	124	33.6	136	31.1
85+	4	40.0	14	23.7	83	22.5	81	18.5




Figure 2: SARI incidence rate per 100,000 hospital catchment population by age group and week of hospital admission, from week 40 2021 to week 11 2024 (n=1778)

Underlying medical conditions and risk factors

The number and proportion of individuals with underlying medical conditions, where known, among those that reported having underlying medical conditions are displayed in Table 2.

Weekly proportions can be based on small numbers and vary from week to week, caution is therefore advised when interpreting changes in weekly proportions.

Table 2: Number and proportion of SARI cases with underlying medical conditions, reported on hospital admission, for the current week, last four weeks (weeks 8 - 11 2024), current 2023/2024 season (weeks 40 2023 - 11 2024) and the previous 2022/2023 season (weeks 40 2022 - 11 2023).

Period	Curre	nt week		t four eeks	Current season		Previous season	
Weeks	W11	2024	W8 2024- W11 2024		W40 2023- W11 2024		W40 2022- W11 2023	
Medical conditions*	n	%	n	n %		%	n	%
Total cases*	9		53		340		410	
Heart disease	3	33.3	28	52.8	143	42.1	156	38.0
Hypertension	2	22.2	19	35.8	118	34.7	168	41.0
Lung disease	4	44.4	19	35.8	111	32.6	154	37.6
Cancer	2	22.2	9	17.0	66	19.4	66	16.1
Neurological disease	1	11.1	6	11.3	50	14.7	89	21.7
Asthma	0	0.0	5	9.4	43	12.6	66	16.1
Diabetes	2	22.2	10	18.9	59	17.4	67	16.3
Kidney disease	1	11.1	7	13.2	41	12.1	27	6.6
Intellectual disability	0	0.0	0	0.0	4	1.2	15	3.7
Immunocompromised	0	0.0	0	0.0	4	1.2	4	1.0
Obesity	0	0.0	0	0.0	2	0.6	11	2.7
Cystic fibrosis	0	0.0	0	0.0	4	1.2	1	0.2
Asplenia**	0	0.0	0	0.0	0	0.0	-	-
Dementia**	3	33.3	7	13.2	35	10.3	-	-
Down syndrome**	0	0.0	0	0.0	1	0.3	-	-
Long COVID**	0	0.0	0	0.0	0	0.0	-	-
Tuberculosis**	0	0.0	0	0.0	0	0.0	-	-
Rheumatological disease**	2	22.2	2	3.8	4	1.2	-	-
Other chronic conditions***	5	55.6	24	45.3	134	39.4	205	50.0

^{*}SARI cases could be reported with one or more underlying medical conditions, only cases where underlying medical conditions are reported are included.

Among female SARI cases aged 15-49 years admitted during the 2023/2024 season (weeks 40 2023 - 11 2024), 1 (3.6%) case was reported as being pregnant at the time of admission. During the same period in the 2022/2023 season (weeks 40 2022 - 11 2023), 3 (12.0%) SARI cases were reported as being pregnant at the time of admission.

^{**}Data collection for these underlying medical conditions began in week 49 2023.

^{***}Data reported on other chronic conditions may include some of the chronic conditions listed above, these data are under review and may change over time

Among those admitted during the 2023/2024 season for whom healthcare worker status is known, 6 (1.6%) cases were reported as being healthcare workers at the time of admission. During the same period in the 2022/2023 season, 4 (0.9%) SARI cases were reported as being healthcare workers.

Symptoms

Information on clinical symptoms, either at or prior to hospital admission, was reported for all SARI cases. The most common symptoms reported were cough and shortness of breath (Table 3).

Table 3: Number and proportion of SARI cases with clinical symptoms, either at or prior to hospital admission, for the current week, last four weeks (weeks 8 - 11 2024), current 2023/2024 season (weeks 40 2023 - 11 2024) and the previous 2022/2023 season (weeks 40 2022 - 11 2023).

Period	Current week			Last four weeks		season	Previous season		
Weeks	W11	W11 2024		W8 2024- W11 2024		W40 2023- W11 2024		W40 2022- W11 2023	
Clinical symptoms*	n %		n	n %		n %		%	
Total cases	10		59		369		437		
Cough	5	50.0	44	74.6	266	72.1	355	81.2	
Shortness of breath	4	40.0	40	67.8	265	71.8	326	74.6	
Fever	6	60.0	22	37.3	166	45.0	214	49.0	
General deterioration	6	60.0	26	44.1	123	33.3	202	46.2	
Malaise	1	10.0	5	8.5	62	16.8	34	7.8	
Headache	0	0.0	2	3.4	16	4.3	20	4.6	
Muscular pain	0	0.0	0	0.0	17	4.6	29	6.6	
Sore throat	0	0.0	1	1.7	24	6.5	24	5.5	
Ageusia	0	0.0	0	0.0	1	0.3	1	0.2	
Anosmia	0	0.0	0	0.0	1	0.3	2	0.5	
Dysgeusia	0	0.0	0	0.0	1	0.3	0	0.0	
Sepsis**	0	0.0	6	10.2	19	5.1	-	-	
Acute confusion**	3	30.0	9	15.3	20	5.4	-	-	
Vomiting**	1	10.0	5	8.5	21	5.7	-	-	
Nausea**	0	0.0	1	1.7	7	1.9	-	-	
Diarrhoea**	0	0.0	1	1.7	8	2.2	-	-	
Apnoea**	0	0.0	0	0.0	1	0.3		-	

^{*}SARI cases could be reported with one or more clinical symptoms

^{**}Data collection for these symptoms began in week 49 2023.

Severe clinical course during hospitalisation

Information on the clinical course during hospitalisation is only available after discharge, and there may be a delay between discharge and data collection, due to the manual data collection methods required. Among those for whom discharge information is available the most common complication reported was pneumonia (Table 4).

Information on ICU admission and respiratory support may be available prior to discharge, see Table 5. However, length of stay in ICU is only available after discharge, therefore, data on ICU length of stay for the current season are not included, due to the small numbers involved.

Data collection is ongoing for those not yet discharged from hospital.

Table 4: Number and proportion of SARI cases by complication, for the current 2023/2024 season (weeks 40 2023 - 11 2024), the previous 2022/2023 season (weeks 40 2022 - 11 2023), and cases admitted between week 40 2022 and week 11 2024.

Season(s)	Current	Current season		Previous season		40 2022
Week/Year	W40 2023- W11 2024		W40 2022- W11 2023		W40 : W11	
Complications*	n	%	n	%	n	%
Total discharged cases	272		437		1033	
Pneumonia	23	8.5	30	6.9	121	11.7
ARDS	12	4.4	13	3.0	50	4.8
Sepsis	14	5.1	10	2.3	33	3.2
Multiorgan failure	5	1.8	4	0.9	13	1.3
Myocarditis	1	0.4	0	0.0	1	0.1
Encephalitis	0	0.0	0	0.0	0	0.0
Bronchiolitis	0	0.0	0	0.0	1	0.1
Acute kidney injury**	16	5.9	-	-	17	1.6
Heart failure**	5	1.8	-	-	7	0.7
Secondary bacterial infection**	0	0.0	-	-	0	0.0
Other complications***	58	21.3	119	27.2	250	24.2
No complications	168	61.8	281	64.3	616	59.6

^{*}SARI cases could be reported with one or more complications.

^{**}Data collection for these complications began in week 49 2023.

^{***}Data reported on "other complications" may include some of the complications listed above, these data are under review and may change over time.

Table 5: Number and proportion of SARI cases by respiratory support and ICU admission, for the current 2023/2024 season (weeks 40 2023 - 11 2024), the previous 2022/2023 season (weeks 40 2022 - 11 2023), and cases admitted between week 40 2022 and week 11 2024.

Season(s)	Current season		Previous season		Since W40 202	
Week/Year	W40 2023- W11 2024		W40 2022- W11 2023		W40 2022- W11 2024	
	n	%	n	%	n	%
Respiratory support status known	274		437		1035	
High-flow oxygen therapy*	186	67.9	274	62.7	659	63.7
Invasive ventilation	7	2.6	14	3.2	25	1.4
No respiratory support	81	29.6	149	34.1	351	33.9
ICU status known	363		437		1125	
ICU/ventilated**	194	53.4	288	65.9	686	61.0
Admitted to ICU	9	2.5	26	5.9	49	4.4
Admitted and discharged	9	2.5	26	5.9	49	4.4
ICU length of stay (days)						
Mean	-		9		8	
Median	-		6		5	
Interquartile range	_		3-11		3-10	
Range	-		1-42		<1-42	

^{*}Non-invasive ventilation

Laboratory testing for SARS-CoV-2, Influenza and RSV

PCR testing

SARI cases are tested by PCR for SARS-CoV-2, influenza and RSV on admission.

In week 11 2024:

- SARS-CoV-2 PCR testing was carried out on 10 (100.0%) SARI cases, one (10.0%) tested positive, compared to one (9.1%) SARS-CoV-2 positive cases in week 10 2024.
- Influenza PCR testing was carried out on 10 (100.0%) SARI cases, two (20.0%) tested positive, compared to one (9.1%) influenza positive cases in week 10 2024.
- Respiratory syncytial virus (RSV) PCR testing was carried out on 10 (100.0%) SARI cases, none tested positive for RSV, the last RSV positive case was week 6 2024.

^{**}SARI cases which required invasive and/or non-invasive ventilation and/or ICU admission

The weekly positivity rate of SARI cases for the three acute respiratory pathogens are presented in Figure 3. Table 6 displays the number and proportion of SARI cases tested by PCR and positive for SARS-CoV-2, influenza and RSV, and the type/subtype for all influenza PCR positive test results.




Figure 3: Percentage of SARI cases with a positive laboratory test result for SARS-CoV-2, influenza and RSV by week, from week 40 2021 to week 11 2024

Table 6: Number of positive SARS-CoV-2, influenza, and RSV SARI cases, and influenza type/subtype for the current week, previous two weeks (week 10 2024, week 9 2024), current 2023/2024 season (weeks 40 2023 - 11 2024), and the 2022/2023 season (weeks 40 2022 - 11 2023).

Period		Individual weeks						Current season		Previous season		
Weeks	W11	2024	W10	2024	W9 2024							2022- 2023
Test result	n	%	n	%	n	%	n	%	n	%		
SARS-CoV-2												
Total tested	10		11		14		357		431			
Positive	1	10.0	1	9.1	1	7.1	43	12.0	77	21.6		
RSV												
Total tested	10		11		14		357		427			
Positive	0	0.0	0	0.0	0	0.0	19	5.3	41	11.5		
Influenza												
Total tested	10		11		14		357		427			
Positive	2	20.0	1	9.1	2	14.3	54	15.1	66	18.5		
Influenza A (H3)	0	0.0	0	0.0	2	14.3	30	8.4	31	8.7		
Influenza A (H1)pdm09	0	0.0	0	0.0	0	0.0	16	4.5	29	8.1		
Influenza A (not subtyped)	2	20.0	1	9.1	0	0.0	8	2.2	4	1.1		
Influenza B (Victoria)	0	0.0	0	0.0	0	0.0	0	0.0	2	0.6		
Influenza B (no lineage)	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0		

In the current season (weeks $40\ 2023-11\ 2024$), two cases with co-infections were reported; one case tested positive for SARS CoV-2 and influenza A(H3), and one case tested positive for SARS CoV-2 and RSV.

SARS-CoV-2 Genomic analysis

There can be a lag-time before WGS results are available. The WGS data presented is up to week 8 2024

Sequencing results have been received for 371 SARS-CoV-2 SARI cases admitted between week 40 2021 and week 8 2024 (Figure 4).

BA.2 sub-lineage JN.1 (and its sub-lineages) is the dominant variant circulating among SARI cases admitted to the hospital site in the current season. Among SARS-CoV-2 positive SARI cases admitted during the current season (weeks 40 2023 - 11 2024), for whom WGS data are available, 19 (54.3%) were variant BA.2 sub-lineage JN.1 (and its sub-lineages), 8 (22.9%) were XBB.1.5-like lineages, and 8 (22.9%) were XBB.1.5-like+F456L mutation.

Further information on SARI variants is available in the appendix (Table A1 and A2). For further information on circulating variants in Ireland, see the COVID-19 virus variants reports on the HPSC website⁴.




Figure 4: Number of SARI cases sequenced and reported, by week of hospitalisation, week 40 2021 to week 8 2024 (n=371)

Note: As described by the ECDC, 'XBB.1.5-like' and 'XBB.1.5-like + F456L' refer to groupings of lineages that share sets of spike protein mutations

Page 15 of 21

^{*}Includes sub-lineage JN.1

^{**}All other variants

⁴ https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/surveillance/summaryofcovid-19virusvariantsinireland/

Outcome

Collection of discharge data is a manual process, therefore there is a significant lag time between patient discharge and data collection.

Of the 369 SARI cases admitted to the SARI hospital site during the current 2023/2024 season (weeks 40 2023 - 11 2024), 272 (73.7%) have discharge data available. Of those admitted during the same period in the 2022/2023 season (weeks 40 2022 - 11 2023), 437 (100.0%) cases have been discharged (Table 7).

Among SARI cases admitted during the current 2023/2024 season (weeks 40 2023 - 11 2024) and discharged with known outcome, 30 (11.0%) deaths have been reported, 15 (50.0%) were male and 15 (50.0%) were female. The median age was 82 years (IQR: 71-89 years).

Among SARI cases admitted during the 2022/2023 season (weeks 40 2022 - 11 2023) and discharged with known outcome, 48 (11.0%) died in hospital, 23 (47.9%) were male and 25 (52.1%) were female. The median age was 78.5 years (IQR: 74-87 years).

Table 7: Number and proportion of discharged SARI cases by outcome and hospital length of stay, for the current 2023/2024 season (weeks 40 2023 - 11 2024), the previous 2022/2023 season (weeks 40 2022 - 11 2023), and cases admitted between week 40 2022 and week 11 2024.

Season(s)	Current season		Previous season		Since W40 2022	
Week/Year	W40 2023- W11 2024		W40 2022- W11 2023		W40 2022- W11 2024	
	n %		n	%	n	%
Known outcome	272		437		1033	
Discharged alive	242	89.0	384	87.9	926	89.6
Transferred*	0	0.0	5	1.1	8	0.8
Died in hospital	30	11.0	48	11.0	99	9.6
Hospital length of stay (days)						
Mean	11		14		12	
Median	6		6		6	
Interquartile range	3-15		3-13		3-13	
Range	1-63		1-271		1-271	

^{*}Transferred to another hospital

Vaccination status

Vaccination data are available approximately one week after cases are notified to HPSC, therefore the vaccination status for the current week's SARI cases is recorded as unknown.

COVID-19 Vaccination status

Amongst SARI cases, admitted in the current season (weeks 40 2023 - 11 2024) who were PCR positive for SARS-CoV-2 and with known COVID-19 vaccination status, 18 (51.4%) had not received a vaccine dose within the six months prior to their episode of illness (Table 8). Characteristics of **all** SARI cases by time since last COVID-19 vaccine dose and symptom onset during the current season are presented in the Appendix (Table A3). Please refer to the technical notes for the full list of definitions on COVID-19 vaccination status.

Table 8: Characteristics of SARS-CoV-2 positive SARI cases by time since last COVID-19 vaccine dose and symptom onset during the current season (weeks 40 2023-11 2024). Note: SARS-CoV-2 positive SARI cases with unknown vaccination status, n=8 (18.6%) are excluded.

Weeks	V	Week 40 2023 - 10 20	24
Characteristic	<180 days, N = 17 ¹	>=180 days, N = 18 ¹	Not vaccinated, N = 0 ¹
Total	17 (48.6%)	18 (51.4%)	0 (0.0%)
Age(years)			
Mean	77	73	-
Median	77	75	-
IQR	73 - 82	69 - 84	-
Range	65 - 91	33 - 92	-
Gender			
Female	6 (46.2%)	7 (53.8%)	0 (0.0%)
Male	11 (50.0%)	11 (50.0%)	0 (0.0%)
Age groups (years)			
15-49	0 (0.0%)	1 (100.0%)	0 (0.0%)
50-69	3 (42.9%)	4 (57.1%)	0 (0.0%)
70+	14 (51.9%)	13 (48.1%)	0 (0.0%)
Patient residence			
Residential care facility	5 (100.0%)	0 (0.0%)	0 (0.0%)
Private residence/home	12 (44.4%)	15 (55.6%)	0 (0.0%)
Other residence	0 (0.0%)	0 (0.0%)	0 (0.0%)
Patient residence not known	0 (0.0%)	3 (100.0%)	0 (0.0%)
Underlying medical conditions			
Yes	17 (48.6%)	18 (51.4%)	0 (0.0%)
No	0 (0.0%)	0 (0.0%)	0 (0.0%)
Unknown	0 (0.0%)	0 (0.0%)	0 (0.0%)

¹n (%)

Influenza vaccination status

Amongst the SARI cases, admitted in the current season (weeks 40 2023 - 11 2024) who were PCR positive for influenza and with known vaccination status, 23 (45.1%) had not received a dose of this current season's vaccine prior to their episode of illness (Table 9).

Table 9: Characteristics of influenza positive SARI cases by influenza vaccination status during the current season (weeks 40 2023 - 11 2024). Note: Influenza positive SARI cases with unknown vaccination status (n=3 (5.6%)) are excluded.

Weeks	Week 40 2023 - 11 2024						
Characteristic	Vaccinated, N = 28 ¹	Not vaccinated, N = 23 ¹					
Total	28 (54.9%)	23 (45.1%)					
Age(years)							
Mean	77	70					
Median	81	72					
IQR	73 - 84	60 - 80					
Range	39 - 90	41 - 100					
Gender							
Female	13 (48.1%)	14 (51.9%)					
Male	15 (62.5%)	9 (37.5%)					
Age groups (years)							
15-49	2 (50.0%)	2 (50.0%)					
50-69	2 (20.0%)	8 (80.0%)					
70+	24 (64.9%)	13 (35.1%)					
Patient residence							
Residential care facility	5 (45.5%)	6 (54.5%)					
Private residence/home	23 (57.5%)	17 (42.5%)					
Other residence	0 (0.0%)	0 (0.0%)					
Patient residence not known	0 (0.0%)	0 (0.0%)					
Underlying medical conditions							
Yes	26 (56.5%)	20 (43.5%)					
No	2 (40.0%)	3 (60.0%)					
Unknown	0 (0.0%)	0 (0.0%)					

¹n (%)

Acknowledgements

Sincere thanks are extended to all those who participate in SARI surveillance, including those in St. Vincent's University Hospital, the UCD Clinical Research Centre and the National Virus Reference Laboratory. Thanks to members of the HSE Integrated Information Services (IIS) for work on the SARI-COVAX data linkages.

This report was produced by the SARI surveillance team at HPSC.

This report was produced using R studio software.

Technical notes

- SARI case
 - A SARI case refers to an individual patient episode of care.
- 2. Vaccination status
 - For the purposes of SARI surveillance, vaccination status of cases is as follows:

Vaccinated⁵: A case who received their last primary COVID-19 vaccine dose ≥ 14 days prior to the date of symptom onset or their last booster COVID-19 vaccine dose ≥7 days prior to the date of symptom onset.

Time since vaccination: For vaccinated cases, time since vaccination is calculated by subtracting the date of vaccination from the date of symptom onset and categorised as <180 days or ≥180 days since vaccination.

Not vaccinated, if the following applies:

- Vaccination record on the National COVID-19 Immunisation system indicates the person was vaccinated after the date of symptom onset.
- The SARI patient was reported as not vaccinated on the SARI hospital clinical questionnaire, and there is no identifiable linked record of COVID-19 vaccination on the National COVID-19 Immunisation system.

Vaccine status unknown, if:

- The SARI patient is reported on the SARI hospital clinical questionnaire as vaccinated, however there is no identifiable linked record of COVID-19 vaccination on the National COVID-19 Immunisation system. Vaccination status is reported as unknown, until verified on the National COVID-19 Immunisation system.
- The SARI patient is reported on the SARI hospital clinical questionnaire as vaccination status unknown, AND there is no identifiable linked record of COVID-19 vaccination on the National COVID-19 Immunisation system

.

⁵ Refer to www.hse.ie for further information on the COVID-19 vaccination rollout

Appendix

Table A1

Number and proportion of SARI cases sequenced and reported, by pango lineage and variant, admitted during the 2023/2024 season, weeks 40-11 (n=35)

Virus Variant			Sequenced cases %
BA.2 lineages	JN.1	18	51.4
	JN.1.8.1	1	2.9
XBB.1.5like+F456L	EG.5.1	1	2.9
	EG.5.1.1	1	2.9
	EG.5.1.3	1	2.9
	HK.3	1	2.9
	HV.1	1	2.9
	XBB.1.16.6	2	5.7
XBB.1.5-like lineages	FL.15	1	2.9
	FL.9	1	2.9
	GE.1	2	5.7
	JD.1.1	1	2.9
	XBB.1.5	1	2.9
	XBB.2.3	1	2.9
	XBB.2.3.11	1	2.9
	XBB.2.3.13	1	2.9
Total		35	

Table A2

Number of SARI cases sequenced and reported by pango lineage and week of admission for the previous five weeks for which sequencing data are available.

Virus variant	Pango lineage	2024- W04	2024- W05	2024- W06	2024- W07	2024- W08	Total
BA.2	JN.1	0	0	0	0	1	1
	JN.1.8.1	0	0	0	1	0	1
Total	-	0	0	0	1	1	2

Table A3

Characteristics of **all** SARI cases by COVID-19 vaccination status by time since last COVID-19 vaccine dose and symptom onset during the current influenza season (weeks 40 2023 - 11 2024). Note: SARI cases with unknown vaccination status are excluded, n=84 (22.8%).

Weeks	Week 40 2023 - 10 2024		
Characteristic	<180 days, N = 163 ¹	>=180 days, N = 120¹	Not vaccinated, N = 2 ¹
Total	163 (57.2%)	120 (42.1%)	2 (0.7%)
Age(years)			
Mean	76	69	35
Median	79	73	35
IQR	72 - 85	59 - 82	31 - 38
Range	18 - 100	18 - 94	28 - 41
Gender			
Female	80 (57.1%)	60 (42.9%)	0 (0.0%)
Male	83 (57.2%)	60 (41.4%)	2 (1.4%)
Age groups (years)			
15-49	11 (33.3%)	20 (60.6%)	2 (6.1%)
50-69	22 (41.5%)	31 (58.5%)	0 (0.0%)
70+	130 (65.3%)	69 (34.7%)	0 (0.0%)
Patient residence			
Residential care facility	40 (93.0%)	3 (7.0%)	0 (0.0%)
Private residence/home	115 (51.3%)	107 (47.8%)	2 (0.9%)
Other residence	0 (0.0%)	0 (0.0%)	0 (0.0%)
Patient residence not known	8 (44.4%)	10 (55.6%)	0 (0.0%)
Underlying medical conditions			
Yes	159 (58.7%)	112 (41.3%)	0 (0.0%)
No	4 (28.6%)	8 (57.1%)	2 (14.3%)
Unknown	0 (0.0%)	0 (0.0%)	0 (0.0%)

¹n (%)