

Bird flu

Public health advice on bird flu (avian influenza) for poultry flock owners and poultry workers

Avoid the risk of infection

- Wash hands (and arms if necessary) before eating, drinking, smoking, using the telephone, taking medication, applying make-up, inserting contact lenses
- Cover all new and existing cuts and grazes with waterproof dressings and/or gloves before starting work with poultry. If cuts and grazes occur, wash immediately with soap and running water and apply a waterproof dressing
- Avoid touching your eyes nose and mouth with your hands e.g. don't put pens or pencils in your mouth or wipe your eyes with your hand
- Follow the bio security advice in the leaflet from the Department of Agriculture and Food

Get flu vaccine every year

Get vaccinated against human flu.

The vaccine prevents infection with human flu and can cut the risk of a person becoming sick with human flu and bird flu at the same time. This can prevent the two types of flu mixing and becoming more easily spread from person-to-person.

The vaccine is offered free to poultry flock owners and poultry workers. Your GP can arrange this for you. It does not protect against bird flu.

If you suspect disease in your poultry

- Ask your vet for advice immediately
- Keep contact with the poultry to a minimum until the vet has assessed the situation
- Don't let anyone come into contact with the potentially infected poultry, i.e. keep people away
- Do not move poultry or poultry products e.g. feathers, eggs etc or manure/litter from the premises

For further information visit www.hpsc.ie



