Influenza Surveillance in Ireland - Weekly Report

Influenza Week 41 2016 (10th – 16th October 2016)











Summary

All indicators of influenza activity in Ireland were at low levels during week 41 2016 (week ending 16th October 2016).

- <u>Influenza-like illness (ILI)</u>: The sentinel GP influenza-like illness (ILI) consultation rate was 4.9 per 100,000 population in week 41 2016, remaining low, and stable compared to the updated rate of 4.4 per 100,000 reported during week 40 2016.
 - o ILI rates were below the Irish baseline threshold (18.3 per 100,000 population)
 - o ILI rates were at low levels in all age groups
- <u>GP Out of Hours:</u> The proportion of influenza—related calls to GP Out-of-Hours services was at low levels during week 41 2016.
- National Virus Reference Laboratory (NVRL):
 - One influenza positive specimen, an influenza A(H3) virus, was detected by the NVRL from non-sentinel sources during week 41 2016.
 - o No sentinel GP specimens were positive for influenza during week 41 2016.
 - o Sporadic positive detections of influenza A(H3), A(H1)pdm09 and B were reported throughout the 2016 summer period.
 - Positive detections of respiratory syncytial virus (RSV) are starting to increase, although remain at low levels.
 - Sporadic detections of adenovirus, human metapneumovirus (hMPV) and parainfluenza virus type 4
 (PIV-4) were reported during week 41 2016.
- Respiratory admissions: Respiratory admissions reported from a network of sentinel hospitals remained at low levels during weeks 40 and 41 2016.
- Hospitalisations: No confirmed influenza hospitalised cases were notified to HPSC during week 41 2016.
- <u>Critical care admissions:</u> No confirmed influenza cases were admitted to critical care units and reported to HPSC during week 41 2016.
- Mortality: There were no reports of any influenza-associated deaths during week 41 2016.
- <u>Outbreaks</u>: Two acute respiratory outbreaks in residential care facilities, one in HSE-NW and one in HSE-W were reported to HPSC during week 41 2016.
- <u>International</u>: As is usual for this time of year, influenza activity is low in the European Region.

1. GP sentinel surveillance system - Clinical Data

- During week 41 2016, 11 influenza-like illness (ILI) cases were reported from sentinel GPs, corresponding to an ILI consultation rate of 4.9 per 100,000 population, remaining low, and stable compared to the updated rate of 4.4 per 100,000 reported during week 40 2016. The ILI rate for week 41 2016 is below the Irish baseline ILI threshold (18.3/100,000 population) (figure 1).
- ILI age specific rates were low in all age groups during week 41 2016 (figure 2).
- HPSC in consultation with the European Centre for Disease Prevention and Control (ECDC) has revised
 the Irish baseline ILI threshold for the 2016/2017 influenza season to 18.3 per 100,000 population; this
 threshold indicates the likelihood that influenza is circulating in the community. The Moving Epidemic
 Method (MEM) has been adopted by ECDC to calculate thresholds for GP ILI consultations in a
 standardised approach across Europe.¹
- The baseline ILI threshold, medium (58.7/100,000 population) and high (113.3/100,000 population) intensity ILI thresholds are shown in figure 1.




Figure 1: ILI sentinel GP consultation rates per 100,000 population, baseline ILI threshold, medium and high intensity ILI thresholds¹ and number of positive influenza A and B specimens tested by the NVRL, by influenza week and season. Source: ICGP and NVRL

¹ For further information on the Moving Epidemic Method (MEM) to calculate ILI thresholds: http://www.ncbi.nlm.nih.gov/pubmed/22897919




Figure 2: Age specific sentinel GP ILI consultation rate per 100,000 population by week during the summer of 2016 and the 2016/2017 influenza season to date. *Source: ICGP*.

2. Influenza and Other Respiratory Virus Detections - NVRL

The data reported in this section for the 2016/2017 influenza season refers to sentinel and non-sentinel respiratory specimens routinely tested for influenza, respiratory syncytial virus (RSV), adenovirus, parainfluenza viruses types 1, 2, 3 & 4 (PIV-1, -2, -3 & -4) and human metapneumovirus (hMPV) by the National Virus Reference Laboratory (NVRL) (figures 3 & 4, tables 1 & 2).

- One (1/153; 0.7%) influenza positive specimen, an influenza A(H3) virus, was detected by the NVRL from non-sentinel sources during week 41 2016. No sentinel GP specimens were positive for influenza during this period.
- Data from the NVRL for week 41 2016 and the 2016/2017 season to date are detailed in tables 1 and 2.
- Fourteen (14/153; 9.2%) respiratory syncytial virus (RSV) positive non-sentinel specimens were reported during week 41 2016. Figure 4 shows the number and percentage of non-sentinel RSV positive specimens detected by the NVRL during the 2016/2017 season, compared to the 2015/2016 season. For the 2016/2017 season to date, no RSV positive specimens from sentinel GP sources have been detected.
- Three adenovirus, three human metapneumovirus (hMPV) and one parainfluenza virus (PIV) type -4 positive specimens were reported by the NVRL during week 41 2016 (table 2).
- The overall proportion (22/153; 14.4%) of non-sentinel specimens positive for respiratory viruses*, increased compared to the previous week (9.2%). * Respiratory viruses tested for by the NVRL are detailed above.




Figure 3: Number of sentinel and non-sentinel specimens tested by the NVRL for influenza and percentage influenza positive by week for the 2016/2017 influenza season. *Source: NVRL*




Figure 4: Number and percentage of non-sentinel RSV positive specimens detected by the NVRL during the 2016/2017 season, compared to the 2015/2016 season. *Source: NVRL*.

Table 1: Number of sentinel and non-sentinel respiratory specimens tested by the NVRL and positive influenza results, for week 41 2016 and the 2016/2017 season to date. Source: NVRL

Week	Specimen type	Total tested	Number influenza	% Influenza		Influenza			
			positive	positive	A (H1)pdm09	A (H3)	A (not subtyped)	Total influenza A	В
41 2016	Sentinel	4	0	0.0	0	0	0	0 0	
	Non-sentinel	153	1	0.7	0	1	0	1	0
	Total	157	1	0.6	0	1	0	1	0
2016/2017	Sentinel	9	0	0.0	0	0	0	0	0
	Non-sentinel	370	2	0.5	0	1	1	2	0
	Total	379	2	0.5	0	1	1	2	0

Table 2: Number of non-sentinel specimens tested by the NVRL for other respiratory viruses and positive results, for week 41 2016 and the 2016/2017 season to date. Source: NVRL

Week	Specimen type	Total tested	RSV	% RSV	Adenovirus	% Adenovirus	PIV- 1	% PIV- 1	PIV- 2	% PIV- 2	PIV- 3	% PIV- 3	PIV- 4	% PIV- 4	hMPV	% hMPV
41 2016	Sentinel	4	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Non-sentinel	153	14	9.2	3	2.0	0	0.0	0	0.0	0	0.0	1	0.7	3	2.0
	Total	157	14	8.9	3	1.9	0	0.0	0	0.0	0	0.0	1	0.6	3	1.9
2016/2017	Sentinel	9	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Non-sentinel	370	28	7.6	5	1.4	0	0.0	0	0.0	2	0.5	0	0.0	3	0.8
	Total	379	28	7.4	5	1.3	0	0.0	0	0.0	2	0.5	0	0.0	3	0.8

[†] Please note that non-sentinel specimens relate to specimens referred to the NVRL (other than sentinel specimens) and may include more than one specimen from each case.

3. Regional Influenza Activity by HSE-Area

The geographical spread of influenza activity is reviewed on a weekly basis using sentinel GP ILI consultation rates, laboratory data and outbreak data.

The geographical spread of influenza/ILI during week 41 2016 is shown in figure 6. Sporadic influenza activity (based on ILI cases only) was reported in HSE-E and -MW and no influenza activity was reported in all other HSE-Areas during week 41 2016 (figure 5).




Figure 5: Map of provisional influenza activity by HSE-Area during influenza week 41 2016

Sentinel hospitals

The Departments of Public Health have established at least one sentinel hospital in each HSE-Area, to report data on total, emergency and respiratory admissions on a weekly basis. For the 2016/2017 influenza season, eight sentinel hospitals are regularly reporting respiratory admissions data in a timely manner.

Respiratory admissions reported from a network of sentinel hospitals were at low levels during weeks 40 and 41 2016, at 257 and 249 respectively (figure 6). It should be noted that seven of eight sentinel hospitals reported data during week 40 2016 and six of eight sentinel hospitals reported during week 41 2016.




Figure 6: Number of respiratory admissions reported from sentinel hospitals and ILI sentinel GP consultation rate per 100,000 population by week and season. Source: Departments of Public Health - Sentinel Hospitals & ICGP. Data were missing from one sentinel hospital during week 40 2016 and two sentinel hospitals during week 41 2016, these weeks are represented by the hatched bars.

4. GP Out-Of-Hours services surveillance

The Department of Public Health in HSE-NE is collating national data on calls to nine of thirteen GP Out-of-Hours services in Ireland. Records with clinical symptoms reported as flu or influenza are extracted for analysis. This information may act as an early indicator of increased ILI activity. However, data are self-reported by callers and are not based on coded influenza diagnoses.

The proportion of influenza—related calls to GP Out-of-Hours services remained at low levels during week 41 2016 at 1.8%, compared to 2.0% during week 40 2016. A slight increase in the proportion of influenza related calls to GP Out-of-Hours services occurred between weeks 37-39 2016; this increase is usually observed each September when schools return from the summer break (figure 7).




Figure 7: Self-reported influenza-related calls as a proportion of total calls to Out-of-Hours GP Co-ops and sentinel GP ILI consultation rate per 100,000 population by week and season. Source: GP Out-Of-Hours services in Ireland (collated by HSE-NE) & ICGP.

5. Influenza & RSV notifications

Influenza and RSV cases notifications are reported on Ireland's Computerised Infectious Disease Reporting System (CIDR), including all positive influenza /RSV specimens reported from all laboratories testing for influenza/RSV and reporting to CIDR.

Influenza and RSV notifications are reported in the <u>Weekly Infectious Disease Report for Ireland</u>. Influenza notifications were at low levels during week 41 2016, with only two cases notified. RSV notifications are starting to increase, with 19 cases notified during week 41 2016, compared to seven during the previous week.

6. Influenza Hospitalisations

No confirmed influenza hospitalised cases were notified to HPSC during week 41 2016.

7. Critical Care Surveillance

The Intensive Care Society of Ireland (ICSI) are continuing with the enhanced surveillance system set up during the 2009 pandemic, on all critical care patients with confirmed influenza. HPSC process and report on this information on behalf of the regional Directors of Public Health/Medical Officers of Health.

No confirmed influenza cases were admitted to critical care and reported to HPSC during week 41 2016.

8. Mortality Surveillance

Influenza-associated deaths include all deaths where influenza is reported as the primary/main cause of death by the physician or if influenza is listed anywhere on the death certificate as the cause of death. HPSC receives daily mortality data from the General Register Office (GRO) on all deaths from all causes registered in Ireland. These data have been used to monitor excess all-cause and influenza and pneumonia deaths as part of the influenza surveillance system and the European Mortality Monitoring Project. These data are provisional due to the time delay in deaths' registration in Ireland. http://www.euromomo.eu/

- There were no reports of any influenza-associated deaths occurring during week 41 2016.
- During week 41 2016, no excess all-cause mortality was reported in Ireland after correcting GRO data for reporting delays with the standardised EuroMOMO algorithm.

9. Outbreak Surveillance

Two acute respiratory outbreaks in residential care facilities were reported to HPSC during week 41 2016, one in HSE-NW and one in HSE-W. No pathogens were identified from these outbreaks.

10. International Summary

As is usual for this time of year, influenza activity is low in the European Region. As of October 17th 2016, globally, influenza activity decreased in Oceania, South Africa and temperate South America. Influenza activity in the temperate zone of the northern hemisphere remained at inter-seasonal levels. See <u>ECDC</u> and WHO influenza surveillance reports for further information.

Further information is available on the following websites:

Northern Ireland http://www.fluawareni.info/
Europe – ECDC http://ecdc.europa.eu/

Public Health England http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/SeasonalInfluenza/

United States CDC http://www.cdc.gov/flu/weekly/fluactivitysurv.htm
Public Health Agency of Canada http://www.phac-aspc.gc.ca/fluwatch/index-eng.php

- Information on Middle Eastern Respiratory Syndrome Coronavirus (MERS), including the latest ECDC rapid risk assessment is available on the <u>ECDC website</u>. Further information and guidance documents are also available on the <u>HPSC</u> and <u>WHO</u> websites.
- Further information on avian influenza is available on the ECDC website.

11. WHO recommendations on the composition of influenza virus vaccines

On February 25, 2016, the WHO vaccine strain selection committee recommended that trivalent vaccines for use in the 2016/2017 influenza season (northern hemisphere winter) contain the following: an A/California/7/2009 (H1N1)pdm09-like virus; an A/Hong Kong/4801/2014 (H3N2)-like virus; a B/Brisbane/60/2008-like virus. http://www.who.int/influenza/vaccines/virus/recommendations/en/

Further information on influenza in Ireland is available at www.hpsc.ie

Acknowledgements

This report was prepared by Lisa Domegan and Joan O'Donnell, HPSC. HPSC wishes to thank the sentinel GPs, the ICGP, NVRL, Departments of Public Health, ICSI and HSE-NE for providing data for this report.