



Enhanced Surveillance of *Clostridium difficile* Infection: Ireland – Q1 2017 National Report

Executive Summary

- During Q1 2017, a total of 411 cases of *C. difficile* infection (CDI) were reported to enhanced surveillance from 54 acute public and private hospitals across Ireland. Therefore, 82% of all CDI cases notified to the Departments of Public Health also have enhanced surveillance data
- The national overall rate of CDI in hospitalised patients was 3.0 cases per 10,000 bed days used (BDU) [298 cases], which lower than that reported for Q1 2016 [326 cases; rate = 3.3]
- There were 200 cases of CDI deemed to be hospital-acquired (HA-CDI), of which 176 were new, representing a national new HA-CDI rate of 1.8 [median rate = 1.3]
- All hospitals reported using a *C. difficile* testing method recommended in the 2014 updated national clinical guidelines for *C. difficile*
- Ribotyping data was available for just 15% of cases, with ribotypes 078, 002 and 014 the most frequently reported
- With regard to acquisition, *C. difficile* was mostly associated with acute hospitals (200; 49%). However, a large proportion of cases were associated with long term care facilities (31; 8%) and the community (84; 20%), whereby patients had no overnight stay in a healthcare facility in 12 weeks prior to symptom onset
- Over one third of all cases (142; 35%) had CDI symptom onset while residing in the community:
 - This emphasises the importance of considering CDI when evaluating any patient with potentially infectious diarrhoea in all healthcare settings, including hospitals, primary care and long-term care facilities. Guidance on CDI for primary and long-term care settings is available at the following link:
<http://www.hpsc.ie/A-Z/Gastroenteric/Clostridiumdifficile/Guidelines/File,14387,en.pdf>
 - It also emphasises the importance for all microbiology laboratories in Ireland to implement the recommendations of the national *C. difficile* clinical guidelines to routinely include *C. difficile* testing for all faeces specimens that take the shape of the container submitted from patients aged ≥2 years, regardless of patient location or physician request. Guidance on *C. difficile* testing is available in Section 2.5, pages 43 – 54 of the national *C. difficile* clinical guidelines
- The excellent participation in enhanced surveillance since it was launched in 2009 indicates the commitment of the microbiology laboratories, multi-disciplinary infection prevention and control and antimicrobial stewardship teams, along with hospital management to understanding the epidemiology of this important infection and minimising the risk of patients acquiring CDI as an unintended consequence of healthcare

Part 1: National CDI Epidemiology – Q1 2017

CDI data was reported to the enhanced surveillance programme from 54 acute public and private hospitals across Ireland ([Appendix A](#)). There were 411 reported CDI cases in patients aged ≥ 2 years, representing a national CDI rate in hospitalised patients of 3.0 cases per 10,000 bed days used (BDU) [298 cases], which is lower than that reported for Q1 2016 [326 cases; rate = 3.3]. The majority were aged ≥ 65 years (66%) and were female (55%). Five cases of severe CDI were reported (1.2%), defined as requiring critical care admission or colectomy due to complications of CDI, versus nine cases (2.0%) for Q1 2016. [Table 1](#) displays the breakdown of all CDI cases for Q1 2017 versus Q1 2016, by case type, origin, onset and severity.

CDI Case Type

The majority were categorised as new infections (83%), with 8% recurrent and for 9%, the CDI case type was unknown.

CDI Origin

The majority were categorised as healthcare-associated (HCA) CDI [n=245; 60%], with community-associated (CA) CDI accounting for 20% [n=84]. For the remainder, the origin was either unknown [n=51; 12%] or could not be determined [n=31; 8%]. Of 245 HCA-CDI cases, the origin was the reporting hospital, termed hospital-acquired (HA) for 200 (82%), a long-term care facility for 31 (13%) and 'other' or 'unknown healthcare facility' for 14 (6%).

CDI Onset


Patient locations at onset of CDI symptoms included; while admitted to a healthcare facility, termed healthcare-onset (HO) for 234 cases (57%), while residing in the community, termed community onset (CO) for 142 cases (35%) and unknown patient location for 35 cases (9%). Of 234 HO CDI cases, the reporting hospital was the onset location for 188 (80%), a long-term care facility for 35 (15%) and other healthcare facilities for 10 (4%).

Table 1. National CDI epidemiology: Q1 2017 versus Q1 2016.

	National CDI Epidemiology Q1 2017 vs Q1 2016	Q1 2017	Q1 2016
CDI case type	Total reported cases:	411	462
	New	342	368
	Recurrent	32	51
	Unknown	37	43
CDI origin	Healthcare-associated (HCA):	245	282
	Reporting hospital	200	207
	Long term care facility (LTCF)	31	55
	Other healthcare facility	10	17
	Unknown healthcare facility	4	3
	Community associated (CA)	84	102
	Discharged within 4 – 12 weeks from healthcare facility	31	27
Unknown origin	51	51	
CDI onset	Healthcare onset (HO):	234	273
	Reporting hospital	188	187
	LTCF	35	62
	Other healthcare facility	10	9
	Community onset (CO)	142	164
Unknown onset location	35	25	
CDI severity	Critical care admission or colectomy	5	9

Part 2: Hospital-acquired CDI (HA-CDI) Epidemiology – Q1 2017

Data on HA-CDI was reported from 54 acute public and private hospitals across Ireland. There were 200 HA-CDI cases in patients aged ≥ 2 years during Q1 2017. Of those, 176 were new HA-CDI cases, representing a national HA-CDI rate of 1.8 [median rate = 1.3], comparable to that reported for Q1 2016 [172 cases; rate = 1.8; median rate = 0.9]. [Figure 1](#) displays quarterly HA-CDI rates since 2010 and [Table 2](#) displays quarterly HA-CDI data from 2015 to 2017.



The national overall CDI rate represents all CDI diagnosed in hospitalised patients per 10,000 BDU, while the HA-CDI rate represents **new** cases of hospital-acquired CDI per 10,000 BDUs. Raw data for this graph is provided in Table 2. The national range is represented by the 5th to 95th percentile of the CDI rate.

Figure 1. Quarterly national HA-CDI rates: 2010 – 2017.

CDI Case Type

The majority of 200 hospital-acquired CDI cases were categorised as new infections (176; 88%), with 20 (10%) recurrent cases. For four CDI cases, the case type was unknown.

CDI Onset

Patient locations at onset of HA-CDI symptoms included; while admitted to a healthcare facility, termed healthcare-onset (HO) for 176 cases (88%) and while residing in the community, termed community onset (CO) for 24 cases (12%).

Of 176 HO CDI cases, the reporting hospital was the onset location for 175 (99%), and was not reported for one case (1%).

Table 2. Quarterly HA-CDI data: 2015 – 2017

YearQ	Number of participating hospitals ^a	Number of cases reported				CDI rate per 10,000 BDUs ^b		
		New	Recurrent	Unknown	Total	Rate	Range ^c	Median
2015Q2	52	215	21	0	236	2.4	0 - 4.5	1.7
2015Q3	52	203	20	0	223	2.3	0 - 4.9	1.8
2015Q4	53	198	27	1	226	2.1	0 - 4.4	1.3
2016Q1	52	172	33	1	206	1.8	0 - 3.2	0.9
2016Q2	51	186	27	1	214	2	0 - 3.2	1.2
2016Q3	52	182	22	2	206	2	0 - 3.8	1.4
2016Q4	51	182	19	2	203	2	0 - 3.3	1.2
2017Q1	54	176	20	4	200	1.8	0 - 3.6	1.3

a Since Q1 2012 97% of all tertiary and general hospitals participated in the enhanced surveillance system.

b The CDI rate is the number of new cases of CDI that were acquired in the reporting hospital - per 10,000 bed days used (BDUs). **c** The national range corresponds to the 5th to 95th percentile of the data.

Data for Q1 2017 is provisional

Part 3: *C. difficile* Testing Methods – Q1 2017

All 54 hospitals participating in the enhanced CDI surveillance system during Q1 2017 reported use of a *C. difficile* testing method recommended by the updated National Clinical Guidelines for Surveillance, Diagnosis & Management of *C. difficile* infection in Ireland (2014). This includes either one of a variety of two-step testing methods (n = 30; 56%) or a single-step method using molecular polymerase chain reaction (PCR) test for *C. difficile* toxin gene (n = 24; 44%) as displayed in [Table 3](#), along with stratification by hospital type.

Table 3. *C. difficile* testing methods utilised in Q1 2017, by hospital type.

Test Category	Hospital Type				Total
	General	Private	Specialist	Tertiary	
1 STEP: PCR for toxin gene	11	2	5	6	24
2 STEP: GDH EIA, followed by confirmatory <i>C. difficile</i> toxin EIA	3	5	2	0	10
2 STEP: Combined GDH with toxin EIA, followed by toxin EIA*	1	0	0	0	1
3 STEP: Combined GDH with toxin EIA, followed by PCR**	5	3	1	1	10
2 STEP: GDH EIA, followed by confirmatory PCR	6	0	0	0	6
2 STEP: PCR, followed by confirmatory toxin EIA	0	1	0	2	3
Total	26	11	8	9	54

PCR for *C. difficile* toxin gene: Polymerase chain reaction (PCR) for the detection of TcdA and/or TcdB genes

GDH EIA Enzyme immunoassay (EIA) for the detection of glutamate dehydrogenase (GDH) of *C. difficile*

GDH AND TOXIN EIA: Enzyme immunoassay (EIA) for the detection of both *C. difficile* GDH and *C. difficile* toxin TcdA and/or TcdB

***2 STEP: Combined GDH with toxin EIA, followed by confirmatory toxin EIA:** Addition of a confirmatory toxin EIA test (using a different EIA kit) if the initial toxin EIA is negative

****2 STEP: Combined GDH with toxin EIA, followed by confirmatory PCR:** Addition of confirmatory PCR if the initial toxin EIA is negative

Part 4: *C. difficile* Ribotyping – Q1 2017

Ribotyping data was available for just 15% of CDI cases reported to CDI enhanced surveillance, a reflection on the continued absence of a national funded *C. difficile* reference laboratory service, which has been a key recommendation of national *C. difficile* guidelines since 2008. Ribotypes 078, 002 and 014 were the most frequently reported. The lack of a robust, prospective system to capture *C. difficile* ribotyping data limits understanding of the epidemiology of this important healthcare-associated infection.

Acknowledgments

The HPSC would like to sincerely thank all who have contributed to this report: Microbiology Surveillance Scientists, Infection Prevention and Control Nurses, Microbiology Laboratory Scientists, Clinical Microbiologists, along with all the staff of the Departments of Public Health across Ireland.

Appendix A: National CDI Enhanced Surveillance Participating Hospitals

Hospital Group	Hospital Name	Category
Dublin Midlands	Adelaide & Meath & National Children's Hospital, Tallaght	Tertiary
	Coombe Women and Infant's University Hospital	Specialist
	Midland Regional Hospital Portlaoise	General
	Midland Regional Hospital Tullamore	General
	Naas General Hospital	General
	St James's Hospital	Tertiary
Ireland East Hospital Group	St Luke's Hospital, Dublin	Specialist
	Cappagh National Orthopaedic Hospital, Dublin	Specialist
	Mater Misericordiae University Hospital	Tertiary
	Midland Regional Hospital Mullingar	General
	National Maternity Hospital, Holles Street	Specialist
	Our Lady's Hospital, Navan	General
	Royal Victoria Eye & Ear Hospital, Dublin	Specialist
	St Columcille's Hospital, Loughlinstown	General
	St Luke's General Hospital, Kilkenny	General
St Michael's Hospital, Dun Laoghaire	General	
RCSI Hospital Group	St Vincent's University Hospital	Tertiary
	Wexford General Hospital	General
	Beaumont Hospital	Tertiary
	Cavan General Hospital	General
	Connolly Hospital, Blanchardstown	General
Saolta Hospital Group	Louth County Hospital, Dundalk	General
	Our Lady of Lourdes Hospital, Drogheda	General
	Letterkenny General Hospital	General
	Mayo General Hospital, Castlebar	General
	Portiuncula University Hospital, Ballinasloe	General
	Roscommon University Hospital	General
South/South West Hospital Group	Sligo General Hospital	General
	University College Hospital Galway	Tertiary
	Bantry General Hospital	General
	Cork University Hospital Group	Tertiary
	Kerry General Hospital, Tralee	General
	Lourdes Orthopaedic Hospital, Kilcreene, Kilkenny	Specialist
	Mallow General Hospital	General
	Mercy University Hospital, Cork	General
South Infirmary - Victoria University Hospital, Cork	General	
UL Hospital Group	South Tipperary General Hospital, Clonmel	General
	Waterford Regional Hospital	Tertiary
	Croom Hospital	Specialist
	Ennis Hospital	General
	Nenagh Hospital	General
	St John's Hospital	General
Private Hospitals	University Hospital, Limerick	Tertiary
	University Maternity Hospital	Specialist
	Aut Even, Kilkenny	Private
	Beacon Hospital, Dublin	Private
	Blackrock Clinic	Private
	Bon Secours, Cork	Private
	Bon Secours, Galway	Private
	Bon Secours, Glasnevin	Private
	Bon Secours, Tralee	Private
	Galway Clinic	Private
	Mater Private, Dublin	Private
	Mater Private, Cork	Private
Children's Hospital Group	St Vincents Private Hospital	Private
	Children's University Hospital, Temple Street	Specialist

Appendix B

Case Definitions for Surveillance of *Clostridium difficile* Infection

For surveillance purposes, a confirmed *Clostridium difficile* infection (CDI) case is a patient two years or older, to whom one or more of the following criteria applies:

- Diarrhoeal* stools or toxic megacolon, with either a positive laboratory assay for *C. difficile* toxin A (TcdA) and/or toxin B (TcdB) in stools or a toxin-producing *C. difficile* organism detected in stool via culture or other means.
- Pseudomembranous colitis (PMC) revealed by lower gastrointestinal endoscopy.
- Colonic histopathology characteristic of *C. difficile* infection (with or without diarrhoea) on a specimen obtained during endoscopy, colectomy or autopsy.

* Diarrhoea is defined as three or more loose/watery bowel movements (which are unusual or different for the patient) in a 24 hour period

CASE TYPE

- **New Case of CDI:**
 - The first episode of CDI, **OR**
 - A subsequent episode of CDI with onset of symptoms **more than eight weeks** after the onset of a previous episode.
- **Recurrent Case of CDI:**
 - A patient with an episode of CDI that occurs **within eight weeks** following the onset of a previous episode **provided that CDI symptoms from the earlier episode resolved with or without therapy.**

ONSET

- **Healthcare onset** » Symptoms start during a stay in a healthcare facility.
- **Community onset** » Symptoms start in a community setting, outside healthcare facilities.
- **No information available** » If no information was available on onset of symptoms

ORIGIN

- **Healthcare-associated case.** This is a CDI patient with either:
 - Onset of symptoms at least 48 hours following admission to a healthcare facility (healthcare-onset, healthcare-associated), **OR**
 - With onset of symptoms in the community within four weeks following discharge from a healthcare facility (community-onset, healthcare-associated).
- **Community-associated case.** This is a CDI patient with either:
 - Onset of symptoms while outside a healthcare facility, and without discharge from a healthcare facility within the previous 12 weeks (community-onset, community-associated), **OR**
 - With onset of symptoms within 48 hours following admission to a healthcare facility without residence in a healthcare facility within the previous 12 weeks (healthcare-onset, community-associated).
- **Discharged 4 – 12 weeks from a healthcare facility**
 - » This is a CDI patient who was discharged from a healthcare facility between four and 12 weeks before the onset of symptoms.
 - **No information available**

SEVERE CDI Case

This is a CDI patient to whom any of the following criteria apply:

- Admission to an intensive care unit for treatment of CDI or its complications (e.g., for shock requiring vasopressor therapy)
- Surgery (colectomy) for toxic megacolon, perforation or refractory colitis
- Death within 30 days after diagnosis if CDI is either the primary or a contributive cause