



ROYAL COLLEGE OF
PHYSICIANS OF IRELAND



Office of the
Nursing & Midwifery
Services Director

Point Prevalence Survey of Healthcare-Associated Infections & Antimicrobial Use in Long-Term Care Facilities (HALT): May 2013

A Summary of Mixed Care LTCF Data

REPUBLIC OF IRELAND: NATIONAL REPORT – MARCH 2014

Report Authors: Fiona Roche, Sheila Donlon & Karen Burns, HPSC

SUGGESTED CITATION: HEALTH PROTECTION SURVEILLANCE CENTRE, POINT PREVALENCE SURVEY OF HEALTHCARE-ASSOCIATED INFECTIONS & ANTIMICROBIAL USE IN LONG-TERM CARE FACILITIES: MAY 2013 – REPUBLIC OF IRELAND NATIONAL REPORT: MARCH 2014

PLEASE NOTE:

This report represents a shortened version of the full national HALT report focusing on data from Mixed Care Facilities.

The full national report including methodology, results on all LTCF care types, discussion and future priorities can be found on the hpsc website [here](#).

Table of Contents

| | |
|---|-----------|
| Executive Summary | 4 |
| 1. RESULTS | 8 |
| 1.1 NATIONAL OVERVIEW | 8 |
| 1.1.1 Description of Participating LTCF | 8 |
| 1.1.2 Governance Structures | 11 |
| 1.1.3 HCAI and Antimicrobial Use | 27 |
| 1.2 MIXED CARE FACILITIES WITH A LOS GREATER THAN 12 MONTHS (MIXED > 12M) | 36 |
| 1.2.1 Description of Care Type | 36 |
| 1.2.2 Description of Residents | 36 |
| 1.2.3 HCAI in Mixed > 12m | 37 |
| 1.2.4 Antimicrobial Use in Mixed > 12m | 41 |
| 2. APPENDICES | 46 |
| 2.1 APPENDIX A: LIST OF ACRONYMS USED IN THIS REPORT | 46 |

Executive Summary

In May 2013, 9,318 residents in 190 Irish long-term care facilities (LTCF) were included in a European point prevalence survey (PPS) of healthcare-associated infections (HCAI) and antimicrobial use. The survey is also known as the HALT survey. This is the national report for Ireland.

Aims of the HALT survey

1. To calculate the prevalence of healthcare-associated infections (HCAI) in residents of Irish LTCF
2. To calculate the prevalence of and indications for antimicrobial use in Irish LTCF
3. To provide the Irish Government, Department of Health, Health Service Executive, the managers, doctors and nurses caring for residents in all of the LTCF that took part, with information for action: to reduce the numbers of residents who develop HCAI and to influence positive antimicrobial stewardship practices in LTCF
4. To provide residents, their families and members of the public with more information about HCAI in Ireland and which types of infections are most commonly seen in Irish LTCF

Participating LTCF

- Of the 190 LTCF, the majority were owned by the Health Service Executive (HSE) [n=128; 67%], followed by private [n=39; 21%] and voluntary services [n=23; 12%]
- The median capacity of participating LTCF was 46 beds (range = 5 – 203) and the median bed occupancy on the HALT survey date was 94%
- Overall, single room accommodation accounted for a median of 34% of available beds. The proportion of single room accommodation was much lower in HSE-owned than privately-owned LTCF (21% versus 76%)
- For the purposes of data analysis and reporting, the HALT steering group stratified the 190 LTCF into the following care type categories, based on the characteristics and estimated length-of-stay (LOS) for the majority of the residents:
 - i. **General nursing homes >12 months (GN>12m):** 103 long-stay facilities with 5,807 residents
 - ii. **Mixed care type facilities >12 months (Mixed>12m):** 26 long-stay facilities with 1,409 residents

- iii. **LTCF caring for residents with intellectual disabilities (Intellectually disabled):** 24 facilities with 1,060 residents
- iv. **LTCF (either general nursing homes or mixed care type facilities) <12 months (LTCF<12m):** 15 short-stay facilities with 374 residents
- v. **LTCF caring for residents with psychiatric conditions (Psychiatric):** 11 facilities with 345 residents
- vi. **Other care types:** Facilities caring for residents with palliative care needs (4 facilities with 89 residents) rehabilitation needs (3 facilities with 139 residents), physical disabilities (2 facilities with 46 residents) and 'other' care types (2 facilities with 49 residents)

Coordination of medical care, infection prevention & control & antimicrobial stewardship

- Overall, resident medical care was provided by the resident's own general practitioner (GP) in 35%, by a directly-employed doctor in 41% and by a mix of GP plus directly-employed doctor care in 24% of LTCF. However, when LTCF were stratified by ownership, GP-led medical care predominated in privately-owned LTCF (82%) versus HSE-owned (35%) LTCF
- A designated coordinating physician, with responsibility for coordination and standardisation of policies/practices for resident medical care within the LTCF was available for only 45% of LTCF overall and for only 26% of privately-owned LTCF. Where a coordinating physician was in post, the reported roles undertaken infrequently included development of local infection prevention and control (IPC) (16%) or antimicrobial prescribing policies (14%)
- One third of LTCF reported having no active local infection prevention and control committee (IPCC)
- Access to a staff member with training in IPC was reported by 62% of LTCF overall and by only 10% of privately-owned LTCF. For the vast majority of LTCF with a trained IPC staff member, that person was an infection prevention and control nurse (IPCN) (93%). For the majority of LTCF, the IPCN was not based in the LTCF on a day-to-day basis (71%)
- Following the HALT survey, additional information was sought to estimate the national number of whole time equivalent (WTE) IPCNs for LTCF: In 2013, it is estimated that there was one WTE IPCN for every 496 LTCF beds in Ireland
- Although the vast majority of LTCF reported the presence of a written local hand hygiene policy (97%), the provision of regular staff hand hygiene training sessions was not universal, with only 88% of LTCF reporting that such a session had been arranged in the past 12

months. Medical and allied health professional staff were less likely to be invited to attend such training sessions than nursing and hygiene services staff. In addition, 19% of LTCF reported having no system in place for the organisation, control and feedback on hand hygiene

- The provision of seasonal influenza vaccination for residents was not universal, with 6% of LTCF overall reporting this was not routine local practice
- The vast majority (95%) reported having no active local antimicrobial stewardship committee (ASC), training on antimicrobial prescribing was not provided by 95% and just over two thirds (68%) of LTCF reported having no local antimicrobial prescribing guidelines
- Prescriber feedback regarding local antimicrobial use and local microbiology laboratory antimicrobial susceptibility data for common pathogens causing infection was available in only a minority of LTCF (13% and 7%, respectively)
- LTCF with a designated coordinating physician were significantly more likely to demonstrate positive local antimicrobial stewardship practices such as; an active ASC, prescribing guidelines, restrictive prescribing policy and provision of antimicrobial consumption data

Resident demographics, nursing care requirements and HCAI risk factors

- Female residents predominated across all care types and the proportion aged ≥ 85 years was highest in GN>12m (47%), Mixed>12m (41%) and LTCF<12m (38%). In contrast, only 1% of intellectually disabled LTCF residents were aged ≥ 85 years
- Indicators of resident nursing care requirements (incontinence, disorientation and impaired mobility) were evident in all care types, but most prevalent in GN>12m, Mixed>12m and LTCF<12m
- HCAI risk factors (presence of urinary or vascular catheter, pressure sores or other wounds) were most prevalent in residents of palliative care LTCF
- It was largely uncommon for residents to have a history of recent surgery, other than in rehabilitation facilities (5%) and LTCF<12m (4%)

HCAI

- The national crude HCAI prevalence was 5.3% and the national median HCAI prevalence was 4.2%. The median prevalence was higher in rehabilitation (7.8%), LTCF<12m (8.3%), Mixed>12m (6.1%) and the highest prevalence was reported in palliative care (18%), which

may reflect the HCAI risk factors encountered in that unique resident cohort. The lowest median HCAI prevalence was reported from GN>12m (4.2%) psychiatric (4.3%) and physically disabled LTCF (no HCAI detected in 46 residents)

- The most prevalent HCAI types were: respiratory tract infections (RTI), urinary tract infections (UTI) and skin infections; affecting 1.9%, 1.7% and 1.3% of all residents, respectively

Antimicrobial use and antimicrobial resistance

- The national crude antimicrobial use prevalence was 9.8% and the national median antimicrobial use prevalence was 9.7%. The median prevalence was higher in LTCF<12m (11.2%). At 24.5%, the prevalence in palliative care was more similar to antimicrobial use prevalence reported from acute hospital settings
- The majority of antimicrobials were prescribed within the LTCF (81%), mainly by GPs and directly-employed doctors
- Whilst the majority of antimicrobials were prescribed to treat infection, the proportion that were prescribed for infection prevention/prophylaxis was particularly high in intellectually disabled LTCF (49%), GN>12m (39%) and Mixed>12m (35%)
- During HALT 2013, 3.2% of GN>12m, 2.9% of Mixed>12m and 2% of intellectually disabled LTCF residents were prescribed antimicrobials for UTI prophylaxis. Prophylaxis against RTI (1.9%) and skin infection (1.4%) was most prevalent in intellectually disabled LTCF
- A relevant microbiological specimen had been obtained for just over one quarter of antimicrobial prescriptions (27%), with *Escherichia coli* (33%) and *Staphylococcus aureus* (22%) the two most frequently reported pathogens. Of those with available antimicrobial susceptibility results, 29% of *E. coli* were resistant to 3rd generation cephalosporins and 44% of *S. aureus* were meticillin/flucloxacillin resistant (i.e., MRSA). There were no carbapenem resistant *Enterobacteriaceae* reported during the HALT survey

1. Results

1.1 National Overview

1.1.1 Description of Participating LTCF

There was an excellent response to participate in the voluntary 2013 HALT survey, with a continued increase in participating LTCF; from 69 (2010) to 108 (2011) to 190 (2013), as displayed in Table 3.1.1. Fifty LTCF have participated in all three HALT surveys to date, 33 participated in 2011 and again in 2013, with 100 (53%) participating in HALT for the first time in 2013. In 2013, a designation for voluntary ownership was included for the first time.

There has also been an annual increase in participating LTCF across all care types. LTCF delivering care to eight major resident care types participated in HALT 2013, with four care types having more than one participant for the first time in 2013 [palliative care, physically disabled, and rehabilitation LTCF] (Table 3.1.1).

Table 1.1.1 Annual increases in HALT participation, by ownership, HSE region and care type.

| Category | 2010 | 2011 | 2013 |
|----------------------------|-----------|------------|------------|
| <i>by Ownership</i> | | | |
| HSE | 61 | 84 | 128 |
| <i>South</i> | 8 | 18 | 38 |
| <i>West</i> | 32 | 34 | 42 |
| <i>Dublin Mid Leinster</i> | 14 | 22 | 23 |
| <i>Dublin North East</i> | 7 | 10 | 25 |
| Private | 8 | 24 | 39 |
| Voluntary | N/A | N/A | 23 |
| <i>by Care Type</i> | | | |
| General nursing homes | 30 | 58 | 112 |
| Mixed care facilities | 16 | 16 | 32 |
| Intellectually disabled | 8 | 15 | 24 |
| Psychiatric | 3 | 5 | 11 |
| Palliative care | 1 | 1 | 4 |
| Physically disabled | 1 | 1 | 3 |
| Rehabilitation | 1 | 1 | 2 |
| Other | 3 | 1 | 2 |
| National | 69 | 108 | 190 |

* Other care types in 2013 included: a young chronically ill unit and a paediatric mixed care unit.

Figure 3.1.1 displays the distribution by county and by percentage of HIQA-registered residential care settings for older people that participated in HALT 2013.




Figure 1.1.1 Distribution, by county and by percentage of HIQA-registered residential care settings for older people that participated in HALT 2013.

Table 3.1.2 displays participating LTCF, by care type and by the estimated LOS for the majority (>75%) of each LTCF’s residents. Most (n=166; 87%) estimated that the majority of their residents were expected to remain in the LTCF indefinitely (i.e., for a period between 12 months until end-of-life).

Table 1.1.2 Breakdown of LTCF care type, by estimated LOS *

| Care Type | Length of stay of majority of residents | | | | | Total |
|-------------------------|---|-------------|------------|-------------------|----------|------------|
| | <3 months | 3-12 months | >12 months | until end-of-life | Other | |
| General nursing care | 5 | 4 | 6 | 97 | 0 | 112 |
| Intellectually disabled | 0 | 0 | 2 | 22 | 0 | 24 |
| Mixed facility | 3 | 3 | 9 | 17 | 0 | 32 |
| Other | 0 | 0 | 0 | 2 | 0 | 2 |
| Palliative care | 2 | 0 | 0 | 1 | 1 | 4 |
| Physically disabled | 0 | 0 | 1 | 1 | 0 | 2 |
| Psychiatric | 0 | 1 | 3 | 5 | 2 | 11 |
| Rehabilitation | 3 | 0 | 0 | 0 | 0 | 3 |
| National | 13 | 8 | 21 | 145 | 3 | 190 |

*Estimated LOS of the majority of residents admitted to the LTCF = expected LOS for >75% of residents.

The HALT Steering Group agreed to further stratify participating LTCF for data analysis, taking into account both care type and LOS. Thus, LTCF were categorised into eight care types, as displayed in Table 3.1.3.

- The two largest LTCF categories, general nursing homes and mixed care type facilities were further divided into three groups based on estimated LOS for majority of residents:
 - General nursing homes with estimated LOS >12 months (long-stay) = GN>12m
 - Mixed care type facilities with estimated LOS >12 months (long-stay) = Mixed >12m
 - LTCF (either general nursing homes or mixed care type facilities) with estimated LOS <12 months (short-stay) = LTCF <12m
- LTCF caring for residents with intellectual disabilities (Intellectually disabled)
- LTCF caring for residents with psychiatric conditions (Psychiatric)
- Other care types: Facilities caring for residents with palliative care needs, rehabilitation needs, physical disability or other care types

GN>12m accounted for the majority of participating LTCF (n=103; 54%), followed by Mixed>12m (n=26; 14%) and intellectually disabled LTCF (n=24; 13%). The remaining 37 LTCF (19%) were distributed among a variety of other care types (Table 3.1.3).

Table 3.1.3 also displays further breakdown of each care type, by ownership, size, overall bed occupancy and availability of single rooms. Overall, there was a median of 46 beds (range = 5 - 203 beds) per LTCF and the median number of beds in privately-owned (n=59) and voluntary (n=53) LTCF was higher than that in HSE-owned LTCF (n=38).

Overall, the median bed occupancy was 94% and the median single room occupancy was 34%. However, there were striking differences in single room availability based on ownership, with much lower proportions of single room availability in HSE-owned (21% of residents) versus LTCF under voluntary (50% of residents) and private (76% of residents) ownership.

Information was also captured on the proportion of LTCF residents who were absent on the survey date due to hospital admission. On average, 2.4% of residents were absent due to hospitalisation.

Table 1.1.3 Breakdown of participating LTCF, by ownership and care type.

| Category | Number of LTCFs | Size of the facility | | | Total residents surveyed | Median proportion of single rooms available | Median percentage of beds occupied | Percentage of residents hospitalised |
|-----------------------------|-----------------|----------------------|----------|------------|--------------------------|---|------------------------------------|--------------------------------------|
| | | n | median | min | | | | |
| <i>by Ownership</i> | | | | | | | | |
| HSE | 128 | 38 | 5 | 186 | 5,622 | 21 | 93 | 0.9 |
| <i>by HSE Region</i> | | | | | | | | |
| South | 38 | 44 | 16 | 137 | 1,872 | 18 | 100 | 0.6 |
| West | 42 | 30 | 9 | 142 | 1,512 | 21 | 100 | 1.3 |
| Dublin Mid Leinster | 23 | 51 | 27 | 166 | 1,290 | 21 | 100 | 1.1 |
| Dublin North East | 25 | 26 | 5 | 186 | 948 | 24 | 100 | 0.7 |
| Private | 39 | 59 | 20 | 203 | 2,536 | 76 | 96 | 1.5 |
| Voluntary | 23 | 53 | 10 | 141 | 1,160 | 50 | 95 | 0.6 |
| <i>by Care Type</i> | | | | | | | | |
| General nursing > 12 months | 103 | 51 | 21 | 203 | 5,807 | 35 | 95 | 1.2 |
| Mixed > 12 months | 26 | 47 | 13 | 142 | 1,409 | 32 | 94 | 1.1 |
| Intellectually disabled | 24 | 34 | 5 | 137 | 1,060 | 56 | 98 | 0.6 |
| LTCFs < 12 months | 15 | 29 | 12 | 78 | 374 | 15 | 89 | 0.5 |
| Psychiatric | 11 | 25 | 12 | 110 | 345 | 24 | 92 | 0.3 |
| Palliative care | 4 | 24 | 12 | 46 | 89 | 43 | 88 | 0.0 |
| Physically disabled | 2 | 28 | 22 | 34 | 46 | 14 | 82 | 0.0 |
| Rehabilitation | 3 | 64 | 35 | 72 | 139 | 14 | 93 | 0.0 |
| Other | 2 | 31 | 29 | 32 | 49 | 34 | 88 | 2.0 |
| National | 190 | 46 | 5 | 203 | 9,318 | 34 | 94 | 2.4 |

1.1.2 Governance Structures

1.1.2.1 Provision of Nursing & Medical Care

Availability of 24-hour qualified nursing care is a prerequisite for participation in the HALT survey. In the majority of units (n= 181, 95%), nursing staff had direct access to residents' healthcare records. For the remaining 5%, this information was not provided.

A variety of models of medical care exist in Irish LTCF, as displayed in Figure 3.1.2. Care was provided by the resident's own GP in 35%, a directly-employed doctor in 41% and in the remaining 24%, a mixed care model was observed, with both GPs and directly-employed doctors providing medical care.

Regional differences were also observed within HSE-owned LTCF, with GP-led care predominant in the West (64%) and much less common in the South (13%) and Dublin-North-East (20%). Notably, the distribution of care types was similar across the regions.

Differences were also observed based on LTCF ownership (GP-led care predominating in 82% of privately-owned versus 22% in those under voluntary ownership) and by care type (GN>12m were more likely to have directly employed doctors than Mixed>12m; 40% versus 23%) as displayed in Figure 3.1.3.




Figure 1.1.2 Models of medical care provision in LTCF, by ownership type and HSE region (for HSE-owned facilities)




Figure 1.1.3 Models of medical care provision in LTCF, by care type

1.1.2.2 Coordination of Medical Care

Participants were asked to provide information regarding the coordination of medical care within the facility. This was defined as having a designated ‘coordinating physician’ to arrange medical activities and take responsibility for standardisation of practices/policies for resident care. Figure 3.1.4 displays the coordination of medical care.

Overall, 55% reported having no coordinating physician and this was higher in privately-owned LTCF (74%). For the 45% with a coordinating physician, a variety of models of care were delivered [internal (20%), external (18%) or a mixture of both (2%)].




Figure 1.1.4 Coordination of medical care, by LTCF ownership.

Differences in coordination of medical care were also observed when facilities were stratified by care type (Figure 3.1.5). Absence of a nominated coordinating physician was more common in both GN>12m and Mixed>12m categories (>60%), whereas a coordinating physician was present in all of the palliative care, rehabilitation and physically disabled LTCF.

As the largest care type, availability of a coordinating physician in GN>12m was analysed, based on ownership. Absence of a coordinating physician was significantly more common in private (78%) than HSE (53%) GN>12m [p=0.02].




Figure 1.1.5 Coordination of medical care, by care type.

For the 85 (45%) LTCF with a designated coordinating physician (whether internal, external or a combination of both), retrospective additional information was sought on the job title of the coordinating physician. Of the facilities that responded (n = 60), the majority of coordinating physicians were either a directly-employed doctor (n = 24, 28%) or a designated GP (n = 20, 24%). Other titles included a hospital specialist [e.g., geriatrician] (n = 12, 14%), a medical doctor who owned the facility (n = 1, 1%) or another type of medical doctor (n = 3, 4%). For the remaining 25 LTCF, the job title of the coordinating physician was not provided. Figure 3.1.6 displays a breakdown of the coordinating physician job title, by care type.




Figure 1.1.6 Type of coordinating physician, by LTCF type.

In 98% (n = 83) of LTCF with a coordinating physician, that person also delivered medical care to the residents and in the majority (96%), the coordinating physician had direct access to residents' healthcare records. In one LTCF direct access to records was not available and for two LTCF this was not reported.

Further information was sought on the duties performed by the coordinating physician. Figure 3.1.7 displays the frequency and variety of tasks undertaken. Frequently-reported duties were; organisation of an on-call service, coordination of resident vaccination, supervision of medical

records and care strategy development. The coordinating physician was less frequently reported to undertake roles specific to prevention of HCAI and antimicrobial resistance, such as development of IPC (16%) or antimicrobial prescribing (14%) policies.




Figure 1.1.7 Duties performed by the coordinating physician.

1.1.2.3 Infection Prevention & Control (IPC) Practices

Tables 3.1.4 & 3.1.5 display the IPC structures, educational practices, protocols, surveillance and additional activities. Further description of these categories is provided subsequently.

Table 1.1.4 Overview of IPC structures, education and protocols, by ownership and care type.

| | IPC Structure | | | IPC Education | | | IPC Protocols | | | | |
|----------------------------------|-------------------------|-------------------|---------------|-----------------------|---|-----------------------------------|---------------|--------------|---------------------------------|----------------------------------|-------------------------------|
| | Staff with IPC training | Expert IPC advice | IPC committee | Hand hygiene training | IPC training of nursing/paramedical staff | IPC training of GPs/medical staff | MRSA | Hand hygiene | Management of urinary catheters | Management of vascular catheters | Management of enteral feeding |
| | % | | | % | | | % | | | | |
| by Ownership type | | | | | | | | | | | |
| Private (n = 39) | 10 | 51 | 46 | 90 | 95 | 10 | 100 | 97 | 92 | 23 | 87 |
| Voluntary (n = 23) | 61 | 61 | 61 | 70 | 87 | 30 | 87 | 96 | 57 | 48 | 78 |
| HSE (n = 128) | 77 | 65 | 75 | 91 | 84 | 13 | 99 | 98 | 91 | 56 | 87 |
| by Care Type | | | | | | | | | | | |
| GN > 12 months (n = 103) | 52 | 66 | 67 | 92 | 88 | 14 | 100 | 98 | 90 | 41 | 88 |
| Private only (n = 32) | 9 | 50 | 44 | 91 | 94 | 9 | 100 | 97 | 91 | 25 | 84 |
| Voluntary only (n = 7) | 57 | 86 | 71 | 86 | 100 | 57 | 100 | 100 | 86 | 43 | 71 |
| HSE only (n = 64) | 73 | 72 | 78 | 94 | 84 | 11 | 100 | 98 | 91 | 48 | 92 |
| Mixed > 12 months (n = 26) | 58 | 46 | 73 | 88 | 92 | 8 | 96 | 96 | 88 | 50 | 92 |
| Intellectually disabled (n = 24) | 71 | 54 | 75 | 71 | 79 | 4 | 92 | 100 | 79 | 50 | 83 |
| LTCFs < 12 months (n = 15) | 87 | 73 | 67 | 93 | 80 | 20 | 100 | 100 | 100 | 73 | 93 |
| Psychiatric (n = 11) | 73 | 64 | 36 | 73 | 64 | 9 | 91 | 82 | 73 | 27 | 36 |
| Palliative care (n = 4) | 100 | 75 | 50 | 100 | 100 | 50 | 100 | 100 | 75 | 100 | 75 |
| Physically Disabled (n = 2) | 100 | 0 | 100 | 100 | 100 | 50 | 100 | 100 | 100 | 100 | 100 |
| Rehabilitation (n = 3) | 100 | 67 | 67 | 100 | 100 | 67 | 100 | 100 | 67 | 100 | 100 |
| National | 62 | 62 | 67 | 88 | 86 | 14 | 98 | 97 | 87 | 48 | 86 |

Table 1.1.5 Overview of IPC surveillance and general activities, by ownership and care type.

| | Surveillance | | | | General IPC activities | | | | | |
|----------------------------------|-------------------|--|---|-------------------------------|--|-------------------------|--|---|-------------------------------|---|
| | HCAI surveillance | Performing audits on IPC policies and procedures | Feedback of surveillance results to staff | Monitoring incidence of MDROs | Offering influenza immunisation to residents | Management of outbreaks | Organisation, control and feedback on hand hygiene | Decisions on transmission-based precautions for residents | Development of care protocols | Supervision of disinfection/sterilisation |
| | % | | | | % | | | | | |
| by Ownership type | | | | | | | | | | |
| Private (n = 39) | 36 | 62 | 49 | 54 | 97 | 92 | 72 | 82 | 82 | 62 |
| Voluntary (n = 23) | 39 | 65 | 61 | 30 | 78 | 87 | 70 | 70 | 78 | 48 |
| HSE (n = 128) | 35 | 55 | 56 | 57 | 95 | 83 | 85 | 80 | 77 | 59 |
| by Care Type | | | | | | | | | | |
| GN > 12 months (n = 103) | 37 | 53 | 55 | 53 | 97 | 87 | 79 | 83 | 80 | 61 |
| Private only (n = 32) | 41 | 59 | 53 | 56 | 97 | 94 | 69 | 84 | 78 | 63 |
| Voluntary only (n = 7) | 57 | 57 | 57 | 43 | 100 | 100 | 43 | 86 | 57 | 71 |
| HSE only (n = 64) | 33 | 50 | 56 | 53 | 97 | 83 | 88 | 83 | 83 | 59 |
| Mixed > 12 months (n = 26) | 42 | 69 | 54 | 62 | 96 | 88 | 92 | 81 | 88 | 65 |
| Intellectually disabled (n = 24) | 21 | 71 | 50 | 46 | 92 | 79 | 75 | 58 | 75 | 42 |
| LTCFs < 12 months (n = 15) | 40 | 60 | 53 | 60 | 100 | 80 | 87 | 73 | 73 | 53 |
| Psychiatric (n = 11) | 18 | 9 | 27 | 45 | 82 | 73 | 55 | 64 | 45 | 45 |
| Palliative care (n = 4) | 50 | 75 | 100 | 25 | 25 | 75 | 100 | 100 | 100 | 75 |
| Physically Disabled (n = 2) | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 50 |
| Rehabilitation (n = 3) | 33 | 67 | 100 | 33 | 67 | 100 | 100 | 100 | 67 | 67 |
| National | 36 | 57 | 55 | 53 | 94 | 85 | 81 | 79 | 78 | 58 |

MDROs: Multi-drug resistant organisms

1.1.2.3.1 Staff with Training in IPC & Access to Advice from External IPC Experts

Overall, 117 (62%) LTCF reported access to a staff member with IPC training. However, LTCF under HSE or voluntary services ownership were more likely to have access to staff with IPC training (77% and 61%, respectively) than LTCF under private ownership (10%).

Of the 117 LTCF reporting a staff member with IPC training, for 83 (71%), that person was not based within the LTCF on a day-to-day basis, for 27 (23%) that person was based within the LTCF on an ongoing basis and for seven (6%) that person attended the LTCF on a sessional basis. Where a staff member with IPC training was available, for the majority of LTCF, that person was a nurse (n=109; 93%). Seven LTCF (6%) reported having both a nurse and a doctor with IPC training and one LTCF (1%) reported having a doctor with IPC training. Four LTCF were governed by an acute hospital. Thus, for those LTCF, the IPC service was provided by the acute hospital’s IPC team.

There was considerable geographic variability in the proportion of LTCF beds with access to an IPCN when distributed by county (range: 0 – 100%), as displayed in Figure 3.1.8.




Figure 1.1.8 Geographical distribution of the proportion of participating LTCF beds covered by IPCNs.

* Includes LTCF beds that participated in HALT only

A total of 36 IPCNs were available to participating LTCF. Excluding the four IPCNs attached to LTCF governed by acute hospitals, the estimated overall WTE ratio of IPCNs to LTCF beds in Ireland was 1:496. This ratio was calculated by including all LTCF beds that the IPCN was responsible for, which included both participating and non-participating HALT LTCF.

In HSE-owned LTCF, the estimated ratio of WTE IPCNs to LTCF beds was highest in HSE South and West (1:673 and 1:659) when compared to Dublin-mid-Leinster and Dublin-North-East (1:387 and 1:354) (Table 3.1.6).

Table 1.1.6 Ratio of WTE IPCN per HSE-owned LTCF, by HSE region.

| HSE Region | Number of HSE LTCFs that participated | Number of facilities with an IPCN | WTE ratio ^a |
|---------------------|---------------------------------------|-----------------------------------|------------------------|
| Dublin Mid Leinster | 22 | 14 | 1:387 |
| Dublin North East | 24 | 20 | 1:354 |
| South | 36 | 24 | 1:673 |
| West | 42 | 37 | 1:659 |

^a The whole time equivalent (WTE) ratio per HSE region was calculated by dividing the total number of HSE beds that IPCNs* within that region are responsible for by the total WTE associated with those IPCNs.

In addition to having access to staff with IPC training, information was sought on access to external expert IPC advice. Overall, 117 LTCF (62%) reported having access to such advice, no access was reported by two LTCF caring for residents with physical disability, and less access for LTCF caring for intellectually disabled (54%) and Mixed>12m (46%).

1.1.2.3.2 Infection Prevention and Control Committee (IPCC)

Just over two-thirds reported having an active local IPCC, with a median number of three meetings per year (range = 0 – 15). Psychiatric (36%) and palliative care facilities (50%) were less likely to have an IPCC.

1.1.2.3.3 Hand Hygiene and Access to Personal Protective Equipment (PPE)

Hand Hygiene Training

Overall, 88% reported that a staff hand hygiene training session had been organised during the previous year. This figure was higher in GN>12m (92%) and LTCF<12m (93%) and lower in intellectually disabled (71%) and psychiatric (73%) LTCF.

Overall, there was variation in the staff categories invited to avail of annual hand hygiene training as displayed in Table 3.1.7. Nursing (100%) and hygiene services staff (92%) were most likely to be invited and medical (27%) and allied health professional (28%) staff were less likely to be invited. By category of ownership, medical and allied health staff were less likely to be invited to attend hand hygiene training in privately-owned LTCF (11% and 14%, respectively) and more likely to be invited in facilities under voluntary ownership (both 53%).

Table 1.1.7 Staff categories invited to avail of annual hand hygiene education.

| Ownership | Nurses and care | | | Hygiene |
|-----------------|-----------------|------------|----------------------|----------------|
| | Medical staff | assistants | Allied health staff* | services staff |
| | % | % | % | % |
| Private | 11 | 100 | 14 | 92 |
| HSE | 28 | 100 | 28 | 93 |
| Voluntary | 53 | 100 | 53 | 80 |
| National | 27 | 100 | 28 | 92 |

* physiotherapist, occupational therapist, speech therapist etc

Access to hand hygiene products

The vast majority of LTCF reported having both alcohol-based hand rub (ABHR) and liquid soap (both 97%) as hand hygiene products (Table 3.1.8). Alcohol-based wipes were reported to be available in 41% of LTCF. ABHR was reported as the preferred hand hygiene method in 53% and the average estimated volume of ABHR consumed during the previous year (2012) was 118 litres. For 36% of LTCF, hand washing with a non-antiseptic soap was the preferred method and for 10% antiseptic soap was the preferred method.

Table 1.1.8 Availability of hand hygiene products and preferred hand hygiene methods.

| Hand hygiene products/methods | % of LTCFs |
|---|------------|
| Product | |
| Alcohol rub | 97 |
| Liquid soap | 97 |
| Wipes | 41 |
| Bar soap | 0 |
| Method | |
| Hand disinfection with an alcohol solution | 53 |
| Hand washing with water and a non antiseptic soap | 36 |
| Hand washing with water and an antiseptic soap | 10 |

Access to personal protective equipment (PPE)

All reported having both gloves and aprons available for healthcare worker PPE. Gowns were available in 99%, surgical masks in 88% and goggles in 66% of LTCF.

1.1.2.3.4 Availability of Written Protocols

Information regarding the availability of written protocols for staff on the following topics was sought (Table 3.1.4):

- *Management of MRSA and other multi-drug resistant organisms (MDRO)*: Available in 98% overall, with lower rates reported from psychiatric (91%) and intellectually disabled LTCF (92%)
- *Hand hygiene*: Available in 97% overall, with lower rates reported from psychiatric LTCF (82%)
- *Management of urinary catheters*: Available in 87% overall, with lower rates reported from rehabilitation (67%), psychiatric (73%) and palliative care facilities (75%)
- *Management of vascular catheters*: Available in 48% overall, with higher rates from palliative care (100%) and rehabilitation (100%) and lower rates reported from Mixed>12m and intellectually disabled LTCF (both 50%)
- *Management of enteral feeding*: Available in 86% overall, with higher rates from LTCF<12m (93%), Mixed>12m (92%) and lower rates reported from psychiatric LTCF (36%)

1.1.2.3.5 HCAI Surveillance Programme

- Some form of a HCAI surveillance programme was reported by 36% (n=68) of LTCF (Table 3.1.5). Ongoing participation in repeated HALT surveys could be regarded as an annual HCAI surveillance programme. Some care types were more likely to report having HCAI surveillance activities; palliative care (50%) and Mixed>12m (42%). Lower levels were reported from rehabilitation (33%), intellectually disabled (21%) and psychiatric (18%) LTCF
- Overall, just over half reported that audits of IPC policies and procedures are conducted (57%), that surveillance results are fed back to staff (55%) and that the incidence of MRDO is monitored (53%)
- Notably, facilities with an active IPCC were much more likely to report having MDRO surveillance programmes than those without
- A designated staff member for reporting and management of infection outbreaks was available in 85% overall, with lower levels reported from psychiatric (73%) and palliative care (75%) LTCF
- A system in place for the organisation, control and feedback on hand hygiene was available in 81% overall, with lower levels reported from psychiatric (55%) and intellectually disabled (75%) LTCF
- A system for management of patients with resistant organisms (e.g., patient isolation, additional IPC precautions) was available in 79% overall, with less availability in intellectually disabled (58%), psychiatric (64%) and LTCF<12m (73%)
- Overall, a system for development of resident care protocols available in 78% and a system for supervision of disinfection/sterilisation of medical equipment in only 58% of LTCF
- Overall, the majority (94%) reported that seasonal influenza vaccine is offered to residents. However, lower figures were reported from rehabilitation (67%) and psychiatric (82%) LTCF

1.1.2.4 Antimicrobial Stewardship Practices

Antimicrobial stewardship practices, stratified by LTCF ownership, care type and presence of a designated coordinating physician are displayed in Table 3.1.9.

Table 1.1.9 Antimicrobial stewardship practices, by LTCF ownership, care type and presence of a coordinating physician.

| | Antimicrobial stewardship committee | Training of prescribers on antimicrobial use | Guidelines for appropriate antimicrobial use | Data about antimicrobial consumption | Microbiological samples taken before antimicrobials | Local antimicrobial resistance profile summaries | Permission for prescribing restricted antimicrobials | Pharmacist giving advice on antimicrobial use | Therapeutic formulary available | Feedback to GPs on antimicrobial consumption |
|--|-------------------------------------|--|--|--------------------------------------|---|--|--|---|---------------------------------|--|
| | % | | | | | | | | | |
| by Ownership type | | | | | | | | | | |
| Private (n = 39) | 0 | 5 | 28 | 13 | 21 | 13 | 0 | 44 | 21 | 10 |
| Voluntary (n = 23) | 13 | 22 | 43 | 22 | 22 | 13 | 0 | 43 | 39 | 9 |
| HSE (n = 128) | 5 | 2 | 31 | 11 | 17 | 4 | 6 | 27 | 37 | 10 |
| by Care Type | | | | | | | | | | |
| GN > 12 months (n = 103) | 6 | 4 | 28 | 16 | 19 | 8 | 5 | 39 | 33 | 13 |
| Private only (n = 32) | 0 | 6 | 22 | 16 | 25 | 16 | 0 | 44 | 25 | 13 |
| Voluntary only (n = 7) | 14 | 14 | 43 | 29 | 29 | 14 | 0 | 29 | 14 | 14 |
| HSE only (n = 64) | 8 | 2 | 30 | 14 | 16 | 3 | 8 | 38 | 39 | 13 |
| Mixed > 12 months (n = 26) | 0 | 0 | 54 | 8 | 19 | 8 | 0 | 27 | 35 | 12 |
| Intellectually disabled (n = 24) | 8 | 13 | 17 | 8 | 21 | 4 | 0 | 17 | 38 | 8 |
| LTCFs < 12 months (n = 15) | 0 | 0 | 40 | 13 | 27 | 0 | 13 | 20 | 27 | 7 |
| Psychiatric (n = 11) | 0 | 0 | 9 | 0 | 0 | 0 | 0 | 18 | 9 | 0 |
| Palliative care (n = 4) | 0 | 25 | 100 | 25 | 0 | 25 | 0 | 100 | 75 | 0 |
| Physically disabled (n = 2) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 100 | 0 |
| Rehabilitation (n = 3) | 0 | 0 | 67 | 0 | 0 | 0 | 0 | 33 | 0 | 0 |
| by Presence of a Coordinating Physician | | | | | | | | | | |
| With a CP | 9 | 8 | 41 | 20 | 19 | 9 | 8 | 40 | 41 | 13 |
| Without a CP | 1 | 2 | 25 | 7 | 18 | 5 | 1 | 27 | 28 | 8 |
| Chi-test (p-value) | 0.02 | 0.052 | 0.02 | 0.01 | 0.9 | 0.2 | 0.02 | 0.54 | 0.052 | 0.24 |
| National | 5 | 5 | 32 | 13 | 18 | 7 | 4 | 33 | 34 | 10 |

* Chi-test p-values that reached significance are highlighted in bold

1.1.2.4.1 Overview of Antimicrobial Stewardship Practices & Guidelines

- The vast majority (95%) reported having no antimicrobial stewardship committee (ASC). Of the nine LTCF with an ASC (5%), none were privately-owned
- Additionally, the vast majority (95%) reported that annual training on antimicrobial prescribing was not provided
- Just under one third (32%) reported having a local antimicrobial prescribing guideline, with less availability reported by psychiatric (9%), intellectually disabled (17%) and GN>12m (28%). Prescribing guidelines were more likely to be available in palliative care (100%), rehabilitation (67%) and Mixed>12m (54%) facilities
- The vast majority (88%) reported having no restrictions on the types of antimicrobials that could be prescribed for residents. Of the 22 that reported having a restricted antimicrobial list, the types of restricted antimicrobials are displayed in Table 3.1.10

Table 1.1.10 Types of restricted antimicrobials.

| Restricted antimicrobials | Number of LTCFs (%) |
|-------------------------------|---------------------|
| 3rd generation cephalosporins | 15 (68) |
| carbapenems | 14 (64) |
| vancomycin | 13 (59) |
| fluoroquinolones | 8 (36) |
| intravenous antimicrobials | 8 (36) |
| broad-spectrum antibiotics | 6 (27) |
| glycopeptides | 5 (23) |
| mupirocin | 3 (14) |
| Total | 22 (100) |

- One-third reported having access to the advice of a pharmacist as required, if residents were prescribed unusual antimicrobials, with more access reported by LTCF under private (44%) or voluntary (43%) ownership
- Local antimicrobial consumption data was collected by the minority (13%), such data was reported back to GPs by a minority (10%) and a minority (7%) reported having access to summary reports of antimicrobial resistance in key pathogens from their local microbiology laboratory
- A minority (18%) reported having a system in place to remind staff of the importance of obtaining relevant clinical specimens from the resident prior to commencing antimicrobial

therapy (e.g., the importance of taking a urine specimen before starting treatment for a suspected UTI)

- Information was sought regarding the frequency with which a urine dipstick test was used for UTI diagnosis. Of the 186 (98%) who answered, urine dipstick was performed routinely in 146 (78%), on occasion in 39 (21%) and never in one LTCF (1%)
- Specific information was also sought on the availability of local antimicrobial prescribing guidelines for three common infection types, as displayed in Table 3.1.11:
 - RTI (35%)
 - UTI (43%)
 - Wound/skin or soft tissue infection (43%)

Table 1.1.11 Written antimicrobial treatment guidelines.

| | Antimicrobial treatment guidelines | | |
|--|------------------------------------|--------------------------|----------------------------------|
| | Respiratory tract infections | Urinary tract infections | Wound and soft tissue infections |
| by Ownership | | | |
| Private | 26 | 42 | 39 |
| HSE | 36 | 41 | 42 |
| Voluntary | 45 | 55 | 55 |
| by Presence of a Coordinating Physician | | | |
| With a CP | 46 | 50 | 51 |
| Without a CP | 27 | 37 | 37 |
| National | 35 | 43 | 43 |

- When LTCF were stratified by the presence or absence of a coordinating physician, the presence of a coordinating physician was significantly associated with a higher prevalence of positive antimicrobial stewardship practices, in particular the existence of an ASC, antimicrobial prescribing guidelines, restrictive prescribing policies and antimicrobial consumption data

1.1.3 HCAI and Antimicrobial Use

1.1.3.1 Description of Residents

Table 3.1.12 displays an overview of the resident demographics, care load indicators and HCAI risk factors, by care type. Female residents predominated across all care types. There was a much higher proportion of residents aged ≥85 years in; GN>12m, Mixed>12m, LTCF<12m and rehabilitation LTCF.

Care load indicators (incontinence, disorientation and impaired mobility) were evident but varied across all care types. Overall, there was a heavy burden of all care load indicators in GN>12m, Mixed>12m and LTCF<12m.

HCAI risk factors were most prevalent in palliative care, where urinary and vascular catheters and ‘other wounds’ were much more common than for other care types. Overall, recent surgery within the past 30 days was uncommon in residents, with rehabilitation (5%) and LTCF<12m (4%) more likely to accommodate those resident types. Residents of psychiatric LTCF were less likely to have HCAI risk factors.

Table 1.1.12 Resident demographics, care load indicators and HCAI risk factors, by care type.

| Facility Type | Gender | Age | Care load indicators | | | HCAI Risk factors | | | | |
|-------------------------|----------------|--------------------|----------------------|----------------|-------------------|-------------------|-------------------|----------------|--------------|--------------------|
| | male residents | resident >85 years | incontinence | disorientation | impaired mobility | urinary catheter | vascular catheter | pressure sores | other wounds | surgery (<30 days) |
| % | | | | | | | | | | |
| GN > 12 months | 35 | 47 | 62 | 57 | 48 | 5 | 0 | 4 | 10 | 1 |
| Mixed > 12 months | 39 | 41 | 65 | 54 | 54 | 8 | 1 | 4 | 11 | 3 |
| LTCFs < 12 months | 36 | 38 | 45 | 33 | 40 | 14 | 1 | 8 | 11 | 4 |
| Intellectually disabled | 45 | 1 | 52 | 54 | 33 | 3 | 0 | 1 | 9 | 0 |
| Psychiatric | 48 | 10 | 43 | 29 | 23 | 2 | 1 | 2 | 2 | 1 |
| Palliative care | 44 | 9 | 27 | 26 | 60 | 31 | 10 | 9 | 36 | 1 |
| Physically Disabled | 48 | 9 | 78 | 59 | 74 | 7 | 0 | 0 | 2 | 0 |
| Rehabilitation | 40 | 29 | 22 | 17 | 21 | 8 | 2 | 1 | 15 | 5 |
| Other | 71 | 0 | 78 | 53 | 84 | 2 | 0 | 6 | 39 | 0 |
| National | 37 | 38 | 59 | 53 | 46 | 6 | 0.5 | 4 | 11 | 1 |

1.1.3.2 HCAI

The national median HCAI prevalence was 4.2%. Table 3.1.13 displays the HCAI prevalence, by care type. Similar to the distribution of HCAI risk factors, as displayed in Table 3.1.12 above, the median prevalence of HCAI was highest in palliative care (18%) and lowest in intellectually disabled (2.2%) and physically disabled LTCF (no reported HCAI in 46 residents). Median HCAI prevalence was higher in rehabilitation (7.8%) and LTCF<12m (8.3%) than Mixed>12m (6.1%) and GN>12m (4.2%). The median HCAI prevalence in GN>12m and psychiatric LTCF was quite similar (4.2% and 4.3%).

Table 1.1.13 HCAI prevalence, by care type.

| Facility Type | Total eligible residents | Number of residents with an infection | HCAI prevalence (%) | |
|-------------------------|--------------------------|---------------------------------------|---------------------|----------------------------|
| | | | Crude ^a | Median (IQR ^b) |
| GN > 12 months | 5,807 | 294 | 5.1 | 4.2 (2 - 7) |
| Mixed > 12 months | 1,409 | 86 | 6.1 | 6.1 (2.6 - 8.5) |
| Intellectually disabled | 1,060 | 46 | 4.3 | 2.2 (0 - 6.8) |
| LTCFs < 12 months | 374 | 28 | 7.5 | 8.3 (0.7 - 10.9) |
| Psychiatric | 345 | 11 | 3.2 | 4.3 (0 - 6.5) |
| Rehabilitation | 139 | 11 | 7.9 | 7.8 (6 - 11.3) |
| Palliative care | 89 | 18 | 20.2 | 18.0 (14.1 - 21.8) |
| Physically disabled | 46 | 0 | 0.0 | 0.0 (0 - 0) |

^a The crude prevalence of residents with a HCAI is the total number of residents with an infection divided by the total number of eligible residents.

^b The interquartile range is the difference between the first quartile (25th percentile) and the third quartile (75th percentile) of an ordered range of data. It represents the middle fifty percent of the data.

HCAI Types

Figure 3.1.9 displays the prevalence of HCAI, by care type.

RTI

- Overall, RTI was the most prevalent HCAI, affecting 1.9% of all residents. RTI were further categorised into; lower RTI (68%), common cold (23%), pneumonia (8%) and flu (2%)
- RTI was the most prevalent (or one of the most prevalent) HCAI reported by Mixed>12m (2.3%), intellectually disabled (1.6%) and rehabilitation (2.9%) LTCF. RTI was less prevalent in psychiatric LTCF (0.9%)

UTI

- Overall, UTI was the second most prevalent HCAI, affecting 1.7% of all residents. In total, 33% were reported as microbiologically-confirmed UTI
- UTI was the most prevalent (or one of the most prevalent) HCAI, reported by LTCF<12m (3.2%), GN>12m (1.7%) and rehabilitation LTCF (2.9%). UTI was less prevalent in intellectually disabled (0.6%) and psychiatric LTCF (0.6%)

Skin infections

- Skin infections were the third most prevalent HCAI, affecting 1.3% of all residents. The vast majority were further categorised as cellulitis (94%)
- Skin infections were the most prevalent (or one of the most prevalent) HCAI reported by intellectually disabled (1.6%) and psychiatric LTCF (1.4%)




Figure 1.1.9 Prevalence of HCAI, by care type.

Only LTCF care types including > 100 eligible residents were included for this analysis

1.1.3.3 Antimicrobial Use

The national median antimicrobial use prevalence was 9.8%. Table 3.1.14 displays antimicrobial use prevalence, by care type. The median prevalence was highest in palliative care (24.5%) and Mixed>12m (11.2%) and lowest in psychiatric (4.7%) and physically disabled (no antimicrobials prescribed) LTCF.

Table 1.1.14 Antimicrobial use prevalence, by care type.

| Facility Type | Total eligible residents | Number of residents on antimicrobials | Antimicrobial prevalence (%) | |
|-------------------------|--------------------------|---------------------------------------|------------------------------|-------------------|
| | | | Crude | median (IQR) |
| GN > 12 months | 5807 | 541 | 9.3 | 9.1 (5.7 - 13.3) |
| Mixed > 12 months | 1409 | 149 | 10.6 | 11.2 (8.2 - 15.3) |
| Intellectually disabled | 1060 | 106 | 10.0 | 7.5 (3.2 - 13.9) |
| LTCFs < 12 months | 374 | 42 | 11.2 | 9.5 (5.5 - 16.7) |
| Psychiatric | 345 | 23 | 6.7 | 4.7 (4.3 - 11.7) |
| Rehabilitation | 139 | 14 | 10.1 | 9.4 (8.9 - 12.1) |
| Palliative care | 89 | 31 | 34.8 | 24.5 (19.7 - 33) |
| Physically Disabled | 46 | 0 | 0.0 | 0.0 (0 - 0) |

Antimicrobial Prescribers and Prescribing Location

Across all care types, the vast majority of antimicrobials were prescribed within the LTCF (81%), as displayed in Figure 3.1.10. A higher proportion of antimicrobials were prescribed in the hospital setting in Mixed>12m (15%), intellectually disabled (14%), and palliative (18%) LTCF.




Figure 1.1.10 Antimicrobials, by prescribing location and care type.

Combined, GPs and directly-employed doctors accounted for the majority of prescribers across the care types, as displayed in Figure 3.1.11. Hospital-based specialists accounted for increasing proportions of antimicrobials in rehabilitation (33%) and palliative care (45%) LTCF.




Figure 1.1.11 Antimicrobials, by prescriber and care type.

Reasons & Sites for which Antimicrobials were Prescribed

The reason for antimicrobials varied across care types, as displayed in Figure 3.1.12. Whilst the majority were for treatment of infection, the proportion prescribed for prophylaxis was highest in intellectually disabled (49%), GN>12m (39%) and Mixed>12m (35%) and lowest in LTCF<12m (17%) and rehabilitation (13%) LTCF.




Figure 1.1.12 Reason for antimicrobials, by care type.

Figure 3.1.13 displays the antimicrobial use prevalence, by body site across the care types.

- The urinary tract was the most prevalent site, accounting for antimicrobials prescribed to 4.6% of all residents. GN>12m (5.1%), Mixed>12m and LTCF<12m (4.8% each) had a slightly higher prevalence, whilst intellectually disabled (2.6%) and psychiatric (2.3%) LTCF had a lower prevalence
- The respiratory tract was the second most prevalent site, accounting for antimicrobials prescribed to 2.9% of all residents. Intellectually disabled (3.9%), Mixed>12m (3.6%) and LTCF<12m (3.2%) had a higher prevalence, whilst psychiatric (2.3%) and GN>12m (2.4%) had a lower prevalence
- Skin or wounds were the third most prevalent site, accounting for antimicrobials prescribed to 1.6% of all residents. Intellectually disabled LTCF had a higher prevalence (3.1%) and GN>12m, Mixed>12m and rehabilitation LTCF had a lower prevalence (1.4% each)




Figure 1.1.13 Prevalence of antimicrobial use, by body site and care type.

Only LTCF care types including > 100 eligible residents were included in this breakdown.

Figures 3.1.14 and 3.1.15 display the breakdown of antimicrobials for treatment and prophylaxis, by care type.

- Treatment of RTI and UTI were jointly the two most prevalent indications: LTCF<12m (3.2% each) and rehabilitation LTCF (2.9% each). UTI treatment was less prevalent in intellectually disabled (0.7%) and psychiatric LTCF (1.2%). Treatment of skin/wound infections was most prevalent in LTCF<12m (1.9%) and intellectually disabled LTCF (1.7%)
- UTI prevention was the most prevalent indication for antimicrobial prophylaxis (3.2% of GN>12m and 2.9% of Mixed>12m residents)
- Intellectually disabled LTCF had the highest prevalence of prophylaxis (5.3%), divided into urinary tract (2.0%), respiratory tract (1.9%) and skin/wound (1.4%) prophylaxis. Indeed, almost half of antimicrobials prescribed in intellectually disabled LTCF were for prevention of skin infection




Figure 1.1.14 Body sites for treatment of infection, by care type.

Only LTCF care types including > 100 eligible residents were included in this breakdown.




Figure 1.1.15 Body sites for prevention of infection (prophylaxis), by care type.

Only LTCF care types including > 100 eligible residents were included in this breakdown.

Prescribed Antimicrobials

Figure 3.1.16 displays the breakdown of prescribed antimicrobials, by care type. Co-amoxiclav was the most commonly prescribed antimicrobial. Nitrofurantoin, trimethoprim, flucloxacillin, ciprofloxacin, amoxicillin, clarithromycin and cefalexin were also frequently prescribed. See the separate reports of each care type for a more detailed description of antimicrobials.




Figure 1.1.16 Prescribed antimicrobials, by care type.

1.2 Mixed Care Facilities with a LOS greater than 12 months (Mixed > 12m)

1.2.1 Description of Care Type

There were 26 mixed care type LTCF in this category, with an estimated LOS for the majority of residents greater than 12 months (Mixed>12m). The majority were HSE-owned (20; 77%), with five under private (19%) and one under voluntary ownership (4%). Mixed>12m were distributed around Ireland: Leinster (9), Munster (11), Connacht (3) and Ulster (3), with each accommodating a median of 47 residents (range: 13 – 142).

1.2.2 Description of Residents

Of the 1,409 residents, females predominated (61%) and 41% were ≥85 years. Figure 3.3.1 displays resident demographics, care load indicators and risk factors for HCAI.

Care Load

Sixty-five percent of residents were incontinent of faeces and/or urine. Residents with an indwelling urinary catheter were not counted as incontinent, as per the HALT protocol. Over half were disoriented in time and/or space (54%) and 54% were suffering from impaired mobility (wheelchair-bound or bed-ridden).

HCAI Risk Factors

Of 1,409 Mixed>12m residents, 8% had a urinary catheter *in situ*. Vascular catheters were uncommon, with only 1% having this device. Pressure sores were present in 4% and 11% were reported to have an 'other wound' (e.g., leg ulcer, insertion site of a suprapubic catheter or gastrostomy tube, colostomy, ileostomy, tracheostomy, traumatic or surgical wound). Three percent had a history of surgery in the past 30 days.




Figure 1.2.1 Resident demographics, care load indicators and HCAI risk factors in Mixed>12m.

1.2.3 HCAI in Mixed > 12m

Of the 1,409 residents, 96 (6.8%) were reported to have signs or symptoms of infection. Of those, 87 ultimately met a HCAI definition. Therefore, the crude HCAI prevalence was 6.2% (median = 6.1%). Five residents had more than one HCAI. In total, there were 92 HCAI. (Table 3.3.1). There was a wide variation in the HCAI prevalence by individual LTCF (0–16.1%), with three Mixed>12m (12%) reporting no residents with HCAI.

Table 1.2.1 HCAI prevalence in Mixed>12m.

| HCAI prevalence data | |
|--|-------------|
| Number of LTCFs that participated in survey | 26 |
| Number of residents surveyed | 1409 |
| Number of residents with signs/symptoms of an infection | 96 |
| Number of residents with infections ^a | 87 |
| Number of infections | 92 |
| Residents with more than one infection | 5 |
| Crude prevalence of residents with a HCAI infection ^b | 6.2% |
| National median prevalence | 6.1% |
| National range (min - max) | 0 - 16.1% |
| National interquartile range ^c | 2.8 - 8.5% |

^a As defined by Stone *et al* 2012.[1]

^b The crude prevalence of residents with a HCAI is the total number of residents with an infection divided by the total number of eligible residents.

^c The interquartile range is the difference between the first quartile (25th percentile) and the third quartile (75th percentile) of an ordered range of data. It represents the middle fifty percent of the data.

Residents with HCAI: Demographics and HCAI Risk Factors

Of 87 residents with HCAI, the mean age was 81 years (range: 48 – 98) and the majority (70%) had been living in the LTCF for one year or longer. Recent hospitalisation (within past three months) was reported for 31% of Mixed>12m residents with HCAI.

Table 3.3.2 displays a comparison of Mixed>12m residents with HCAI and the overall Mixed>12m population, with regard to demographics, care load indicators and HCAI risk factors.

Residents with HCAI were significantly less likely to be aged >85 years (30% vs 41%), significantly more likely to have pressure sores (9% vs 4%) and ‘other wound’ types (28% vs 11%) when compared to the overall Mixed>12m population.

Table 1.2.2 Comparison of Mixed>12m resident demographics, care load indicators and risk factors for residents with HCAI compared to the total Mixed>12m population.

| Resident Characteristic | Total resident population (%) | Residents with an infection (%) | p value | Odd's ratio |
|--------------------------|-------------------------------|---------------------------------|---------------|-------------|
| <i>Gender/Age</i> | | | | |
| male | 39 | 41 | 0.6133 | |
| > 85 | 41 | 30 | 0.0323 | 0.6 |
| <i>Care Load</i> | | | | |
| Incontinent | 65 | 62 | 0.6227 | |
| Disoriented | 54 | 56 | 0.5971 | |
| Immobile | 54 | 62 | 0.1227 | |
| <i>HCAI Risk Factors</i> | | | | |
| Urinary Catheter | 8 | 10 | 0.4261 | |
| Vascular Catheter | 1 | 1 | 0.6866 | |
| Pressure Sores | 4 | 9 | 0.0161 | 2.5 |
| Other Wounds | 11 | 28 | 0.0000 | 3.3 |
| Recent Surgery | 3 | 3 | 0.6205 | |

*An odd’s ratio was only provided where the p-value reached significance (< 0.05).

HCAI Types

Three HCAI categories accounted for 84% of reported infections: RTI (33; 36%), UTI (24; 26%) and skin infections (19; 21%). Figure 3.3.2 displays the distribution of the 92 HCAI, by infection category and Table 3.3.3 provides a further breakdown of the HCAI types within each category.




Figure 1.2.2 Distribution of HCAI in Mixed>12m, by infection category.

Table 1.2.3 HCAI breakdown in Mixed>12m, by HCAI type and prevalence (**Appendix C**).

| Infection Type | Number of residents with infections | % of residents with infection |
|--|-------------------------------------|-------------------------------|
| Respiratory Tract Infection | 33 | 2.3% |
| <i>Cold</i> | 4 | 0.3% |
| <i>Pneumonia</i> | 5 | 0.4% |
| <i>Flu</i> | 1 | 0.1% |
| <i>Lower respiratory tract infection</i> | 23 | 1.6% |
| Urinary Tract Infection | 24 | 1.7% |
| <i>Confirmed</i> | 7 | 0.5% |
| <i>Probable</i> | 17 | 1.2% |
| Skin | 19 | 1.3% |
| <i>Cellulitis</i> | 18 | 1.3% |
| <i>Herpes</i> | 1 | 0.1% |
| <i>Fungal</i> | 0 | 0.0% |
| Eye, Ear, Nose, Mouth | 8 | 0.6% |
| <i>Eye</i> | 7 | 0.5% |
| <i>Ear</i> | 1 | 0.1% |
| <i>Mouth</i> | 0 | 0.0% |
| <i>Sinusitis</i> | 0 | 0.0% |
| Gastrointestinal | 3 | 0.2% |
| <i>Gastro</i> | 1 | 0.1% |
| <i>C. difficile</i> | 2 | 0.1% |
| Other | 4 | 0.3% |
| <i>Bloodstream infection</i> | 0 | 0.0% |
| <i>Fever</i> | 1 | 0.1% |
| <i>Other</i> | 3 | 0.2% |
| Total number of residents with infections | 86 | 6.1% |

RTI

RTI were the most prevalent HCAI, affecting 2.3% of residents. Of the 33 RTI, most (70%) were further categorised as 'lower respiratory tract infections'. 'Pneumonia' confirmed by chest X-ray was the second most prevalent RTI type (15%), followed by 'common cold or pharyngitis' (12%). One case meeting the 'flu' surveillance definition was reported (3%). It is notable that influenza activity in Ireland during May 2013 was at a low level.[2]

UTI

UTI were the second most prevalent HCAI, affecting 1.7% of residents. Of the 24 UTI, most (71%) were further categorised as 'probable UTI', based on absence of a positive urine microbiology culture result.

Skin Infections

Skin infections were the third most prevalent HCAI, affecting 1.3% of residents. Of the 19 infections, the vast majority (95%) were categorised as 'cellulitis/soft tissue/wound infections'. There was also one case of herpes infection (i.e., herpes simplex/cold sore or herpes zoster/shingles).

Eye, Ear, Nose & Mouth Infections

This infection category was the fourth most prevalent, affecting 0.6% of residents. Of the eight infections, conjunctivitis accounted for the majority (88%) and one case of ear infection was reported.

Gastrointestinal Infections

There were three cases in the category gastrointestinal infection, affecting 0.2% of residents. *C. difficile* infection (2) and gastroenteritis (1).

Other Infection Types

There were four additional infection types reported. Unexplained fever (1) and 'other' infections not categorised in the HCAI subtypes (3) above. There were no cases of bloodstream infection reported from Mixed>12m residents.

1.2.4 Antimicrobial Use in Mixed > 12m

Of the 1,409 Mixed>12m residents, 149 were prescribed systemic antimicrobials. Therefore, the crude prevalence of antimicrobial use was 10.6% (median = 11.2%). Seven residents were prescribed more than one antimicrobial. In total, information on 157 antimicrobials was recorded (Table 3.3.4). The majority were administered via the oral route (92%), with 6% intravenous. There was a wide variation in antimicrobial use prevalence by individual LTCF (0 – 20.8%), with one Mixed>12m facility (4%) reporting no residents on antimicrobials.

Table 1.2.4 Antimicrobial use prevalence in Mixed>12m.

| Antimicrobial prevalence data | |
|--|--------------|
| Number of residents surveyed | 1409 |
| Number of residents on antimicrobials | 149 |
| Number of antimicrobials prescribed | 157 |
| Number of residents on more than one antimicrobial | 7 |
| Crude prevalence of residents on antimicrobials ^a | 10.6% |
| National mean prevalence | 11.3% |
| National median prevalence | 11.2% |
| National range (min - max) | 0 - 20.8% |
| National interquartile range | 8.2 - 15.3% |

Residents Prescribed Antimicrobials: Demographics, Care Load Indicators and HCAI Risk Factors

Of 149 residents prescribed antimicrobials, the mean age was 80 years (range: 40 – 98) and 34% were ≥85 years. Table 3.3.5 displays a comparison of Mixed>12m residents on antimicrobials and the overall Mixed>12m population, with regard to demographics, care load indicators and risk factors.

Mixed>12m residents on antimicrobials were significantly more likely to have a urinary catheter *in situ* (15% vs 8%) and to have ‘other’ wound types (21% vs 11%).

Table 1.2.5 Comparison of Mixed>12m resident demographics, care load indicators and risk factors for residents on antimicrobials compared to the total Mixed>12m population.

| | Total resident population (%) | Residents on an antimicrobial (%) | p value | Odd's ratio |
|--------------------------|-------------------------------|-----------------------------------|--------------|-------------|
| <i>Gender/Age</i> | | | | |
| male | 39 | 44 | 0.147 | |
| > 85 | 41 | 34 | 0.084 | |
| <i>Care Load</i> | | | | |
| Incontinent | 65 | 60 | 0.267 | |
| Disoriented | 54 | 54 | 0.978 | |
| Immobile | 54 | 61 | 0.070 | |
| <i>HCAI Risk Factors</i> | | | | |
| Urinary Catheter | 8 | 15 | 0.002 | 2.2 |
| Vascular Catheter | 0.8 | 2.0 | 0.071 | |
| Pressure Sores | 4 | 6 | 0.232 | |
| Other Wounds | 11 | 21 | 0.000 | 2.3 |
| Recent Surgery | 3 | 5 | 0.094 | |

Antimicrobial Prescribers & Prescribing Location

For each prescribed antimicrobial, information was sought regarding the prescribing location and the prescriber (Table 3.3.6). The majority of antimicrobials in Mixed>12m (76%) were prescribed in the LTCF, followed by the hospital (15%).

GPs prescribed the majority of antimicrobials (48%), followed by directly-employed doctors (29%), hospital-based specialists (14%) and other healthcare professionals [e.g., nurse prescribers] (9%).

Table 1.2.6 Antimicrobial prescriptions in Mixed>12m, by prescribing location and prescriber.

| Where are antimicrobials prescribed? | % |
|--------------------------------------|----|
| In this facility | 76 |
| In the hospital | 15 |
| Elsewhere | 10 |
| Unknown | 0 |

| Who prescribes the antimicrobials? | % |
|---|----|
| GP | 48 |
| Medical doctor employed by the facility | 29 |
| Specialist | 14 |
| Other | 9 |
| Unknown | 1 |

Reasons & Sites for which Antimicrobials were Prescribed

The reason for each of the 157 antimicrobials prescribed to Mixed>12m residents was recorded. The majority were for treatment of infection (102; 65%), with the remainder for prevention of infection (hereafter known as prophylaxis) (55; 35%). Figure 3.3.3 displays the breakdown by the body site for which the 157 antimicrobials were prescribed. Combined, three body sites accounted for 91% of prescriptions.




Figure 1.2.3 Prescribed antimicrobials, by body site indication.

The vast majority of prophylaxis was for UTI prevention (76%). Treatment of RTI (45%) and UTI (25%) were the most frequent indications for therapeutic antimicrobials (Figure 3.3.4).




Figure 1.2.4 Breakdown of prophylaxis and treatment, by body site.

Urinary Tract

The urinary tract (UTI treatment or prophylaxis) accounted for the highest proportion of antimicrobials prescribed to Mixed>12m residents (68; 43%), with the majority of those prescribed for UTI prophylaxis (42; 62%). The prevalence of antimicrobial use for the urinary tract in Mixed>12m was 4.8% (prophylaxis 2.9% and treatment 1.8%). Residents prescribed UTI prophylaxis were significantly more likely to have an indwelling urinary catheter (20% vs 8%: p-value: 0.007, OR 2.9). Of the 26 residents prescribed antimicrobial therapy for suspected UTI, prior to commencing treatment a urine dipstick was performed for 22 (85%) and for 16 (62%), a urine specimen was sent to the microbiology laboratory for culture and susceptibility testing.

Respiratory Tract

The respiratory tract (RTI treatment or prophylaxis) accounted for 34% (n=53) of antimicrobials prescribed to Mixed>12m residents. The prevalence of antimicrobial use for the respiratory tract was 3.6% (prophylaxis 0.5% and treatment 3.1%).

Skin or Wound

The skin or wound (treatment or prophylaxis for skin/wound infection) accounted for 14% (n=22) of antimicrobials prescribed to Mixed>12m residents. The prevalence of antimicrobial use for skin/wound was 1.4% (prophylaxis 0.1% and treatment 1.3%).

Prescribed Antimicrobials

Table 3.3.7 displays the most frequently prescribed antimicrobials in Mixed>12m:

- i. Co-amoxiclav was the most common antimicrobial (24.8%). It was mostly prescribed to treat RTI (62%) and UTI (21%)
- ii. Nitrofurantoin was the second most common antimicrobial (13.4%). It was only prescribed for urinary tract indications, in particular for UTI prophylaxis (71% of prescriptions)
- iii. Trimethoprim was the third most common antimicrobial (11.5%). It was only prescribed for urinary tract indications, in particular for UTI prophylaxis (67% of prescriptions)

Table 1.2.7 Most frequently prescribed antimicrobials in Mixed>12m.

| Antimicrobial name | Number of prescriptions (%) |
|---------------------------|------------------------------------|
| co-amoxiclav | 39 (24.8) |
| nitrofurantoin | 21 (13.4) |
| trimethoprim | 18 (11.5) |
| flucloxacillin | 11 (7) |
| clarithromycin | 9 (5.7) |
| doxycycline | 7 (4.5) |
| amoxicillin | 6 (3.8) |
| methenamine | 6 (3.8) |
| ciprofloxacin | 5 (3.2) |
| cefalexin | 4 (2.5) |
| other | 31 (19.7) |
| Total | 157 (100) |

2. Appendices

2.1 Appendix A: List of Acronyms used in this Report

| | |
|------------------|---|
| ABHR | Alcohol-Based Hand Rub |
| ASC | Antimicrobial Stewardship Committee |
| CDC | US Centers for Disease Control & Prevention |
| CDI | <i>Clostridium difficile</i> infection |
| CRE | Carbapenem resistant <i>Enterobacteriaceae</i> |
| ECDC | European Centre for Disease Prevention and Control |
| ESBL | Extended Spectrum Beta Lactamase |
| FAQ | Frequently-Asked Questions |
| GN>12m | General nursing homes with LOS > 12 months |
| GP | General Practitioner |
| HAI | Hospital-acquired Infection |
| HALT | Healthcare-Associated Infections in Long-Term Care Facilities |
| HCAI | Healthcare-Associated Infection |
| HIQA | Health Information & Quality Authority |
| HPSC | Health Protection Surveillance Centre |
| HSE | Health Service Executive |
| IPC | Infection Prevention & Control |
| IPCC | Infection Prevention & Control Committee |

| | |
|---------------------|--|
| IPCN | Infection Prevention & Control Nurse |
| LOS | Length-of-Stay |
| LTCF | Long-Term Care Facility |
| LTCF<12m | LTCF (either general nursing home or mixed care type) with LOS < 12 months |
| MHC | Mental Health Commission |
| Mixed>12m | Mixed care facilities with LOS > 12 months |
| MRSA | Meticillin Resistant <i>Staphylococcus aureus</i> |
| MSSA | Meticillin Susceptible <i>Staphylococcus aureus</i> |
| MDRO | Multi-Drug Resistant Organisms |
| PPE | Personal Protective Equipment |
| PPS | Point Prevalence Survey |
| RCPI | Royal College of Physicians of Ireland |
| RTI | Respiratory Tract Infection |
| SHEA | Society for Healthcare Epidemiology of America |
| UTI | Urinary Tract Infection |
| VRE | Vancomycin Resistant Enterococci |
| WIV-ISP | Scientific Institute for Public Health, Brussels, Belgium |
| WTE | Whole Time Equivalent |

PLEASE NOTE:

This report represents a shortened version of the full national HALT report focusing on data from Mixed Care Facilities.

The full national report including methodology, results on all LTCF care types, discussion and future priorities can be found on the hpsc website [here](#).