

# Weekly Report on Severe Acute Respiratory Infections (SARI), Ireland

**Week 50 2024 (week ending 15/12/2024)**

**Report prepared on 18/12/2024**

## About this report

Three sentinel hospital sites are now participating in the severe acute respiratory infections (SARI) surveillance programme in Ireland. Along with St Vincent's University Hospital (SVUH) (commenced on 5<sup>th</sup> July 2021), both St James's Hospital (SJH) and University Hospital Limerick (UHL) (paediatric cases only) commenced SARI surveillance on 30<sup>th</sup> September 2024 (Week 40 2024).

Data were extracted from the HPSC SARI surveillance database on **18/12/2024**. Data are provisional and subject to ongoing review, validation and update. As a result, figures presented in this report may differ from previously published figures.

**Two of the three** SARI sentinel hospital sites (67%) reported data for the current week (W50 2024). Variations in the number of sentinel sites reporting each week, should be considered when comparing incidence rates and case numbers from previous weeks.

## Key messages

Based on data from two of the three sentinel hospital sites, the number of SARI cases remained stable with 64 cases reported in week 50 compared to 62 cases in week 49. Influenza positivity increased from 20.0% in week 49 to 35.6% in week 50 with increases seen in all age groups. RSV positivity decreased slightly from 20.0% to 18.7%, and SARS-CoV-2 positivity remained at low levels decreasing from 2.9% to 1.7%.

## Summary

- **SARI case numbers and incidence:** 64 SARI cases were admitted to two SARI sentinel sites in week 50 2024, compared to 79 cases from three sites in week 49 2024.
  - **SARI cases <15 years:** 27 cases in week 50 2024 compared to 29 in week 49 2024 (6.9% decrease)
  - **SARI cases ≥15 years:** 37 cases in week 50 2024<sup>1</sup>.
- The incidence rates per 100,000 hospital catchment population were as follows:
  - **All SARI cases:** 24.6 in week 50 2024<sup>1</sup>.
  - **SARI cases <15 years:** 37.3 in week 50 2024, compared to 40.1 in week 49 2024 (6.9% decrease).
  - **SARI cases ≥15 years:** 19.7 in week 50 2024<sup>1</sup>.
- The incidence rates per 1,000 hospital admissions via emergency departments were as follows:
  - **All SARI cases:** 162.0 in week 50 2024<sup>1</sup>.
  - **SARI cases <15 years:** 369.9 in week 50 2024, compared to 537.0 in week 49 2024 (31.1% decrease).
  - **SARI cases ≥15 years:** 114.9 in week 50 2024<sup>1</sup>.

- **Age profile (W50 2024):** 37.5% (n=24) cases were aged ≥65 years and 25.0% (n=16) cases aged <4 years (Table 1):
  - **All SARI cases:** median age 51 years; IQR: 4-76 years.
  - **SARI cases <15 years:** median age 2 years; IQR: 0-5 years.
  - **SARI cases ≥15 years:** median age 75 years; IQR: 59-79 years.
  
- **Underlying medical conditions (W50 2024):** 70.3% (n=45) of cases reported at least one underlying medical condition; among those aged <15 years 37.0% (n=10) and among those aged ≥15 years 94.6% (n=35) of cases reported at least one underlying medical condition (Table 5).
  
- **Virus positivity rate among SARI cases (W50 2024):**
  - Among those tested (92.2%, n=59)
    - 1.7% (n=1) tested positive for SARS-CoV-2, compared to 2.9% (n=2) in week 49 2024
    - 35.6% (n=21) tested positive for influenza (20 A (not subtyped); 1 B), compared to 20.0% (n=14) in week 49 2024
    - 18.6% (n=11) tested positive for RSV, compared to 20.0% (n=14) in week 49 2024
  - See Figures 5, 6a & 6b and Table 2 for further details
  
- **Genomic surveillance (W47 2024-W50 2024):** Among SARI SARS-CoV-2 positive specimens sequenced over the last four weeks (n=3), XEC variant was the most frequent variant identified, accounting for 66.7% (n=2) of samples sequenced, followed by KP.3.1.1, accounting for 33.3% (n=1) of samples sequenced. (Figure 8 & Table 3)
  
- **Vaccination status of SARI cases admitted during the current season (W40 2024-W50 2024)**
  - Amongst SARI cases positive for SARS CoV-2, aged ≥6 months and with known vaccination status (n=12), 83.3% (n=10) had not received a COVID-19 vaccine dose in the six months prior to the reported episode of illness (Table 9).
  - Amongst SARI cases positive for influenza, aged ≥6 months with known vaccination status (n=22), 72.7% (n=16) had not received the 2024/2025 influenza season vaccine prior to the reported episode of illness (Table 10).
  
- **Severe outcomes among SARI cases admitted during the current season (W40 2024-W50 2024)**
  - 3.2% (n=18) of SARI cases were admitted to ICU. The median length of stay was 7 days, IQR 4-15 days (Table 8). Among SARI cases admitted to ICU, 5.6% (n=1) were positive for SARS-CoV-2, 22.2% (n=4) for influenza and 11.1% (n=2) for RSV.
  - 1.5% (n=10) of SARI cases died in hospital. The median age was 79.5 years, IQR 74-87.5 years, 90.0% (n=9) were ≥65 years of age (Table 8)


**Note<sup>1</sup>:** Case number and percentage comparisons are not reported between weeks 49 and 50 2024 due to the difference in number of reporting sites for these weeks.

## Table of contents

About this report .....	1
Key messages .....	1
Summary .....	1
SARI cases and incidence rates .....	4
Demographics .....	6
Laboratory testing for SARS-CoV-2, Influenza and RSV .....	7
PCR testing: .....	7
Influenza typing: .....	9
Genomic analysis: SARS-CoV-2 .....	10
Symptoms .....	11
Underlying medical conditions and risk factors .....	12
Clinical course and outcome .....	13
Complications .....	13
Respiratory support .....	14
Severe outcomes .....	14
Vaccination status .....	15
COVID-19 vaccination status .....	15
Influenza vaccination status .....	16
Links to other national respiratory virus reports .....	17
Appendix .....	18
Technical Notes .....	19
1. SARI Surveillance objectives .....	19
2. Sentinel hospital SARI surveillance sites .....	19
3. Case definition .....	19
4. Denominator data .....	20
5. Laboratory testing .....	20
6. Data collection and reporting .....	20
7. Influenza season .....	21
8. Reference dates .....	21
9. Vaccination status definitions .....	22
Acknowledgements .....	22


## SARI cases and incidence rates

The number of SARI cases admitted per sentinel hospital site by week of admission is displayed in Figure 1, along with the combined SARI incidence rate per 1,000 admissions via emergency department for all hospital sites.




**Figure 1:** Number and incidence of SARI cases per 1,000 hospital admissions via emergency departments, by sentinel hospital site and week of admission, W40 2024-W50 2024 (n=686)

**Note:** UHL data relates to paediatric cases <15 years of age, while SJH and SVUH data relates to adult cases, 15 years of age and older.



**Figure 2:** SARI age-specific incidence rates per 1,000 hospital admissions via emergency departments by week of admission, W40 2024-W50 2024 (n=686)<sup>1</sup>



**Figure 3:** SARI age-specific incidence rates per 100,000 hospital catchment population by week of hospital admission, W40 2024-W50 2024 (n=686)<sup>1</sup>


**Note**<sup>1</sup>: SARI surveillance on those aged under 15 years began in Week 40 2024

## Demographics

**Table 1:** Number and proportion of SARI cases by sex and age, for the current week (W50 2024), last four weeks (W47 2024-W50 2024) and current 2024/2025 season (W40 2024-W50 2024)

Characteristic	Category	Current week W50 2024 N = 64 <sup>1</sup>	Last four weeks W47 2024-W50 2024 N = 269 <sup>1</sup>	Current season W40 2024-W50 2024 N = 686 <sup>1</sup>
Gender	Female	31 (48.4)	120 (44.6)	320 (46.6)
	Male	33 (51.6)	149 (55.4)	366 (53.4)
Age <15 years (in years)	Median (IQR)	2 (0 - 5)	2 (0 - 6)	2 (1 - 5)
	Range	0 - 11	0 - 14	0 - 14
Age ≥15 years (in years)	Median (IQR)	75 (59 - 79)	74 (62 - 81)	72 (59 - 81)
	Range	20 - 95	20 - 99	18 - 101
Age groups (years)	<1	11 (17.2)	29 (10.8)	58 (8.5)
	1-4	7 (10.9)	39 (14.5)	122 (17.8)
	5-14	9 (14.1)	30 (11.2)	72 (10.5)
	15-34	1 (1.6)	7 (2.6)	18 (2.6)
	35-64	12 (18.8)	44 (16.4)	125 (18.2)
	65-79	15 (23.4)	72 (26.8)	172 (25.1)
	80+	9 (14.1)	48 (17.8)	119 (17.3)


<sup>1</sup>n (%)



**Figure 4:** Age- and sex-specific incidence rates per 100,000 hospital catchment population for the current 2024/2025 season (W40 2024-W50 2024)


## Laboratory testing for SARS-CoV-2, Influenza and RSV

### PCR testing:




**Figure 5:** Number of SARI cases PCR positive for SARS-CoV-2, influenza and RSV by week and overall positivity rate for the three pathogens, W40 2024-W50 2024

**Note:** SARI cases are tested, on-site in each hospital site, by PCR for SARS-CoV-2, influenza and RSV on admission.



**Figure 6a:** Percentage of SARI cases PCR positive for SARS-CoV-2, influenza and RSV by week, W40 2024-W50 2024



**Figure 6b:** Weekly positivity rate of SARI cases PCR positive for SARS-CoV-2, influenza and RSV, by age group, W40 2024-W50 2024

**Note:** Y-axis scale may differ for each age group




**Table 2:** Number and proportion of SARI cases PCR positive for SARS-CoV-2, influenza, and RSV for the current week, last four weeks (W47 2024-W50 2024) and season total (W40 2024-W50 2024)

	Current week		Last four weeks		Season total	
	W50 2024		W47 2024-W50 2024		W40 2024-W50 2024	
	<15y N = 27 <sup>1</sup>	≥15y N = 32 <sup>1</sup>	<15y N = 95 <sup>1</sup>	≥15y N = 129 <sup>1</sup>	<15y N = 247 <sup>1</sup>	≥15y N = 278 <sup>1</sup>
SARS-CoV-2	1 (3.7)	0 (0.0)	1 (1.1)	5 (3.9)	4 (1.6)	23 (8.3)
Influenza	9 (33.3)	12 (37.5)	17 (17.9)	28 (21.7)	23 (9.3)	34 (12.2)
RSV	6 (22.2)	5 (15.6)	24 (25.3)	15 (11.6)	33 (13.4)	20 (7.2)

<sup>1</sup>n (%)


A further breakdown of SARI positivity by hospital site is available in the Appendix.

### Influenza typing:



**Figure 7:** Number of SARI cases PCR positive for influenza type/subtype, W40 2024-W50 2024

## Genomic analysis: SARS-CoV-2



\*Includes sub-lineage JN.1, excludes lineage KP.3

**Figure 8:** Number of SARS CoV-2 positive SARI cases sequenced, by week of hospital admission, W40 2024-W50 2024 (n=23)

**Table 3:** Number of SARS CoV-2 positive SARI cases sequenced and reported by Pango lineage and week of hospital admission, for the last four weeks (W47 2024-W50 2024), the preceding four-week period (W43 2024-W46 2024) and the percentage difference in frequency.

Pangolin lineage	W47 2024-W50 2024		W43 2024-W46 2024		% difference <sup>1</sup>
	(n)	(%)	(n)	(%)	
KP.3.1.1	1	33.3	1	9.1	24.2
XEC	1	33.3	3	27.3	6.0
XEC.6	1	33.3	0	0.0	33.3
KP.1.1.3	0	0.0	2	18.2	-18.2
KP.2.15.1	0	0.0	1	9.1	-9.1
LF.7.1.3	0	0.0	1	9.1	-9.1
MC.13	0	0.0	2	18.2	-18.2
XEC.2	0	0.0	1	9.1	-9.1
Total	3		11		

<sup>1</sup> Red indicates  $\geq 5\%$  increase; green indicates  $\geq 5\%$  decrease

**Note:** There is typically a lag time of 1 to 3 weeks between a case being admitted, positive samples selected for sequencing and sequencing being completed.

For further information on circulating variants in Ireland, see [Summary of COVID-19 virus variants in Ireland - Health Protection Surveillance Centre \(hpsc.ie\)](https://hpsc.ie/summary-of-covid-19-virus-variants-in-ireland).

## Symptoms

**Table 4:** Number and proportion of SARI cases' clinical symptoms, either at or prior to hospital admission, for the last four weeks (W47 2024-W50 2024), and current season (W40 2024-W50 2024)

Condition	Last four weeks		Season total	
	W47 2024-W50 2024		W40 2024-W50 2024	
	<15y, N = 98	≥15y, N = 171	<15y, N = 252	≥15y, N = 434
Cough	78 (79.6)	160 (93.6)	184 (73.0)	392 (90.3)
Shortness of breath	45 (45.9)	146 (85.4)	110 (43.7)	365 (84.1)
Fever	79 (80.6)	90 (52.6)	196 (77.8)	222 (51.2)
General deterioration	0 (0.0)	79 (46.2)	0 (0.0)	172 (39.6)
Nausea/Vomiting	27 (27.6)	25 (14.6)	81 (32.1)	50 (11.5)
Malaise	31 (31.6)	21 (12.3)	72 (28.6)	54 (12.4)
Sore throat	28 (28.6)	11 (6.4)	94 (37.3)	31 (7.1)
Muscular pain	0 (0.0)	43 (25.1)	0 (0.0)	91 (21.0)
Diarrhoea	10 (10.2)	14 (8.2)	23 (9.1)	32 (7.4)
Acute confusion	0 (0.0)	15 (8.8)	0 (0.0)	43 (9.9)
Headache	6 (6.1)	11 (6.4)	15 (6.0)	23 (5.3)
Sepsis	0 (0.0)	3 (1.8)	4 (1.6)	13 (3.0)
Ageusia/Dysgeusia/Anosmia	0 (0.0)	0 (0.0)	4 (1.6)	1 (0.2)

**Note:** The following symptoms have been removed from the table, as there are no cases reporting these symptoms in the above time-periods: Apnoea

## Underlying medical conditions and risk factors

SARI cases may be reported with one or more underlying medical conditions, weekly proportions can be based on small numbers and vary from week to week, caution is therefore advised when interpreting changes in weekly proportions (Table 5).

**Table 5:** Number and proportion of SARI cases with underlying medical conditions reported on hospital admission (among those who reported having underlying medical conditions), for the last four weeks (W47 2024-W50 2024), and current season (W40 2024-W50 2024)

Condition	Last four weeks		Season total	
	W47 2024-W50 2024		W40 2024-W50 2024	
	<15y, N = 98	≥15y, N = 171	<15y, N = 252	≥15y, N = 434
Lung disease	1 (1.0)	71 (41.5)	2 (0.8)	183 (42.2)
Heart disease	7 (7.1)	66 (38.6)	12 (4.8)	166 (38.2)
Hypertension	0 (0.0)	61 (35.7)	0 (0.0)	140 (32.3)
Cancer	0 (0.0)	43 (25.1)	0 (0.0)	114 (26.3)
Rheumatological disease	0 (0.0)	40 (23.4)	0 (0.0)	106 (24.4)
Asthma	13 (13.3)	23 (13.5)	34 (13.5)	52 (12.0)
Neurological disease	3 (3.1)	29 (17.0)	9 (3.6)	67 (15.4)
Immunocompromised	0 (0.0)	29 (17.0)	1 (0.4)	71 (16.4)
Diabetes	0 (0.0)	26 (15.2)	1 (0.4)	57 (13.1)
Liver disease	0 (0.0)	15 (8.8)	0 (0.0)	44 (10.1)
Kidney disease	0 (0.0)	11 (6.4)	3 (1.2)	37 (8.5)
Obesity	0 (0.0)	8 (4.7)	0 (0.0)	29 (6.7)
Dementia	0 (0.0)	9 (5.3)	0 (0.0)	26 (6.0)
Intellectual disability	6 (6.1)	2 (1.2)	16 (6.3)	8 (1.8)
Down syndrome	2 (2.0)	0 (0.0)	5 (2.0)	1 (0.2)
Cystic fibrosis	0 (0.0)	0 (0.0)	0 (0.0)	2 (0.5)
Asplenia	0 (0.0)	0 (0.0)	0 (0.0)	2 (0.5)

**Note:** The following conditions have been removed from the table, as there are no cases reporting these conditions in the above time-periods: Long COVID, Tuberculosis

## Clinical course and outcome

### Complications

Information on the clinical course during hospitalisation is only available after patient discharge, and there may be a delay between discharge and data collection, due to the manual data collection methods required. Furthermore, data collection is ongoing for those not yet discharged from hospital.

SARI cases could be reported with one or more complications; among those for whom discharge information is available the most common complication reported was pneumonia (Table 6).

**Table 6:** Number and proportion of SARI cases and complication among discharged SARI cases, for the last four weeks (W47 2024-W50 2024), and current season (W40 2024-W50 2024)

Complication	Last four weeks		Season total	
	W47 2024-W50 2024		W40 2024-W50 2024	
	<15y, N = 95	≥15y, N = 53	<15y, N = 249	≥15y, N = 206
Pneumonia	19 (20.0)	43 (81.1)	43 (17.3)	170 (82.5)
Bronchiolitis	16 (16.8)	0 (0.0)	39 (15.7)	0 (0.0)
Heart failure	0 (0.0)	7 (13.2)	0 (0.0)	35 (17.0)
Acute kidney injury	1 (1.1)	2 (3.8)	2 (0.8)	17 (8.3)
ARDS	6 (6.3)	0 (0.0)	12 (4.8)	1 (0.5)
Sepsis	0 (0.0)	1 (1.9)	4 (1.6)	9 (4.4)
Multi organ failure	0 (0.0)	1 (1.9)	0 (0.0)	2 (1.0)
Other complications	2 (2.1)	1 (1.9)	3 (1.2)	1 (0.5)
No complications	54 (56.8)	10 (18.9)	155 (62.2)	29 (14.1)

**Note:** The following complications have been removed from the table, as there are no cases reporting these conditions in the above time-periods: Myocarditis, Encephalitis, Secondary bacterial infections, PIMS\*

\*Paediatric inflammatory multisystem syndrome

## Respiratory support

Among SARI cases who have been discharged, the highest level of respiratory support received during hospitalisation is described in Table 7.

**Table 7:** Number and proportion of SARI cases by level of respiratory support received, among discharged SARI cases, for the last four weeks (W47 2024-W50 2024), and current season (W40 2024-W50 2024)

Respiratory support	Last four weeks W47 2024-W50 2024		Current season W40 2024-W50 2024	
	<15y, N = 95 <sup>1</sup>	≥15y, N = 53 <sup>1</sup>	<15y, N = 249 <sup>1</sup>	≥15y, N = 206 <sup>1</sup>
No respiratory support given	67 (71%)	22 (42%)	181 (73%)	70 (34%)
Low-flow oxygen therapy	15 (16%)	23 (43%)	38 (15%)	98 (48%)
Non-invasive ventilation	11 (12%)	7 (13%)	26 (10%)	34 (17%)
Invasive ventilation	2 (2.1%)	1 (1.9%)	4 (1.6%)	4 (1.9%)

<sup>1</sup>n (%)

## Severe outcomes

SARI cases are considered to have severe outcomes if they were admitted to ICU and/or died during their hospital stay.

**Table 8:** Number and proportion of SARI cases with severe outcomes, for the last four weeks (W47 2024-W50 2024), and current season (W40 2024-W50 2024)

	Last four weeks W47 2024-W50 2024 N = 269 <sup>1</sup>	Current season W40 2024-W50 2024 N = 686 <sup>1</sup>
Length of stay in hospital (days)		
Median (IQR)	3 (2 - 5)	3 (2 - 6)
Range	1 - 20	1 - 61
Admitted to ICU	10 (4.3%)	18 (3.2%)
ICU length of stay (days)		
Median (IQR)	2 (1 - 9)	7 (2 - 20)
Range	1 - 9	1 - 33
Died in hospital	4 (1.5%)	10 (1.5%)

<sup>1</sup>n (%)

**Note:** Paediatric cases may be reported as admitted to ICU, if transferred to an ICU in a paediatric hospital. However, these cases are excluded from the calculation of length of stay in ICU.

## Vaccination status

Vaccination data are available approximately one week after cases are notified to HPSC, therefore the vaccination status for the current week's SARI cases is recorded as unknown.

### COVID-19 vaccination status

During the current season (W40 2024-W50 2024) among SARI cases PCR positive for SARS-CoV-2, aged  $\geq 6$  months and with known COVID-19 vaccination status ( $n=12$ ), **83.3% ( $n=10$ )** had not received a vaccine dose in the six months prior to the reported episode of illness (Table 9).

**Table 9:** Characteristics of SARI cases positive for SARS-CoV-2 during the current season (W40 2024-W50 2024) by time since last COVID-19 vaccine dose

Characteristic	Category	W40 2024-W50 2024	
		<180 days, N = 2 <sup>1</sup>	$\geq 180$ days, N = 10 <sup>1</sup>
Gender	Female	0 (0.0%)	3 (100.0%)
	Male	2 (22.2%)	7 (77.8%)
Age (years)	Median (IQR)	86 (85 - 86)	79 (71 - 89)
	Range	84 - 87	36 - 94
Age groups (years)	0-14	0 (0.0%)	0 (0.0%)
	15-59	0 (0.0%)	2 (100.0%)
	60-69	0 (0.0%)	1 (100.0%)
	70-79	0 (0.0%)	2 (100.0%)
	80+	2 (28.6%)	5 (71.4%)
Underlying medical conditions	Yes	2 (16.7%)	10 (83.3%)
	No	0 (0.0%)	0 (0.0%)
Patient residence	Residential care facility	0 (0.0%)	0 (0.0%)
	Private residence/home	2 (16.7%)	10 (83.3%)

<sup>1</sup>n (%)

**Note:** Due to small numbers of cases reported as not vaccinated, this group has been included in the  $\geq 180$  days group.

#### Excluded from analysis:

- SARS-CoV-2 positive SARI cases with unknown vaccination status, 13 (48.1%) are excluded.
- SARS-CoV-2 positive SARI cases aged <6 months, 2 (7.4%) are excluded.

## Influenza vaccination status

During the current season (W40 2024-W50 2024) among SARI cases PCR positive for influenza, aged  $\geq 6$  months and with known influenza vaccination status ( $n=22$ ), **72.7% ( $n=16$ )** had not received the 2024/2025 influenza season vaccine prior to the reported episode of illness (Table 10).

**Table 10:** Characteristics of SARI cases positive for influenza during the current season (W40 2024-W50 2024) by vaccination status for the current season's influenza vaccine

Characteristic	Category	W40 2024-W50 2024	
		Vaccinated, N = 6 <sup>1</sup>	Not vaccinated, N = 16 <sup>1</sup>
Gender	Female	3 (27.3%)	8 (72.7%)
	Male	3 (27.3%)	8 (72.7%)
Age (years)	Median (IQR)	76 (72 - 78)	56 (21 - 71)
	Range	62 - 83	7 - 86
Age groups (years)	0-14	0 (0.0%)	4 (100.0%)
	15-59	0 (0.0%)	4 (100.0%)
	60-69	1 (33.3%)	2 (66.7%)
	70-79	4 (57.1%)	3 (42.9%)
	80+	1 (25.0%)	3 (75.0%)
Underlying medical conditions	Yes	6 (31.6%)	13 (68.4%)
	No	0 (0.0%)	3 (100.0%)
Patient residence	Residential care facility	0 (0.0%)	0 (0.0%)
	Private residence/home	6 (27.3%)	16 (72.7%)

<sup>1</sup>n (%)

### Excluded from analysis:

- Influenza positive SARI cases with unknown vaccination status, 30 (53.6%) are excluded
- Influenza positive SARI cases aged  $< 6$  months, 4 (7.1%) are excluded



## Links to other national respiratory virus reports

### Respiratory viruses

- [Integrated Respiratory Virus Bulletin](#)
- [Respiratory Virus Notification Hub](#)

### COVID-19

- [Summary of COVID-19 virus variants in Ireland.](#)
- [National Wastewater Surveillance Programme](#)

## Appendix

**Table A1:** Number of SARI cases, number tested and positivity by hospital site for the current week (W50 2024), previous week (W49 2024) and season total (W40 2024-W50 2024)

	Cases	SARS CoV-2 tested	SARS CoV-2 positive	Influenza & RSV tested	Influenza positive	RSV positive
Site	n	n	n (%)	n	n (%)	n (%)
<b>W50 2024</b>	<b>64</b>	<b>59</b>	<b>1 (1.7)</b>	<b>59</b>	<b>21 (35.6)</b>	<b>11 (18.6)</b>
SVUH	0	-	-	-	-	-
SJH	37	32	0 (0)	32	12 (37.5)	5 (15.6)
UHL	27	27	1 (3.7)	27	9 (33.3)	6 (22.2)
<b>W49 2024</b>	<b>79</b>	<b>70</b>	<b>2 (2.9)</b>	<b>70</b>	<b>14 (20)</b>	<b>14 (20)</b>
SVUH	17	16	1 (6.3)	16	5 (31.3)	2 (12.5)
SJH	33	26	1 (3.8)	26	3 (11.5)	2 (7.7)
UHL	29	28	0 (0)	28	6 (21.4)	10 (35.7)
<b>W40 2024- W50 2024</b>	<b>686</b>	<b>525</b>	<b>27 (5.1)</b>	<b>509</b>	<b>57 (11.2)</b>	<b>53 (10.4)</b>
SVUH	150	147	18 (12.2)	147	13 (8.8)	9 (6.1)
SJH	284	131	5 (3.8)	115	21 (18.3)	11 (9.6)
UHL	252	247	4 (1.6)	247	23 (9.3)	33 (13.4)

## Technical Notes

### 1. SARI Surveillance objectives

Severe acute respiratory infection (SARI) is of major relevance to public health worldwide. Surveillance of SARI is essential to monitor the (co-) circulation of respiratory pathogens and to assess disease severity. Data collected as part of SARI surveillance can provide important early warning information in the context of respiratory disease outbreaks and pandemics. SARI data can also be used as a platform to measure vaccine and antiviral effectiveness and impact. The objectives of SARI surveillance are:

- To describe the number and incidence of SARI cases by aetiology, time, place and person
- To describe and monitor trends, intensity of activity and severity of SARI infections
- To identify groups at risk of severe disease
- To detect unusual and unexpected events
- To assess the SARI burden of disease in the participating hospital
- To assess and monitor vaccine effectiveness

### 2. Sentinel hospital SARI surveillance sites

SARI surveillance was implemented in one tertiary care adult hospital, St. Vincent's University Hospital (SVUH), Dublin on the 5<sup>th</sup> of July 2021. In September 2024 a second tertiary care adult hospital, St James's Hospital (SJH), was included, both sites reporting on SARI cases aged 15 years and older.

A third tertiary care hospital, University Hospital Limerick (UHL), reporting on SARI cases aged under 15 years of age only, began surveillance in September 2024.

### 3. Case definition

SARI cases are identified from new admissions through the Emergency Department, based on clinical symptoms. Patients that develop SARI during their admission, or are admitted through alternate routes, are not included.

*Clinical SARI case:*

The European Centre for Disease Prevention and Control (ECDC) clinical SARI case definition is used for SARI surveillance in Ireland since week 34 2021

SARI case definition: A person hospitalised for at least 24 hours with acute respiratory infection, with at least one of the following symptoms: cough, fever, shortness of breath OR sudden onset of anosmia, ageusia or dysgeusia with onset of symptoms within 14 days prior to hospital admission.

A SARI case refers to an individual patient episode of care

## 4. Denominator data

Denominator data for the hospital catchment area are based on the Census of Population, 2022. The hospital catchment data were prepared and provided by the Health Intelligence Unit (HIU) of the Health Service Executive (HSE) and were extracted from Health Atlas Ireland on 07/05/2024.

Weekly denominator data on all-cause hospital admissions, through the Emergency Department, are provided by the sentinel hospital sites.

## 5. Laboratory testing

SARS-CoV-2, influenza, and RSV PCR testing is carried out on admission.

SARI samples that are positive for SARS-CoV-2 and have a cycle threshold (Ct) value <25 are referred for whole genome sequencing (WGS). The molecular laboratories in SVUH, SJH and UHL are spoke WGS testing sites as part of the national SARS-CoV-2 WGS surveillance programme, for further information please see [Whole Genome Sequencing Programme - Health Protection Surveillance Centre \(hpsc.ie\)](https://hpsc.ie/whole-genome-sequencing-programme). SARI WGS testing is performed on-site at SVUH, SJH and UHL.

Samples that are PCR positive for influenza are sent to the National Virus Reference Laboratory (NVRL) for influenza typing/subtyping/genetic and antigenic characterisation.

## 6. Data collection and reporting

**St Vincent's University Hospital:** Clinical data are collected and managed using REDCap electronic data capture tools hosted at University College Dublin. Laboratory data are extracted from APEX, the laboratory information management system (LIMS), using IBM Cognos software hosted at SVUH.

**St. James's Hospital:** Clinical data are collected and managed on a specifically adapted electronic form within the patient's electronic patient record (EPR). Laboratory data are extracted from Telepath LIMS.

**University Hospital Limerick:** Clinical data are collected manually on the hard copy of the UHL SARI Case Report Form (CRF) and then recorded in the electronic SARI questionnaire on ICNET. Details of laboratory results are obtained from ICNET and are also recorded in the electronic SARI questionnaire on ICNET.

Case-based data are reported by SVUH, SJH and UHL to the HSE Health Protection Surveillance Centre (HPSC) on a weekly basis. Data are also reported by HPSC to ECDC via The European Surveillance System (TESSy) on weekly basis as part of the European SARI surveillance programme.

COVID-19 vaccination data are obtained from the National COVID-19 Vaccination Management System (COVAX) and linked to SARI cases by the HSE-Integrated Information Service (IIS), where data are available.

## 7. Influenza season

The influenza surveillance season runs from week 40 (early October) to week 20 (end of May). During this time, seasonal respiratory viruses usually circulate at higher levels, compared to the summer period (weeks 21 to 39). The seasonal comparisons used in this report refer to the influenza surveillance season.

## 8. Reference dates

### SARI Surveillance

05/07/2021 (Week 27 2021) – commenced of SARI surveillance at first sentinel hospital site  
30/09/2024 (Week 40 2024) - commenced SARI surveillance at the second and third sentinel sites

### Vaccination campaign

27/09/2021 (Week 39 2021) – first COVID-19 booster vaccination campaign commenced  
22/04/2022 (Week 16 2022) – second COVID-19 booster vaccination campaign commenced  
03/10/2022 (Week 40 2022) – Autumn 2022 COVID-19 booster vaccination campaign commenced  
28/04/2023 (Week 17 2023) – Spring 2023 COVID-19 booster vaccination campaign commenced  
02/10/2023 (Week 40 2023) – Autumn 2023 COVID-19 booster vaccination campaign commenced  
22/04/2024 (Week 17 2024) – Spring 2024 COVID-19 booster vaccination campaign commenced  
30/09/2024 (Week 40 2024) – Autumn 2024 COVID-19 booster vaccination campaign commenced

### Winter respiratory virus seasons

04/10/2021 (Week 40 2021) - start of the 2021/2022 season  
03/10/2022 (Week 40 2022) - start of the 2022/2023 season  
02/10/2023 (Week 40 2023) - start of the 2023/2024 season  
30/09/2024 (Week 40 2024) - start of the 2024/2025 season

Week number refers to the week of hospital admission. Weeks are from Monday to Sunday, as per the international ISO week<sup>1</sup>.

---

<sup>1</sup> Monday to Sunday (ISO week) used as per ECDC/WHO/International reporting protocol.

## 9. Vaccination status definitions

For the purposes of SARI surveillance, vaccination status of cases is as follows:

**Vaccinated COVID case:** A confirmed case of COVID-19 who received any dose of a COVID-19 vaccine,  $\geq 14$  days before onset of symptoms.

**Unvaccinated COVID-19 case:** A confirmed case of COVID-19 who did not receive any dose of a COVID-19 vaccine i.e. was never vaccinated.

**Time since vaccination:** For a vaccinated COVID-19 case, this is the time between the date of last dose vaccination and the date of symptom onset and categorised as  $< 180$  days or  $\geq 180$  days since vaccination.

**Vaccinated influenza case:** A confirmed case of influenza will be considered as vaccinated against influenza if they received one dose of the influenza vaccine as part of the current season's influenza vaccination campaign  $\geq 14$  days before onset of symptoms.

**Unvaccinated influenza case:** A confirmed case of influenza will be considered as unvaccinated if they did not receive an influenza vaccine as part of the current season's influenza vaccination campaign or if they were vaccinated after onset of symptoms.

**Vaccine status unknown:** The SARI patient is reported on the SARI hospital clinical questionnaire as vaccinated, however there is no identifiable linked record of COVID-19 vaccination and/or influenza vaccination on the National Immunisation system. Vaccination status is reported as unknown, until verified on the National Immunisation system.

## Acknowledgements

Sincere thanks are extended to all those who participate in SARI surveillance, including those in St. Vincent's University Hospital, St James's Hospital, University Hospital Limerick, the UCD Clinical Research Centre and the National Virus Reference Laboratory. Thanks to members of the HSE Integrated Information Services (IIS) for work on the SARI-COVAX data linkages.

This report was produced by the SARI Surveillance Team at HPSC, using R studio software.