



# Public Health & Infection Prevention & Control Guidelines on the Prevention and Management of Cases and Outbreaks of COVID-19, Influenza & other Respiratory Infections in Residential Care Facilities

V1.1 30.11.2021

Note: If you have any queries on this guideline please contact the AMRIC team at [hse.amrteam@hse.ie](mailto:hse.amrteam@hse.ie)

Ver.	Date	Changes from previous version
1.1	30.11.2021	<p>Editorial changes and updated to reflect current status of vaccination programme</p> <p>Explicit statement re risk assessment for staff who decline vaccination (in addition to External Contractors)</p> <p>Update to reflect Government policy on restriction of movement for fully vaccinated household close contacts</p> <p>Update on derogation from exclusion of contacts from work</p> <p>Increased emphasis on testing in the context of transfers and admissions of fully vaccinated people</p> <p>Rewording to emphasise that healthcare workers must have access to respirator masks at all times when caring for people with suspected or confirmed COVID-19</p> <p>Reintroduction of requirement for testing for fully vaccinated people before transfer or admission to LTRCF</p>
1.0	26.10.2021	<p>This document replaces Version 6.7 of "COVID-19 Interim Public Health &amp; Infection Prevention and Control Guideline on the prevention and management of cases and outbreaks of COVID-19 in Residential Care Facilities"</p>

*All guidance should be read and interpreted in conjunction with the [Government's Framework of Restrictions](#)*

### **Acknowledgements**

The following guidance documents were referred to in developing this guidance:

- Coronavirus Disease 2019 (COVID-19) Infection Prevention and Control Guidance including Outbreak Control in Residential Care Facilities developed by the Communicable Diseases Network Australia (CDNA)
- COVID-19: Information and Guidance for Social or Community Care & Residential Settings Health Protection Scotland
- Public Health Guidelines on the Prevention and Management of Influenza Outbreaks in Residential Care Facilities in Ireland 2019/2020
- World Health Organization. Infection Prevention and Control Guidance for long-term care facilities in the context of COVID-19: interim guidance, 21 March 2020 World Health Organization; 2020
- HIQA-Rapid Review of Public Health guidance on infection prevention and control measures for residential care facilities in the context of COVID-19 30/30/20

## Table of Contents

1	Introduction .....	5
2	Roles and responsibilities.....	6
2.1	Residential Care Facility .....	6
2.2	Regional Department of Public Health .....	8
3	COVID-19 Background information .....	8
3.1	Sources of Infection with COVID-19.....	8
3.2	Routes of Transmission .....	9
3.3	Control of Transmission .....	11
3.4	Vaccination.....	12
3.5	Incubation period for COVID-19 .....	13
3.6	Survival of Respiratory Virus in the environment.....	13
3.7	Clinical features of COVID-19 and Influenza .....	14
3.8	Laboratory testing.....	15
3.9	COVID-19 and Immunity after Recovery.....	17
3.10	Planning.....	17
3.11	Education .....	18
3.12	Social activity, physical distancing measures & Pods .....	20
3.13	Group Activities.....	21
3.14	Group Controls to minimise risk of inadvertent introduction of virus .....	22
3.15	Increased surveillance and early identification of cases of COVID-19 infection .....	27
4	Management of an outbreak of COVID-19 or Influenza.....	28
4.1	Declaring an outbreak.....	29
4.2	Outbreak Control Team (OCT) .....	30
4.3	Management of a possible or confirmed case of COVID-19 or Influenza.....	32
4.4	Cohorting residents with possible or confirmed COVID-19 or Influenza.....	34
4.5	Management of close contacts of a possible or confirmed case of COVID-19.....	36
4.6	Management of close contacts of a possible or confirmed case of Influenza.....	37
4.7	Infection prevention and control measures .....	38
4.7.1	Standard precautions.....	38
4.7.2	Hand hygiene .....	38
4.7.3	Respiratory hygiene and cough etiquette.....	39
4.7.4	Personal Protective Equipment (PPE).....	40
4.7.5	Transmission-based Precautions for COVID-19 and Influenza virus.....	41
4.8	Duration of transmission based precautions.....	42

4.8.1	Care Equipment .....	43
4.8.2	Management of blood and body fluid spillages.....	43
4.8.3	Management of waste .....	43
4.8.4	Safe management of linen (laundry) .....	44
4.8.5	Environmental hygiene .....	45
4.8.6	Routine cleaning in the context of COVID-19 and Influenza .....	45
4.8.7	Frequency of cleaning in the context of COVID-19 and Influenza .....	46
4.8.8	Terminal cleaning.....	46
4.8.9	Staff uniforms/clothing .....	47
4.9	Communication.....	47
4.10	Support services for staff and residents .....	48
5	Care of the person with suspected of confirmed COVID-19 or a Contact of COVID-19 who is dying	48
6	Care of the recently deceased .....	49
6.1	Hygienic preparation.....	49
6.2	Handling personal possessions of the deceased .....	49
6.3	Transport to the mortuary.....	50
7	Monitoring outbreak progress.....	50
8	Declaring the outbreak over .....	51
	Appendix A: Prevention and control of outbreaks of COVID-19 in RCF .....	52
	Appendix B: Details for line listing .....	54
	Appendix C: Part 1 – Respiratory outbreak line listing Form – Residents ONLY*.....	55
	Appendix C: Part 2 –Residents ONLY .....	56
	Appendix C: Part 3 – Respiratory outbreak line listing form – Staff ONLY* .....	57
	Appendix C: Part 4 –Staff ONLY* .....	58
	Appendix D: Checklist for outbreak management.....	59
	Appendix E Hand Hygiene poster .....	61
	Appendix F Donning and Doffing PPE .....	62
	Appendix G Transmission based precautions.....	63
	Appendix H Admissions, transfers and discharges to and from residential care facilities .....	64

## 1 Introduction

This document replaces version 1.0 of this document. It seems increasingly likely that the winter of 2021/2022 will be characterised by circulation of respiratory viruses other than SARS-CoV-2 including Influenza virus. The clinical features caused by infection with respiratory viruses are often difficult to differentiate and the public health and infection prevention and control management is very similar. For these reasons this document is framed as general guidance for this group of infections.

Managing the risk of COVID-19, Influenza and other respiratory viruses in a residential care setting can be thought of as three elements. The first is to take all practical measures to reduce unintended introduction of the virus into the residential care facility. If the virus is not introduced by a person with infection, then it cannot spread. Even when all practical precautions are taken it is still possible that the virus will be introduced unintentionally, therefore the second element is to take all practical measures to reduce the risk of the virus spreading if introduced. The third element is having processes in place to minimise the risk of harm to residents and staff if both other elements fail and the virus is introduced and spreads. This guideline addresses measures needed to achieve all of the above elements. Controlling the risk of introduction, spread and harm from COVID-19 and Influenza is challenging particularly as there is a need to balance the management of risk with respect for the autonomy and rights of residents. Vaccination of residents and staff plays a central part in managing all aspects of the risk.

Please note that experience and the evidence base related to COVID-19 are increasing rapidly. Therefore, it is essential that you confirm that you are using the latest version of guidance.

<https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/residentialcarefacilities/>

Application of this guidance document should take account of the current level of COVID-19 and Influenza disease in the country and in the specific locality at any time.

<https://www.gov.ie/en/campaigns/resilience-recovery-2020-2021-plan-for-living-with-covid-19/>

This guidance is supplemented by a checklist for RCFs to aid implementation of this guidance.  
The checklist is available at the following link

[hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/residentialcarefacilities](https://hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/residentialcarefacilities)

## **2 Roles and responsibilities**

### **2.1 Residential Care Facility**

This guidance applies to residential care facilities (RCF) where residents are provided with overnight accommodation. The anticipated duration of such accommodation may vary within and between different types of RCF. For example, some RCFs for older persons may offer a blend of long-term nursing home and shorter-term respite and convalescence care.

This guidance was developed primarily for congregated care settings providing care for relatively large numbers of residents who are at high risk or very high risk of severe disease. Experience shows that spread of COVID-19 in these settings had profound consequences prior to the vaccination campaign and continues to impact some residents severely. Although Influenza was not observed in RCFs in Ireland in the winter of 2020/2021 there is reason to expect that it will circulate this year and the potential impact on residents is clear from previous years. While the principles of this guidance can be applied in all residential care settings, the risks are lower in the context of residential care provided in the setting of community housing for groups of five to six people or fewer. In that context, a pragmatic approach is required and in particular restricting people to their room for extended periods is likely to be impractical and should only be considered in the context of very specific risk.

Facilities providing acute inpatient rehabilitation services are advised to refer to the 'Acute Hospital Infection Prevention and Control Precautions for Possible or Confirmed COVID-19 in a Pandemic Setting':

<https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/>

The primary responsibility for managing the risk of infection with COVID-19 and Influenza and for control of outbreaks lies with the RCF, within their responsibilities for resident care and infection prevention and control (IPC). This responsibility is referred to in the 2016 National Standards for Residential Care Settings for Older People in Ireland. The 2018 National Standards for infection prevention and control in community services are also relevant. All RCFs should have in-house IPC expertise and should have outbreak management plans in place.

**Congregated care settings, such as nursing homes, should have at a minimum one designated on-site IPC link practitioner who has protected time and the support of management to promote good IPC practice within the facility. An IPC link practitioner generally does not have a formal IPC qualification but should be supported in participating in link practitioner training at the earliest opportunity and avail of ongoing training as much as possible. The IPC link-practitioner should provide ongoing training to staff with a particular emphasis on Standard Precautions including hand hygiene, respiratory hygiene, cough etiquette and environmental cleaning.**

[Under the Infectious Diseases Regulations 1981, Amendment February 2020](#), any medical practitioner who is aware of a case of COVID-19 or an outbreak of any infectious disease including influenza, is obliged to notify the Medical Officer of Health (MOH) at the regional Department of Public Health. Contact details can be found [here](#) on the HPSC website.

Registered providers must notify the Chief Inspector (HIQA) of an outbreak of a notifiable disease within three working days, (Statutory Notifications Guidance for registered providers and persons in charge of designated centres. January 2016).

## 2.2 Regional Department of Public Health

The Regional Departments of Public Health are responsible for investigating cases and outbreaks of COVID-19, influenza or other infectious disease and providing overall leadership and oversight for outbreak management. The IPC link practitioner is a key resource in supporting the Public Health Department in fulfilling its role.

## 3 COVID-19 Background information

### 3.1 Sources of Infection with COVID-19

COVID-19 and Influenza virus infection is acquired as a result of exposure to a person shedding infectious virus. It is generally accepted that the highest risk of transmission occurs at about the time an infected person develops symptoms. Spread from **symptomatic people** is generally considered to be the greatest risk.

With respect to COVID-19 it is accepted that infection can be transmitted from people with minimal symptoms, from people before they develop symptoms (**pre-symptomatic transmission**) and from people who never develop symptoms (**asymptomatic transmission**). However, symptomatic people are generally more infectious. HIQA have provided a useful summary of the evidence related to asymptomatic transmission at:

<https://www.hiqa.ie/reports-and-publications/health-technology-assessment/evidence-summary-asymptomatic-transmission>

### Transmission in the Healthcare Setting

The spread of COVID-19 and Influenza in the healthcare setting is a specific concern. Experience in Ireland and elsewhere indicates that transmission in residential care facilities and hospitals can occur readily when the virus is introduced from the community into the healthcare setting. Even with high levels of vaccination the virus can spread rapidly particularly if IPC precautions are suboptimal. Transmission typically occurs when an unrecognised infectious person enters the facility. Control of entry to minimise risk of unrecognised introduction is therefore a key priority in preventing outbreaks. This requires a particular focus when rates of infection in the



community served are high. In the context of long-term residential care facilities, the key group of people who move regularly between community and the facility is staff. Visitors also represent a risk of introduction of COVID-19 and influenza. Guidance on managing risk associated with visiting is provided on the HPSC website.

Outbreaks of infection involving both residents and healthcare workers (HCW) have been frequent in RCFs during the major community surges in COVID-19. This has also been observed previously with influenza. The control of spread of the SARS-CoV-2 virus, the cause of COVID-19, in RCFs in this context was very challenging prior to vaccination and continues to be challenging even after vaccination. The emergence of SARS-CoV-2 variants with higher transmissibility or that are less effectively prevented by vaccination remains an ongoing concern.

Vaccination, including booster vaccination, of a high proportion of residents and staff in RCFs against COVID-19 has had a major impact on reducing the impact of COVID-19 in RCFs. In this context, it is possible to manage the risk of spread of COVID-19 effectively with less restriction on the lives of residents. There is however, a continuing need for vigilance to prevent infectious staff members or other people from entering the RCF. It remains important to ensure that patients with symptoms of COVID-19 and influenza are detected promptly and that transmission-based IPC precautions, including appropriate use of PPE, are implemented in the care of infectious residents to further reduce the risk of spread. RCFs must have systems in place to ensure that, to the greatest extent possible; residents with COVID-19 and influenza are rapidly identified and are cared for with appropriate transmission-based IPC precautions.

### **3.2 Routes of Transmission**

The transmission of COVID-19 and influenza occurs mainly as a result of scattering of liquid respiratory particles into the air from an infectious person. Respiratory particles are generated from the nose and mouth by actions such as breathing, coughing, sneezing, talking or laughing. Transmission to others may result from direct impact of infectious droplets on the mucosa of persons nearby and through contact with surfaces contaminated with infectious respiratory

droplets and subsequent transfer of infectious material to the mucous membranes (droplet transmission).

With respect to COVID-19 the World Health Organisation (WHO) states that *“Recent clinical reports of health workers exposed to COVID-19 index cases, not in the presence of aerosol-generating procedures, found no nosocomial transmission when contact and droplet precautions were appropriately used, including the wearing of surgical masks as a component of the personal protective equipment (PPE). These observations suggest that aerosol transmission did not occur in this context. Further studies are needed to determine whether it is possible to detect viable SARS-CoV-2 in air samples from settings where no procedures that generate aerosols are performed and what role aerosols might play in transmission.”* <https://www.who.int/news-room/commentaries/detail/transmission-of-sars-cov-2-implications-for-infection-prevention-precautions> (accessed 29 May 2021). Of note this Scientific Brief was published in July 2020 prior to the emergence of the alpha or delta variants.

Longer range transmission of SARS-CoV-2 (airborne transmission) has been acknowledged since the start of the pandemic, in particular in the context of certain medical procedures referred to as aerosol generating procedures associated with an increased risk of infection (AGPs). More recently, in the context of the emergence and spread of more infectious SARS-CoV-2 virus variants such as the alpha and delta variants the risk of airborne spread has increased. This is reflected in the April 2021 update to the WHO website Frequently Asked Questions as follows. *“The virus can spread from an infected person’s mouth or nose in small liquid particles when they cough, sneeze, speak, sing or breathe. These particles range from larger respiratory droplets to smaller aerosols. Current evidence suggests that the virus spreads mainly between people who are in close contact with each other, typically within 1 metre (short-range). A person can be infected when aerosols or droplets containing the virus are inhaled or come directly into contact with the eyes, nose, or mouth. The virus can also spread in poorly ventilated and/or crowded indoor settings, where people tend to spend longer periods of time. This is because aerosols remain suspended in the air or travel farther than 1 metre (long-range). People may also become infected by touching surfaces that have been contaminated by the virus when touching their eyes, nose or*

*mouth without cleaning their hands.”* <https://www.who.int/news-room/q-a-detail/coronavirus-disease-covid-19-how-is-it-transmitted> (accessed 29 May 2021).

Recent experience in hospitals in Ireland also highlights particular concerns regarding spread of SARS-CoV-2 infection over longer distances from infectious patients supported by high flow oxygen devices (which is a recognised AGP) in particular in multi-bed areas.

Higher levels of SARS-CoV-2 virus have been detected in patients with severe illness compared to mild cases. Peak levels of SARS-CoV-2 and influenza virus are generally found around the time of symptom onset but people may be infectious before they develop symptoms (pre-symptomatic spread) and some people who never notice symptoms may be infectious (asymptomatic spread).

### **3.3 Control of Transmission**

#### **Vaccination**

There is evidence that vaccination against SARS-CoV-2 reduces the risk of transmission in addition to preventing severity of disease in those vaccinated. This serves to emphasise the importance of vaccination, including booster vaccination, of healthcare workers not only to protect themselves but also in protecting the people that they care for. Vaccination against influenza is likewise important for protection of the healthcare worker and the people they care for.

#### **Standard Precautions**

In addition to vaccination, Standard Precautions applied to all people cared for in all settings at all times plays a key role in managing the risk of infection for residents and for healthcare workers in every long-term residential care facility. For further information on Standard Precautions please see Interim Guidance on Infection Prevention and Control for the HSE (2020). Note the recommendations of the National Public Health Emergency Team (NPHE) on use of surgical masks in healthcare are reflected in the HSE guidance.

#### **Transmission-based Precautions**

Transmission-based Precautions are measures taken in addition to Standard Precautions to manage risk of transmission of infection when caring for people with known or suspected

infectious disease for which Standard Precautions alone are not sufficient. Transmission-based Precautions include Contact, Droplet and Airborne Precautions. For details on Transmission-based precautions, please see the Interim Guidance on Infection Prevention and Control for the HSE (2021) available at:

<https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/hseinfectionpreventionandcontrolguidanceandframework/>

### **3.4 Vaccination**

Vaccination for COVID-19 began in Ireland in late December 2020. The programme has been highly successful, with uptake of vaccines in adults at greater than 90%. The vast majority of residents and staff in RCFs for older people have now been offered vaccination. The vast majority of residents in RCFs have been vaccinated. A programme to administer booster doses has been implemented and additional doses are being administered to immunocompromised people aged 12 and older.

Vaccination offers a high degree of protection against severe disease to residents and healthcare workers when they are fully vaccinated and have had booster vaccine. There is also evidence that vaccination reduces the risk of transmission of virus by people who are vaccinated.

Individuals are considered to be fully vaccinated as follows:

1. 15 days after the second dose of AstraZeneca (Vaxzevria);
2. 7 days after the second Pfizer-BioNTech dose (Comirnaty);
3. 14 days after the second Moderna dose (Spikevax);
4. 14 days after Janssen (one dose vaccination course).

If other COVID-19 vaccines become available the requirement for vaccination will be as advised by HSE.ie.

RCFs should ensure that there is a risk assessment process in place for staff who decline vaccination. This should be equivalent to that which the HSE applies to HSE staff.

Influenza vaccine is recommended to and offered to residents of RCFs and to all healthcare workers before winter each year.

When people have had vaccination the vaccine will confer a high degree of protection from severe disease. However, vaccine protection is not perfect and the vaccine may not work so well in people who have certain conditions or who are on a treatment that interferes with their immune system. In the context of potentially intense exposure related to caring for a person with infectious COVID-19 or influenza, vaccine does not eliminate risk of infection. Even fully vaccinated healthcare workers and residents are advised to continue to adhere to all IPC measures in this guideline.

### **3.5 Incubation period for COVID-19**

Current estimates suggest that the time between exposure to the virus and developing symptoms (incubation period) is from five to six days for most people, but can range from one to 14 days. Individuals are usually considered most infectious to others around the time they develop symptoms. How infectious an individual is and how long they remain infectious is related to some degree to the severity and stage of illness and may be influenced by the immune function of the individual.

### **3.6 Survival of Respiratory Virus in the environment**

Survival on environmental surfaces depends on the type of surface and the environmental conditions. One study using a SARS-CoV-2 strain showed that it can survive on plastic for up to 72 hours, for 48 hours on stainless steel and up to eight hours on copper when no cleaning is performed. However, the levels of virus declined very quickly over the time period. Common household cleaning products and many disinfectants, including bleach, easily kill SARS-CoV-2 and

influenza virus. Based on experience through the pandemic, infection as a result of persistence of viable virus on surfaces for long periods appears to be very uncommon.

### **3.7 Clinical features of COVID-19 and Influenza**

Most otherwise healthy people with COVID-19 or influenza will have mild disease and will recover. A minority will develop more serious illness.

The HSE has defined categories of people who are considered very high risk for COVID-19 (also known as extremely medically vulnerable) and those at high risk for severe disease. See the following link <https://www2.hse.ie/conditions/coronavirus/people-at-higher-risk.html>

For more information on symptoms and signs of COVID-19, refer to the latest case definition <https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/casedefinitions/>

And on <https://www2.hse.ie/conditions/covid19/symptoms/overview/>

**It is important to remember that older people with COVID-19 very often do not have fever and respiratory symptoms and may only have symptoms such as:**

- 1. lethargy;**
- 2. increased confusion;**
- 3. change in baseline condition;**
- 4. loss of appetite.**

Clinical judgement with a high index of suspicion should be used when assessing residents.

**It is important to note that people who are fully vaccinated and have had their booster and who become infected may have very mild symptoms but be infectious. This poses a risk in particular for residents who are not vaccinated or who may not have had a good response to vaccine.**

RCF residents with influenza virus infection may have very similar clinical features to residents with COVID-19. It is important to consider and test for both viruses when clinically relevant.

### 3.8 Laboratory testing

See also Guidance on testing for Acute Respiratory Infection in Residential Care Facilities – Winter 2021/2022 at [www.hpsc.ie](http://www.hpsc.ie)).

1. Laboratory testing is necessary to confirm a diagnosis of COVID-19 or influenza infection;
2. Testing is performed in a similar way for both viruses. Note that if a resident has symptoms of viral respiratory tract infection they should generally be tested for both SARS-CoV-2 and Influenza virus;
3. Check with the laboratory who provides the service in advance of any requirement to test that you have the correct type of swab required by that laboratory
4. The viral swab collected may be deep nasal/mid-turbinate swab or a nasopharyngeal swab. A deep nasal swab is often less distressing for people and is almost equally likely to detect the virus if it is present. Deep nasal sampling is particularly useful where people are undergoing repeated testing or are distressed by nasopharyngeal sampling. Please note a HSE video demonstrating the sample collection technique is available at the following link <https://bit.ly/3efCPO2>;
5. Anterior nasal swabs are not a good sample and should not be submitted.
6. **When testing is performed, ensure the correct swab type is taken (viral swab), sealed tightly to prevent leakage and is appropriately labelled** with two matching resident identifiers on both the swab and request form, to include the resident's name and date-of-birth (DOB). Ensure that the name and contact details for the resident's doctor are on the request form, together with the address of the RCF and any other contact details required. These should include the name and telephone number (mobile preferably) for the designated person who will receive the laboratory result clearly visible on the request form. Deliver the sample to the testing laboratory as soon as possible. Confirm in advance that you are sending the sample to the designated laboratory to perform the test for your RCF and that samples taken from residents of RCF are being prioritised for

testing, particularly in a suspected outbreak. Indicate clearly if testing for Influenza virus is also required;

7. Current PCR based laboratory tests are accurate, but no diagnostic test is perfect. If a test result comes back as “SARS-CoV-2 not detected” and “Influenza virus not detected” and the resident remains unwell with no alternative diagnosis, then a diagnosis of COVID-19 or Influenza is still possible. If there is any concern, the resident’s condition should be discussed with their doctor;
8. Additional information is available in the section on Duration of Transmission Based Precautions;
9. Testing for respiratory viruses including SARS-CoV-2 and influenza virus by another method called **antigen testing** is also used in some situations. Antigen testing is generally less likely to detect virus at low levels than PCR testing.



### 3.9 COVID-19 and Immunity after Recovery

There is still limited experience with immunity after recovery and therefore caution is required in interpretation. In general, people who have recovered from COVID-19 have evidence of an immune response and that offers them significant protection, at least in the short-term (up to nine months following recovery). However, it is recommended that healthcare workers who have recovered from COVID-19 continue to follow the same IPC precautions as all other HCWs when in contact with patients to reduce the risk of transmission of COVID-19.

Currently, antibody testing is not recommended for routine use to assess immunity to infection

### General measures to prevent a COVID-19 and Influenza outbreaks

#### 3.10 Planning

1. Identify a lead for COVID-19 and Influenza preparedness and response in the RCF. The lead should be a person with sufficient authority to ensure that appropriate action is taken and requires at a minimum the support of one designated on-site IPC link practitioner (see above). In some smaller RCFs the lead may also fulfil the role of the IPC link practitioner. In larger RCFs there may be a requirement for a liaison person on each unit in the RCF in addition to lead and link IPC practitioner roles;
2. RCF settings must have COVID-19 and influenza preparedness plans in place to include planning for cohorting of potentially infectious residents separate from non-infectious residents, enhanced IPC, staff training, establishing surge capacity and promoting resident and family communication;
3. Maintain an up-to-date line list of all residents in the RCF and all staff working in the RCF, along with contact telephone numbers;
4. Each RCF should have an area identified where a resident with suspected or confirmed COVID-19 or influenza could be isolated;
5. Where possible, each ward or floor should try to operate as a discrete unit or zone, meaning that staff and equipment are designated to a specific area and are not rotated from other areas (this includes night duty). This practice may reduce exposure to risk for staff and residents in the event COVID-19 or influenza is introduced into the facility. This

may also allow outbreak response measures to be targeted in zones, rather than having to be implemented facility-wide;

6. The risk associated with movement or rotation of staff is lower if staff are fully vaccinated and have had booster vaccine against COVID-19 and influenza or are in the nine month period after they have had and recovered from COVID-19;
7. Facilities should ensure the availability of supplies, including tissues, alcohol-based hand rub (ABHR), hand wipes, cleaning products, disinfectants and personal protective equipment (PPE) and liaise with relevant supply lines if there is difficulty in obtaining such supplies;
8. Supplies of PPE should be sufficient to ensure that single-use items of PPE, including visors and goggles, are used only once and then disposed of safely;
9. Note: that the Health and Safety Authority indicate that where a risk assessment indicates that workers need to use a close-fitting respirator mask for their protection that every effort should be made to comply with the requirement for fit testing of the workers, as far as is reasonably practicable. When fit testing of all staff is not immediately possible, then fit testing should be prioritised for those at greatest risk;
10. A summary table of key interventions for the prevention and management of a COVID-19 or Influenza outbreak can be found in Appendix A.

### **3.11 Education**

#### **3.11.1 Staff**

1. All staff should be aware of the early signs and symptoms of COVID-19 and influenza. They should know who to alert if they have a concern. Staff should be able to contact an appropriate escalation pathway 24/7. Please see the HPSC website for the most up to date case definition for COVID-19;
2. All staff should have training in standard precautions, in particular hand hygiene, respiratory hygiene and cough etiquette, along with training in transmission-based

precautions (contact, droplet and airborne), including the appropriate use of PPE for each situation;

3. RCFs should ensure that one or more staff members are trained to collect a viral swab sample for testing for SARS-CoV-2 and Influenza virus. Please refer to guidelines and video in relation to same available [HERE](#).

### **3.11.2 Residents**

1. Residents should be consulted on and kept informed of the measures being taken and the reason for these measures during this time. This is particularly important where visiting has been restricted;
2. Residents should be encouraged and facilitated to clean their hands and actively assisted with this practice where necessary;
3. Key messages around cough etiquette (where appropriate) include:
  - a. Cover your mouth and nose with a disposable tissue when coughing and sneezing to contain respiratory secretions;
  - b. Discard used tissues after use and clean your hands;
  - c. If you don't have a tissue, cough into your forearm or the crook of your elbow;
  - d. Clean your hands.
4. In line with guidance from the National Public Health Emergency Team (NPHE), the importance of maintaining a physical distance from others in accordance with Public Health Guidance where possible should be observed. However, if two residents who are fully vaccinated against COVID-19 are visiting with each other in one room they do not need to wear a mask or maintain distance;
5. Where possible and appropriate, residents should be made aware of the need to report any new symptoms of illness to staff members;
6. Residents who may leave a RCF should be made aware of the general principles of staying well, details of which can be found <https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/peopleatincreasedriskofsevereilln>

[ess/othersatincreasedriskofsevereillness/Guidance%20For%20Older%20People%20and%20Others-Reducing%20COVID-19%20Exposure%20Risk.pdf](#) ;

7. Residents who are in the high risk or the very high risk groups for severe disease with COVID-19 should be supported in taking additional measures to reduce their risk of infection over and above any general measures applied in the RCF if they wish to do so;
8. Residents may wish to wear a mask at certain times or in certain places within the LTRCF, and this should be facilitated. In addition, if tolerated, residents who are not vaccinated should be encouraged to wear a mask in busy areas of the RCF or during transport to and from the facility.

### **3.12 Social activity, physical distancing measures & Pods**

1. Social activity is an essential part of community life within the LTRCF and should not be suspended or limited unless there is a specific infection prevention and control or public health reason not to do so. If social activity is suspended or limited this should be based on a documented risk assessment;
2. Social activity is associated with some risk of spread of infection but it is important to the overall wellbeing of residents;
3. Residents with symptoms of COVID-19, influenza or other viral respiratory tract infection should be asked not to join in social activities until they are no longer infectious. **This continues to apply to people who have been fully vaccinated;**
4. Through the COVID-19 pandemic it has been advised that social activity should be organized on the basis of limiting contact to consistent groups of residents to the greatest extent practical through organization of pods of 4 to 6 people. Stepwise movement to larger groups should be take account of booster vaccination and current intensity of community transmission;
5. Residents engaged in social activity should be encouraged to practice hand hygiene and cough etiquette. Residents engaged in social activity should also be advised to limit direct contact with other people other than partners;
6. In order to support physical distancing, mealtimes may need to be staggered to ensure that distance between people is maintained when dining;

7. In the context of social interaction it is appropriate, with due regard to the weather and comfort, to use well-ventilated indoor space or outdoor space where available;
8. Staff members should also be required to maintain physical distancing measures during their break and meal times;
9. Table 2 of Section 4.4 provides some suggestions on particular group activities.

### 3.13 Group Activities

1. See details above in section 4.3. Suggestions on specific activities are outlined in Table 2;
2. Before any group activity confirm on that day that participants have no symptoms that suggest viral infection (COVID-19 or influenza);
3. Weather permitting, outdoor group activities are likely to be lower risk than indoor activities;
4. Ensure adequate supplies of hand sanitiser and appropriate cleaning products (for example detergent wipes) are available in each activity room/area;
5. Ensure staff and volunteers know that they should be fully vaccinated and wear a surgical face mask when they cannot maintain physical distance from residents and that they should perform hand hygiene regularly especially after assisting a resident.

**Table 2 Examples of group activities**

	<b>Comment</b>
<b>Chair aerobics/ yoga</b>	Clean chairs between each session with detergent wipes
<b>Ball games</b>	Hand hygiene before and after
<b>Bingo</b>	Individual answer sheets & markers
<b>Dancing</b>	Dancing alone with distance maintained is very low risk – if dancing with partners if limited to one consistent partner from the same pod this can reduce risk
<b>Card games e.g. bridge</b>	Hand hygiene before and after
<b>Computer skills</b>	Cover keyboard and clean after use
<b>Sing songs</b>	Maintain as much distance and ventilation as practical and keep groups small
<b>Knitting</b>	Each person uses their own equipment
<b>Art</b>	Each person uses own equipment or equipment cleaned between users
<b>Flower arranging</b>	Hand hygiene before and after
<b>Table quiz</b>	Hand hygiene before and after
<b>Jigsaws</b>	Hand hygiene before and after
<b>Pottery</b>	Clean potter's wheel between users
<b>Films</b>	Physical distancing
<b>Wii Keep fit</b>	Clean hand set and controls between residents

### 3.14 Group Controls to minimise risk of inadvertent introduction of virus

#### 3.14.1 Staff

1. Vaccination of staff against SARS-CoV-2, including booster vaccination, and influenza is expected to play a key part in reducing the risk of inadvertent introduction virus. LTRCFs should do all that is practical to encourage high level of vaccine uptake in staff;
2. External contractors should ensure that their staff who may enter clinical areas are also vaccinated and if not vaccinated against COVID-19 that there is a risk-assessment process in place for their staff equivalent to that which the HSE- applies to HSE staff;
3. Staff should participate in testing if this is required by Public Health or IPC.

#### 3.14.2 Movement across facilities

1. See Section 4.1.

#### 3.14.3 Staff occupational health & workforce planning

1. Staff working in a facility that is experiencing an outbreak of COVID-19 or influenza should not work in any other facility;
2. Staff should be informed that they must not attend work if they have fever, cough, shortness of breath, or any kind of new respiratory symptoms. **This continues to apply to staff after vaccination.** They should be aware of their local policy for reporting illness to their manager. Also, if one of their household contacts have respiratory symptoms the staff member should contact their manager for advice before attending work;
3. The COVID-19 NPHEP requires that all staff have their temperature measured twice a day, once being at the start of each shift;
4. In addition, at the start of each shift, all staff should confirm with their line manager that they do not have any symptoms of respiratory illness, such as fever, cough, shortness-of-breath or myalgia. **This continues to apply to staff after vaccination.** Where relevant staff should be asked to confirm that they are not currently working in a facility where there is an outbreak;
5. Staff members who become unwell at work should immediately report to their line manager and should be sent home and advised to contact their GP by telephone. If they

cannot go home immediately, they should be isolated in a separate room until they can go home;

6. Occupational health guidance for healthcare workers is available at:

<https://www.hse.ie/eng/staff/workplace-health-and-wellbeing-unit/covid-19-guidance/> :

- a. Staff members who test positive for COVID-19 may return to work after the end of the isolation period. For staff who have not required hospitalisation for treatment of COVID-19, this will be 10 days after symptom onset (or date when test was taken if no symptoms) provided they have had no fever during the last five days and are medically well. For staff who have required hospitalisation for treatment of COVID-19, this will be after 14 days from symptoms onset (or date when test was taken if symptom onset less clear) provided they have had no fever during the last five days and are medically well. Repeat testing at the end of the illness is generally not appropriate;
- b. Staff members who test positive for influenza may return to work 48 hours after acute symptoms have resolved;
- c. Staff members who are fully vaccinated against COVID-19 and who are household contacts of a case should restrict movement including remaining off work for 5 days from the day on which they are informed that they are a contact. The period of restricted movement ends after 5 full days provided they remain asymptomatic and virus has not been detected on any tests performed during the 5 days.
- d. Fully vaccinated household contacts will receive antigen tests and instructions for use of those tests. If the period of restricted movement (5 days) has elapsed before the person has not completed the recommended schedule of self-testing they should continue to follow the recommended schedule of testing after they end the period of restricted movement.
- e. Staff members who are not fully vaccinated and who have not had a confirmed diagnosis of COVID-19 in the previous 9 months and who have been identified

as contacts of a case of COVID-19 either in the household, community or the occupational setting should restrict their movements and remain off work for 14 days. Exit from restricted movement can happen before 14 days is complete subject to receipt of a PCR test result taken day 10 or later that does not detect virus.

- f. Note that, derogation from exclusion from work may apply to fully vaccinated contacts, including fully vaccinated household contacts, at the discretion of a senior manager and in specific circumstances. . Only senior management can make the decision to derogate a HCW who is a close contact from exclusion from work and this must be based on a risk assessment. Derogation must only be used in **exceptional circumstances** given the inherent risks;
  - g. Any consideration regarding derogations to return to work should be in the context of current occupational health and public health guidance, available on the HSE website.
7. Available Occupational Health supports are detailed in [Appendix B](#).

#### 3.14.4 Visitors, Pastoral Support and other Essential/Important Service Providers

These issues are addressed in a specific document on Guidance on visitation in long-term residential care facilities available at the following link:

<https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/residentialcarefacilities/Normalising%20visiting%20in%20LTRCF.pdf>

COVID-19 Guidance on visits to and from residential facilities for people with disabilities is available on the following link, it also includes detail on how to carry out a risk assessment in relation to visiting:

<https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/disabilityservicesguidance/Normalising%20visiting%20in%20disability%20services.pdf>



An information leaflet for residents and their visitors is available at the following link:

<https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/residentialcarefacilities/Normalising%20visiting%20in%20nursing%20homes%20%20and%20residential%20care%20facilities.pdf>

### 3.14.5 Resident transfers

1. Guidance on resident transfers related to COVID-19 is addressed as an appendix to this document. (**Appendix H**);
2. A video link for self-isolation for people in Residential Care Facilities who have been discharged from hospital is available at: <https://youtu.be/wug2188UNC4> ;
3. For guidance on admission to facilities such as community hospitals and acute rehabilitation units please refer to the document - Acute Hospital Infection Prevention and Control Precautions for Possible or Confirmed COVID-19 in a Pandemic Setting located [here](#);
4. Guidance on admission to RCF applies to residents who routinely use ventilatory support such as CPAP or BiPAP. Use of CPAP or BiPAP is considered an aerosol generating procedure by many (see the following link <https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/aerosolgeneratingprocedures/> );
5. Recent experience in the acute hospital setting suggests there is a high risk of spread if a person on respiratory support with AGP develops infection therefore particular attention to infection prevention and control precautions and a high level of awareness for features of COVID-19 infection is required particularly in residents who have not vaccine protection or have not had COVID-19 in the previous 9 months;
6. Such residents should be admitted to single rooms with a window that can be opened to improve ventilation (subject to weather and security; the goal is gentle air circulation rather than strong air movements) and the door should remain closed as much as possible when ventilatory support is in use;
7. All residents on CPAP or BiPAP, including those fully vaccinated, should have a test for SARS-CoV-2 in the 3 days before admission. They should remain in their room with the door closed all of the time until the result of an additional test taken after admission is available.;
8. In the case that the test result is not available OR if they have any new clinical features to suggest viral infection any care delivered by staff during the use of CPAP or BiPAP, care

- provided by staff should be delivered with airborne precautions (minimise numbers and time in the room, maximise ventilation as far as is practical and use of appropriate PPE);
9. If the test results for SARS-CoV-2 are reported as not detected, the person is fully vaccinated and not a COVID-19 close contact and there are no clinical features to suggest viral infection, care provided by staff should be delivered with Standard Precautions plus use of a surgical mask;
  10. All newly admitted residents who are not fully vaccinated and have not had COVID-19 in the previous 9 months, should not participate in social activities within the LTRCF and should stay in their room most of the time for the initial 14 days. A link to a video for residents of LTRCF on self-isolation following discharge from hospital is available [here](#);
  11. If a person who is not fully vaccinated or in the 9 months after diagnosis of COVID-19, is on CPAP or BiPAP, is not a contact of COVID19, and if the samples taken before and after admission are reported as COVID-19 not detected, they can move around outside their room and participate in activities subject to confirming each day that there is no deterioration in their condition that could suggest COVID-19 or other viral respiratory tract infection;
  12. If at any point during admission to the RCF a person who uses CPAP or BiPAP develops symptoms consistent with COVID19 or influenza, appropriate transmission-based precautions should be introduced immediately while arrangements are made for assessment by their doctor. Staff should follow the guidance on use of PPE for AGPs while managing a suspected or confirmed case of COVID19, found here: <https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/aerosolgeneratingprocedures/>

### **3.15 Increased surveillance and early identification of cases of COVID-19 infection**

1. Surveillance (monitoring for illness) is an essential component of any effective infection prevention and control programme;
2. RCFs should ensure that they have means in place to identify a new case of COVID-19 and control transmission, through active monitoring of residents and staff for new symptoms of infection, rapid application of transmission-based precautions to those with suspected

COVID-19, prompt testing of symptomatic residents and referral of symptomatic staff for evaluation. Current case definition can be found [here](#);

3. The RCF should ensure that there is twice daily active monitoring of residents for signs and symptoms of respiratory illness or changes in their baseline condition (e.g., increased confusion, falls, and loss of appetite or sudden deterioration in chronic respiratory disease);
4. There should be early identification of staff absence/s, which may be due to COVID-19 or influenza infection.

#### **4 Management of an outbreak of COVID-19 or Influenza**

When there is a suspicion of cases of COVID-19, influenza or other acute respiratory virus infection the MOH should perform a risk assessment to determine whether there is either possible or confirmed active transmission in the facility. An isolated positive result of SARS-CoV-2 or influenza in a resident or staff member is not in itself proof of current active transmission.

When an outbreak is suspected laboratory testing should be arranged as quickly as possible. However, it is not appropriate to wait for laboratory test results before beginning initial investigation, contacting Public Health or implementing control measures. There should be heightened awareness among staff, so that other residents with symptoms are quickly identified.

A local incident management meeting should be arranged promptly and involve key staff members including housekeeping, nursing staff, allied healthcare professional and medical staff.

This group should:

1. Try and establish whether it is likely that an outbreak is occurring, taking in to account the following:
  - a. Could onward transmission have already occurred? (e.g., resident had widespread contact with others in the 48 hours before symptom onset):
    - i. Are they in a single room or sharing?

- ii. Is the resident ambulatory?
  - iii. Have they spent time with others in communal areas or group activities?
  - iv. Are there behavioural characteristics, which might be increased risk of transmission?
  - v. Are all or most residents vaccinated against COVID-19 and influenza?
2. Identify if any other residents are symptomatic and if so, what are their symptoms?
  3. Identify are any staff symptomatic or has there been an increase in staff absence?
  4. Identify residents and staff who were in close contact with the symptomatic resident/s in the 48 hours before symptom onset or before isolation and transmission-based precautions were implemented.

The initial management of the possible case and, where relevant, contacts should be the same as for a confirmed case until an alternative diagnosis has been identified.

The contact tracing guidance varies when VOCs are suspected/ confirmed. Please see <https://www.hpsc.ie/a-z/respiratory/coronavirus/novelcoronavirus/sars-cov-2variantsofconcern/> for further details

#### **4.1 Declaring an outbreak**

For surveillance purposes, the following outbreak definition applies:

##### **Confirmed**

A cluster/outbreak, with two or more cases of laboratory confirmed infection with the same pathogen (COVID-19, Influenza or other respiratory virus).

##### **OR**

A cluster/outbreak, with two or more cases of illness with symptoms consistent with the same pattern of infection related illness, and at least one person is laboratory confirmed.

## **Suspected**

A cluster/outbreak, with two or more cases of illness with symptoms consistent with the same pattern of infection related illness.

### **4.2 Outbreak Control Team (OCT)**

All outbreaks of infectious disease including COVID-19 or influenza in a RCF must be reported to the regional Medical Officer of Health (MOH) at the Department of Public Health at the earliest opportunity.

Public Health doctors from the Regional Department of Public Health will provide overall leadership for the management of the outbreak in the RCF.

Ideally, the OCT should have regular, active involvement of a Public Health Doctor. However, if that is not practically possible, following initial consultation and advice from Public Health, the OCT should liaise on a regular ongoing basis with the regional Public Health Department to provide updates on outbreak progress and seek further advice as appropriate.

The OCT membership should be decided at local level and will depend on available expertise.

An OCT Chairperson should be agreed.

Members of the OCT may include any of the following. However, in many settings it may not be possible to include all the expertise referred to below:

1. Specialist in Public Health Medicine and/or Public Health Department Communicable Disease Control Nurse Specialist;
2. GP/Medical officer/Consultant to RCF (dependent on nature of RCF);
3. Director of Nursing or Nurse Manager from RCF;
4. Management representative from the RCF i.e. manager or CEO;
5. Community Infection Prevention and Control Nurse (CIPCN) where available;
6. Administration support.

Other members who may need to be included, particularly if it is an extensive or prolonged outbreak include:

1. Community Services General Manager;

2. Administrative support;
3. Occupational Medicine Physician;
4. Consultant Clinical Microbiologist;
5. Representative from HPSC;
6. Communications officer.

Every member involved should have a clear understanding of their role and responsibility.

The frequency required for the OCT meeting should be decided and they should be carried out in consideration of physical distancing requirements via teleconference/videoconference facilities. Public Health will formulate a case definition, assign an outbreak code and decide as to whether an on-site visit is required or not.

The RCF should inform HIQA or Mental Health Commission, as appropriate and the local CHO as per usual protocols.

Before the first meeting of the OCT, the local incident team should gather as much information as possible to include:

1. A line list of all residents and staff. Template can be found in Appendix C;
2. The vaccination history (COVID-19 and influenza) of all residents and staff;
3. Identify the total number of people ill (residents & staff), dates of illness onset and the spectrum of symptoms;
4. Identify staff and residents who have recently recovered, developed complications, been transferred to acute hospitals and those who have died;
5. Information on laboratory tests available including the number of tests taken to date and the date sent to the laboratory, along with the tests requested and reported results;
6. Determine if the number of symptomatic residents/staff involves more than one unit/floor/ward or if the outbreak is confined to one area only;
7. Use the case definitions for possible, probable and confirmed COVID-19 available on the HPSC website if relevant [HERE](#).
8. A checklist for outbreak management can be found in Appendix D.

### 4.3 Management of a possible or confirmed case of COVID-19 or Influenza

1. The initial assessment of the resident should be performed by their doctor;
2. If COVID-19 or influenza is suspected, the doctor will arrange testing;
3. If the clinical condition does not require hospitalisation, the resident should not be transferred from the facility on infection prevention and control grounds;
4. Where there is capacity and it is appropriate to their care needs, a resident with possible or confirmed COVID-19 or influenza should be placed in a single room with transmission-based precautions and appropriate use of PPE by staff (**Appendix F & Appendix G**). Staff assigned to care of a resident in these circumstances should preferably be staff who have vaccine protection. This is because vaccination is the most effective way to protect staff although good IPC practice provides substantial protection to staff;
5. Room doors should be kept closed where possible and safe to do so;
6. Practical measures to increase ventilation should be taken consistent with comfort and weather. Note: the intention is to achieve gentle air circulation rather than strong air movements;
7. When this is not possible, ensure the resident's bed is moved to the furthest safe point in the room to try and achieve at least 1 m distance to the door;
8. Display signage to reduce entry into the room, but confidentiality must be maintained;
9. Take time to explain to the resident the importance of the precautions that are being put in place to manage their care and advise them against leaving their room;
10. Ideally, the resident's single room should have ensuite facilities;
11. If ensuite facilities are not available, try to designate a commode or toilet facility for the resident's use;
12. In the event of a commode being used, the HCW should exit the resident's room while wearing appropriate PPE, transport the commode directly to the nearest sluice (dirty utility) and remove the PPE in the sluice after placing the contents directly into the bed pan washer or pulp disposal unit. A second person should be available to assist with opening and closing doors to the single room and sluice room. If a second person is not



available, change gloves and perform hand hygiene and put on a clean pair of disposable gloves;

13. If the resident must use a communal toilet, ensure it is cleaned after every use;
14. Listen and respond to any concerns residents may have to ensure support and optimal adherence is achieved during their care;
15. If well enough, a resident who has infection should be facilitated in going outside alone if appropriate or accompanied by a staff member maintaining adequate distance from both staff and other residents. If the staff member can maintain this distance, they do not need to wear PPE;
16. If the resident passes briefly through a hallway or other unoccupied space to go outside, there is no requirement for any additional cleaning of that area beyond normal good practice;
17. Residents with confirmed COVID-19 or influenza will require appropriate healthcare and social support, including access to their doctor or GP for medical management and on-site support;
18. Residents with influenza or who are exposed to influenza may require treatment with antiviral medication
19. A care planning approach that reflects regular monitoring of residents with COVID-19 or influenza infection for daily observations, clinical symptoms and deterioration should be put in place. Where appropriate there should be advance planning in place with residents and / or advocates reflecting preferences for end of life care and / or transfer to hospital in event of deterioration. Staffing levels / surge capacity planning should reflect the need for an anticipated increase in care needs during an outbreak;
20. **Residents with confirmed COVID-19 infection** should remain in isolation on Contact and Droplet precautions until 14 days after the first date of onset of symptoms and they are fever free for the last five days. A respirator mask (such as an FFP2 mask) and eye protection must be available for use by staff caring for such a resident during the period when they are infectious. In some circumstances a healthcare worker may choose to use a surgical mask and face shield. This also affords substantial protection. If this option is

used it must be based on the healthcare workers risk assessment and not because they lack access to respirator masks. Samples from residents with confirmed COVID-19 after completion of vaccination should be submitted for further testing to identify the SARS-CoV-2 variant if the sample is suitable. The laboratory that performed the test will forward the sample if the sample has been stored;

21. Antiviral treatment for patients with influenza should be discussed promptly with the resident's doctor or with Public Health. Treatment is likely to be more effective if started early;
22. **Residents with confirmed influenza virus infection** should remain in isolation for 7 days from the onset of symptoms and until 48 hours after resolution of fever;
23. Healthcare workers with confirmed influenza should remain off work until 5 days after onset of respiratory symptoms and until 48 hours after resolution of fever;
24. Staff should be mindful that prolonged isolation is stressful for most residents and to encourage relatives and other residents where practical to communicate with them regularly via phone or video calls and where possible window visits;
25. If residents need to see someone, for example because they are distressed or approaching end of life, this should be facilitated if the person they wish to see understands that there is a risk of infection and can follow recommendations to lower the risk of infection.

#### **4.4 Cohorting residents with possible or confirmed COVID-19 or Influenza**

1. Placement of residents with possible or confirmed COVID-19 or influenza in a designated zone, with designated staffing to facilitate care and minimise further spread is known as cohorting;
2. As the lay-out for each RCF will differ, cohorting requires planning for each facility, and the zoned area might be a floor, a wing or a separate annex. In these zoned areas, heightened infection prevention and control measures are critical and practical measures to improve ventilation should be implemented;
3. It is often not practical in a LTRCF to ensure that all of the staff who work in the cohort area are assigned exclusively to work in that area. In such circumstances every practical

effort should be made to ensure that the number of individual staff members and the frequency with which individual staff members are required to move between the cohort area and other areas during the same shift is minimised. Staff members who are required to move between cohort and other areas should be fully vaccinated.

4. In the rare event of outbreak of two viral infections outbreaks do not cohort people with COVID-19 and those with influenza together
5. Cohorting includes residents who are placed in single rooms close together, or in multi-occupancy areas within the building or section of a ward/unit;
6. Where possible, residents with probable or confirmed COVID-19 or influenza should be isolated in single rooms with ensuite facilities. If there are multiple residents and if it is practical to do so, these single rooms should be located in close proximity to one another in one zone, for example on a particular floor or area within the facility;
7. Where single room capacity is exceeded and it is necessary to cohort residents in a multi-occupancy room:
  - a. residents with **a confirmed diagnosis of COVID-19** can be cohorted together;
  - b. Residents with **a confirmed diagnosis of influenza** can be cohorted together
  - c. Residents with suspected COVID-19 or influenza should not be cohorted with those who are confirmed positive;
  - d. The risk of cohorting **suspected cases** in multi-occupancy areas is much greater than that of cohorting confirmed positive residents together, as the suspected cohort is likely to include residents with and without the specific virus infection;
8. Where residents are cohorted in multi-occupancy rooms, every effort should be made to minimise cross-transmission risk:
  - a. Maintain as much physical distance as practical between beds (minimum of 1m); if possible reduce the number of residents/beds in the area to facilitate physical distancing;
  - b. Close privacy curtains if available between the beds to minimise opportunities for close contact.

9. There should be clear signage indicating that the area is a designated zone to alert staff about cohorting location in the RCF. A zone may have multi-occupancy rooms or a series of single rooms;
10. A designated cohort area should ideally be separated from non-cohort areas by closed doors;
11. Minimise unnecessary movement of staff in cohort areas and ensure that the number of staff entering the cohort area is kept to a minimum;
12. Staff working in cohort areas should be fully vaccinated and should not be assigned also to work in non-outbreak areas, where possible;
13. In so far as is possible, the cohort area should not be used as a thoroughfare by other residents, visitors or staff, including residents being transferred, staff going for meal breaks and staff entering and exiting the building;
14. A respirator mask (such as an FFP2 mask) must be available for use by staff working in a cohort area for COVID-19. A surgical mask and face shield also affords substantial protection against transmission of COVID-19 if the healthcare worker prefers this option after risk assessment. A surgical mask is generally appropriate when caring for person with influenza virus infection;
15. Improve ventilation in the cohort area in so far as practical consistent with comfort and safety; the goal is gentle air circulation rather than strong air movements.

#### **4.5 Management of close contacts of a possible or confirmed case of COVID-19**

1. Residents who are close contacts of a confirmed case should be accommodated in a single room with their own bathing and toilet facilities, unless they are fully vaccinated or have had COVID-19 in the previous 9 months. If this is not possible, cohorting in small groups (two to four) with other close contacts is acceptable;
2. Although the risk for close contacts who are fully vaccinated or have had COVID-19 in the previous 9 months is much lower than for residents who do not have vaccine protection, a high degree of caution as in bullet point 1 above remains appropriate in the initial stages of dealing with an outbreak;

3. Residents who are close contacts should be advised to avoid communal areas and stay in their room where it is practical to do so until 10 days after exposure if they have vaccine protection or have had COVID-19 in the previous 9 months. Where a person is not fully vaccinated the longer period of 14 days should apply; this may also apply on the advice of a public health doctor or infection prevention and control doctor;
4. Residents who are close contacts may go outside if appropriate, alone or accompanied by a staff member maintaining adequate distance. An accompanying staff member in this situation is not required to wear PPE if distance can be maintained;
5. Note: testing of residents who are close contacts for COVID-19 should be performed in accordance with current public health guidance;
6. Regardless of the outcome of any tests already performed, the resident should be referred to their doctor for assessment at any time if they develop symptoms of infection;
7. It is understood that some residents may, due to underlying conditions (e.g. dementia with wandering behaviours) have significant difficulties with isolation and / or restricted movement. In these instances, the creation of a 'safe zone' may be the most appropriate support to prevent distress arising from confinement. Separate access to outdoor spaces or communal rooms not used by other residents may be appropriately used when followed by environmental cleaning and disinfection if required. The associated risks are lower if the residents concerned are fully vaccinated;
8. If entry to an occupied shared space is unavoidable, the resident should be encouraged to perform hand hygiene and wear a surgical mask or to cover their mouth and nose with a tissue;
9. A respirator mask (such as an FFP2 mask) must be available for use by staff working in the cohort area for COVID-19 during the period when the residents are infectious. Use of a surgical mask and face shield also affords substantial protection if the healthcare worker prefers this option after risk assessment.

#### **4.6 Management of close contacts of a possible or confirmed case of Influenza**

1. Residents who are close contacts of a confirmed case of influenza should be accommodated in a single room with their own bathing and toilet facilities. If this is not possible, cohorting in small groups (two to four) with other close contacts is acceptable;
2. Antiviral prophylaxis may be appropriate for contacts as per advice from their Public Health;
3. Testing of asymptomatic close contacts of influenza cases is not required.

## **4.7 Infection prevention and control measures**

### **4.7.1 Standard precautions**

**Note that recommendations with respect to Infection Prevention and Control Practice are the same for those who have had COVID-19 and influenza vaccine and for those who have not.**

Standard Precautions are the minimum infection prevention practices that apply to the care of all people, regardless of suspected or confirmed infection status of the person, in any setting where health care is delivered. For further information on Standard Precautions and the chain of infection refer to HSEland online learning or [www.hpsc.ie](http://www.hpsc.ie). With regard to COVID-19 and influenza, key elements include:

### **4.7.2 Hand hygiene**

1. Hand hygiene is the single most important action to reduce the spread of infection in health and other social care settings and is a critical element of standard precautions;
2. Facilities must provide ready access for staff, residents and visitors to hand hygiene facilities and alcohol-based hand rub (ABHR);
3. Staff should adhere to the WHO five moments for hand hygiene:
  - a. Hand hygiene must be performed immediately before every episode of direct resident care and after any activity or contact that potentially results in hands becoming contaminated, including the removal of PPE, equipment decontamination, handling of waste and laundry.

4. Residents should be encouraged and facilitated to clean their hands after toileting, after blowing their nose, before and after eating and when leaving their room. If the resident's cognitive state is impaired, staff must help with this activity;
5. **Gloves should not be used in routine care of residents to whom Standard Precautions apply unless contact with blood or body fluids (other than sweat), non- intact skin or mucous membranes is anticipated. When gloves are required, they are not a substitute for hand hygiene. Hand hygiene is required before putting on gloves and immediately after they have been removed;**
6. HSEland hand hygiene training is available online and staff should be encouraged to do refresher training at [www.hseland.ie](http://www.hseland.ie).

Refer to hand hygiene information posters Appendix E.

#### **4.7.3 Respiratory hygiene and cough etiquette**

1. Respiratory hygiene and cough etiquette refer to measures taken to reduce the spread of viruses via liquid respiratory particles produced when a person coughs or sneezes;
2. Disposable single-use tissues should be used to cover the nose and mouth when sneezing, coughing or wiping and blowing the nose;
3. Used tissue should be disposed of promptly in the nearest foot operated waste bin;
4. Some residents may need assistance with containment of respiratory secretions. Those who are immobile will need a waste bag at hand for immediate disposal of the tissue. Hands should be cleaned with either soap and water or ABHR after coughing sneezing, using tissues or after contact with respiratory secretions and contaminated objects;
5. Staff and residents should be advised to try to avoid touching their eyes, mouth and nose.

#### 4.7.4 Personal Protective Equipment (PPE)

As part of Standard Precautions, it is the responsibility of every HCW to undertake a risk assessment PRIOR to performing a clinical care task, as this will inform the level of IPC precautions needed, including the choice of appropriate PPE for those who need to be present.

Full guidelines on the appropriate selection and use of PPE Appendix F and G and <https://www.hpsc.ie/az/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/ppe/>. Current guidance for the use of surgical masks by HCW in the context of COVID-19 states that:

1. HCWs should wear surgical masks when providing care to residents within 1 to 2 m of a resident. If a resident has suspected or confirmed COVID-19 or is a COVID-19 contact staff should have access to a respirator mask (for example FFP2) and other appropriate PPE;
2. HCWs should wear surgical masks for all encounters with other HCWs in the workplace where a distance of 2m cannot be maintained and the encounter is expected to last longer than 15 minutes;
3. HCWs are also required to wear a surgical mask in busy public areas of healthcare facilities;
4. In the context of a ward or unit-based outbreak it is appropriate to consider all patients in that setting as suspected or confirmed COVID-19 or influenza cases while active transmission is ongoing;
5. Decisions regarding when all patients should be considered as suspected or confirmed COVID-19 cases requiring use of general use of FFP2 masks and other appropriate PPE should be made by the IPC team and outbreak control team;
6. In some circumstances a healthcare worker may choose to use a surgical mask and face shield. This also affords substantial protection. If this option is used it must be based on the healthcare workers risk assessment and not because they lack access to respirator masks;
7. Facemasks that are worn by HCW's in circumstances other than contact with residents with suspected or confirmed COVID-19 or COVID-19 contacts may be disposed of in the domestic waste stream;



8. Educational videos are also available on [www.hpsc.ie](http://www.hpsc.ie) at
9. <https://www.hpsc.ie/az/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/videoresourcesforipc/>;
10. All staff must be trained in the proper use of all PPE that they may be required to wear;
11. The Health and Safety Authority indicate that where a risk assessment indicates that workers need to use a close-fitting respirator mask for their protection that every effort should be made to comply with the requirement for fit testing of the workers, as far as is reasonably practicable. When fit testing of all staff is not immediately possible, then fit testing should be prioritised for those at greatest risk;
12. Note that in outbreak situations or other exceptional circumstances where extended use of some items of PPE (other than gloves) when moving between people care for with a confirmed diagnosis of COVID-19 might be considered, it is important to make every effort to avoid generalised use of PPE throughout the facility without considering the level of risk. Note that extended use of PPE for the sole purpose of reducing PPE use is not appropriate, as PPE supplies should be sufficient to meet requirements;
13. In the event of extended use of PPE being necessary, define clean and contaminated zones. PPE should be donned before entering the contaminated zone and doffed and hand hygiene performed before entering clean zones. Where staff members are having meals on a unit to minimise staff interaction, it is essential that the staff refreshment area is a clean zone. Corridors between units should be designated clean zones. Clinical stations should normally be clean zones;
14. Transiting through the hallway of a contaminated zone without providing resident care does not require use of PPE, if the residents are in their rooms and there is no physical contact with other staff wearing PPE.

#### **4.7.5 Transmission-based Precautions for COVID-19 and Influenza virus**

1. Transmission-based Precautions are IPC measures which are implemented in addition to Standard Precautions when Standard Precautions alone are insufficient to prevent the onward transmission of specific infectious diseases. See Appendix G. They include

contact, droplet and airborne precautions. In general, COVID-19 and influenza are spread mainly through the air over short distances in respiratory liquid particles. Transmission may also occur through contact with the surfaces contaminated with respiratory secretions of someone with infection.

2. Transmission of virus through the air over long distance can occur in some circumstances for example during aerosol generating procedures (AGP) or in crowded and poorly ventilated settings;
3. <https://www.hpsc.ie/az/respiratory/coronavirus/novelcoronavirus/guidance/infectionpreventionandcontrolguidance/aerosolgeneratingprocedures/> ;
4. Transmission-based Precautions should be applied immediately to all suspected cases of COVID-19 and influenza.

#### 4.8 Duration of transmission based precautions

1. **For COVID-19** transmission based precautions can be **discontinued fourteen days after symptom onset, where a person has been fever free for five days, with the exception of people on respiratory support devices that are aerosol generating. For residents who are on respiratory support devices that are aerosol generating the period should be extended to 21 days;**
2. A test of clearance is generally not appropriate for residents who have been diagnosed with COVID-19;
3. Note: some people who meet the above criteria (14 days post onset with 5 days fever free) may have a persistent cough. There is no evidence that such people pose a specific infection risk or that transmission-based precautions should be continued. An extended period of contact and droplet precautions may be considered in some such cases if there is clinical concern. In such cases the period of Transmission based precautions of 21 days may be reasonable. In any case Transmission based precautions should not be extended beyond 28 days;
4. **For Influenza** transmission based precautions can be discontinued after 7 days provided fever has resolved for 2 days or more at that time.

#### **4.8.1 Care Equipment**

1. Where possible, use single-use equipment for the resident and dispose of it as healthcare risk waste into a designated healthcare risk waste bin inside the room;
2. Where single use equipment is not possible, use designated care equipment in the resident's room or cohort area. In a cohort area, the equipment must be decontaminated immediately after use and before use on any other resident following routine cleaning protocols;
3. If it is not possible to designate pieces of equipment to the resident or cohort area these must be decontaminated immediately after use and before use on any resident following standard cleaning protocols;
4. There is no need to use disposable plates or cutlery. Crockery and cutlery should be washed after use in a dishwasher or by handwashing, using household detergent and hand-hot water.

#### **4.8.2 Management of blood and body fluid spillages**

1. Should be managed in line with local policy.

#### **4.8.3 Management of waste**

1. Dispose of all waste from residents with confirmed or suspected COVID-19 or influenza virus as healthcare risk waste during the period when transmission based precautions applies (also referred to as clinical risk waste);
2. When removing waste, it should be handled as per usual precautions for healthcare risk waste;
3. The external surfaces of the bags/containers do not need to be disinfected;
4. All those handling waste should wear appropriate PPE and clean their hands after removing PPE;
5. Hands-free healthcare risk waste bins should be provided in single rooms and cohort areas;
6. If a healthcare risk waste service is not available in the RCF, then all consumable waste items that have been in contact with the individual, including used tissues, should be put

in a plastic rubbish bag, tie the bag, place in a second bag and leave for 72 hours. This should be put in a secure location prior to usual waste collection;

7. Bodily waste, such as urine or faeces from individuals with possible or confirmed COVID-19 does not require special treatment and can be discharged into the sewage system.

#### **4.8.4 Safe management of linen (laundry)**

1. All towels, clothing or other laundry used in the direct care of residents with suspected and confirmed COVID-19 should be managed as 'infectious' linen;
2. Linen must be handled, transported and processed in a manner that prevents exposure to the skin and mucous membranes of staff, contamination of their clothing and the environment;
3. Disposable gloves and an apron should be worn when handling linen;
4. All linen should be handled inside the resident room/cohort area. A laundry skip/trolley should be available as close as possible to the point-of-use for linen deposit, for example immediately outside the cohort area/isolation room;
5. When handling linen, the HCW should not:
  - a. rinse, shake or sort linen on removal from beds/trolleys;
  - b. place used/infectious linen on the floor or any other surfaces (e.g., a bedside locker/table top);
  - c. handle used/infectious linen once bagged;
  - d. overfill laundry receptacles; or
  - e. Place inappropriate items in the laundry receptacle (e.g., used equipment/needles).
6. When managing infectious linen, the HCW should:
  - a. Place linen directly into a water-soluble/alginate bag and secure;
  - b. Place the alginate/water-soluble bag into the appropriately coloured linen bag (as per local policy);
  - c. Store all used/infectious linen in a designated, safe area pending collection by a laundry service;

- d. If there is no laundry service, laundry should be washed using the hottest temperature that the fabric can withstand and standard laundry detergent;
- e. Laundry may be dried in a dryer on a hot setting.

#### **4.8.5 Environmental hygiene**

1. The care environment should be kept clean and clutter free in so far as is possible, bearing in mind this is the resident's home and they are likely to want to personalise their space with objects of significance to them;
2. Ventilation should be maintained in so far as practical taking account of comfort and weather. Note that the goal is to achieve reasonable air exchange with gentle air movement. Strong airflow into the room from outside that is readily felt and causes discomfort is not required and may contribute to airflow out of the room.
3. Residents' observation charts, medication prescription and administration records (drug kardexes) and healthcare records should not be taken into the resident's room, to limit the risk of contamination.

#### **4.8.6 Routine cleaning in the context of COVID-19 and Influenza**

1. Decontamination of equipment and the care environment must be performed using either:
  - a. A combined detergent/disinfectant solution at a dilution of 1,000 parts per million available chlorine (ppm available chlorine (av.cl.)); or;
  - b. A general-purpose neutral detergent in a solution of warm water, followed by a disinfectant solution of 1,000 ppm av.cl;
  - c. Only cleaning (detergent) and disinfectant products supplied by employers are to be used. Products must be prepared and used according to the manufacturer's instructions and recommended product "contact times" must be followed.

2. Vacuuming of carpet floor in a resident's room should be avoided during an outbreak and while the person is infectious. When the resident is recovered the carpet should be steam cleaned;
3. All shared spaces should be cleaned with detergent and disinfectant;
4. Equipment used in the cleaning/disinfection of the isolation area should be single use where possible and stored separately to equipment used in other areas of the facility;
5. Household and care staff should be trained in the appropriate use and removal of PPE (**Appendix F**);
6. In practical terms, single room cleaning may be undertaken by staff who are also providing care to the resident while in the single room.

#### **4.8.7 Frequency of cleaning in the context of COVID-19 and Influenza**

1. All surfaces in the resident room/zone of people who have infectious COVID-19 or influenza should be cleaned and disinfected at least daily and when visibly contaminated. These include high-touch items; bedrails, bedside tables, light switches, remote controls, commodes, doorknobs, sinks, surfaces and equipment close to the resident (e.g., walking frames, sticks, phone or other mobile device);
2. Handrails and table tops in facility communal areas, along with nurses' station counter tops and equipment require regular cleaning;
3. Cohort areas and clinical rooms must be cleaned and disinfected at least daily and when visibly contaminated and a documented cleaning schedule should be available to confirm this.

#### **4.8.8 Terminal cleaning**

1. Terminal cleaning should always be performed after a resident who has had infectious COVID-19 has vacated the room and is not expected to return. In addition to the routine cleaning protocols, a terminal clean is needed;
  - a. Removal of all detachable objects from a room or cohort area, including laundry and curtains;

- b. Removal of waste;
  - c. Cleaning (wiping) of lighting and ventilation components on the ceiling;
  - d. Cleaning of the upper surfaces of hard-to-reach fixtures and fittings;
  - e. Cleaning of all other sites and surfaces working from those at higher level down to floor level.
2. A terminal clean checklist is good practice to support cleaning or household staff to effectively complete all environmental cleaning tasks, which should be signed off by the cleaning supervisor before the room reopens for occupancy;
3. The use of novel technologies for room disinfection have not been shown to add value beyond standard cleaning and disinfection and are not recommended. If they are used, they must be used in addition to and not as a substitute for cleaning.

#### **4.8.9 Staff uniforms/clothing**

1. Staff uniforms are not considered to be personal protective equipment;
2. The appropriate use of PPE will protect staff uniforms from contamination in most circumstances;
3. Uniforms should be laundered:
  - a. separately from other household linen;
  - b. in a load not more than half the machine capacity;
  - c. at the maximum temperature the fabric can tolerate.
4. The risk of virus transmission from contaminated footwear is likely to be extremely low. Shoe covers should not be used. However, HCW could consider designating a pair of comfortable, closed, cleanable shoes for wearing in a COVID-19 care area;
5. Staff should avoid bringing personal items, including mobile phones into isolation or cohort areas.

#### **4.9 Communication**

1. Good communication is essential for residents, family and staff members;

2. Provide regular information sessions and education on measures required for staff members and assign someone to do these.

#### **4.10 Support services for staff and residents**

1. The effect on staff and residents during outbreak events should not be underestimated especially where there have been deaths in the RCF. Every effort should be made to support those who are impacted by outbreak events;
2. One of the key supports to staff is to promote vaccination. It is important that staff with questions about the benefits and risks of vaccination have access to appropriate support.

### **5 Care of the person with suspected or confirmed COVID-19 or a Contact of COVID-19 who is dying**

1. A compassionate, pragmatic and proportionate approach is required in the care of those who are dying;
2. The presence of a person close to the resident should be facilitated. They should be aware of the potential infection risk;
3. If the person who wants to be with the resident is fully vaccinated or has had COVID-19 in the previous 9 months, the risk is much reduced;
4. Pastoral care team where requested by the person or their family and who are willing to attend should NOT be restricted from entering the facility;
5. All persons in attendance should be advised to wear a mask and plastic apron. A respirator mask should be available to those who wish to use them although it is unlikely that it will be practical to train most people in their correct use in this context. A surgical mask also affords substantial protection. Gloves are not essential, so long as those in attendance understand the risks; perform hand hygiene after touching the person and before leaving the room. Visitors should be instructed on how to put on and take off the PPE and how to perform hand hygiene. Where practical, visitors should be supervised when donning and doffing PPE;



6. The use of PPE is less critical if the person accompanying the dying person is fully vaccinated or has had COVID-19 in the previous 9 months, but people should have access to PPE;
7. For the anointing of the sick or other rites where only transient physical contact is required, gloves are not necessary, so long as hand hygiene is performed immediately after anointing or touching the person;
8. Visitors should avoid interacting with residents other than the person they are accompanying.

## **6 Care of the recently deceased**

### **6.1 Hygienic preparation**

1. Any IPC precautions that have been advised before death must be continued in handling the deceased person after death. In relation to COVID-19 specifically if transmission based precautions have been discontinued before death, then they are not required after death – see section on duration of transmission based precautions;
2. Hygienic preparation includes; washing of the face and hands, closing the mouth and eyes, tidying the hair and in some cases, shaving the face;
3. Washing or preparing the body for religious reasons is acceptable if those carrying out the task wear long-sleeved gowns, gloves, a surgical face mask and eye protection, if there is a risk of splashing.

### **6.2 Handling personal possessions of the deceased**

1. Most jewellery including watches, rings, bracelets, earrings and items like photo frames can be wiped down using a detergent/disinfectant wipe. Alternatively, items of jewellery (with the exception of watches) can be placed in hot, soapy water and cleaned first, then rinsed and dried using disposable paper towels;
2. Items of clothing and soft toys should be placed directly into a washing machine and washed;

3. Paper materials (e.g. books, prayer books/bible) or items that cannot be wiped should be placed in a bag and left aside for 72 hours before handling;
4. Clothing that needs to be hand washed should be placed in water and detergent at a suitable temperature with minimal handling. Handling before placing in water and washing should be performed with gloved hands;
5. Personal belongings that family members wish to discard should be placed in a plastic bag and tied securely and can go out for collection in the appropriate general waste stream.

### **6.3 Transport to the mortuary**

1. An inner lining is not required in terms of COVID-19 risk or influenza risk, but may be required for other practical reasons such as maintaining dignity or preventing leakage affecting the mortuary environment;
2. A surgical face mask or similar should be placed over the mouth of the deceased before lifting the remains into the inner lining;
3. Those physically handling the body and placing the body into the coffin or the inner lining should wear, at a minimum, the following PPE:
  - a. Gloves;
  - b. Long sleeved gown;
  - c. Surgical face mask.
4. Pay close attention to hand hygiene after removal of PPE;
5. The family should be advised not to kiss the deceased and should clean their hands with alcohol hand rub or soap and water after touching the deceased.

**PPE is not required for transfer, once the body has been placed in the coffin.**

## **7 Monitoring outbreak progress**

1. Monitoring the outbreak will include ongoing surveillance for symptoms in residents and staff to identify new cases and to update the status of ill residents and staff;
2. The nominated RCF liaison person should update the line listing with new cases or developments as they occur and communicate this to the OCT on a daily basis or more

frequently if major changes occur, in line with Public Health recommendations until the outbreak is declared over;

3. The review of this information should examine issues of ongoing transmission and the effectiveness of control measures;
4. Institute active daily surveillance for fever, respiratory symptoms, including cough and other symptoms suggestive of COVID-19 or influenza, in residents and staff until the outbreak is declared over.

## **8 Declaring the outbreak over**

In order to formally declare that the outbreak is over, the RCF should not have experienced any new cases of infection (resident or staff) considered as likely to have been acquired in the RCF which meet the case definition for a period of two incubation periods.

In most practical terms the RCF can resume normal function after 1 incubation period has elapsed without evidence of continuing transmission but there should be a high level of vigilance regarding symptoms of infection until the outbreak is formally closed.

As above, an isolated positive result of SARS-CoV-2 in a resident or staff member is not of itself evidence of ongoing transmission.

## Appendix A: Prevention and control of outbreaks of COVID-19 in RCF

	Domain	Action	Comment
<b>Pre-Outbreak Measures</b>	Planning and Administration	Written Policies	Immunisation policies Standard and Transmission based Precautions including droplet and contact Written outbreak management plan
		RCF Lead (Named person)	To oversee development, implementation and review of policies and procedures
		Training and Education	For all staff Ongoing training – Standard and Transmission-based Precautions, PPE Measures to improve compliance
		Provision of supplies	Hand hygiene supplies, PPE, cleaning and disinfection materials, viral swabs, request forms and arrangements for prioritised testing of samples
	Vaccination	Regularly review the uptake of vaccination in residents and staff.	
	Standard Precautions	Standard infection control procedures	Standard Precautions and mask use should be practised by all staff at all times
	Surveillance	Awareness of signs and symptoms of COVID	Formal process to record any new symptomatic residents twice daily
<b>Early recognition</b>	Case Definition	As per HPSC guidance	Case definition may change as pandemic progresses
	Outbreak Definition	Action threshold for outbreak control measures	One suspected or confirmed case for public health action
	Communication of suspected outbreak	Notification of senior management, medical and public health staff, CHO and NH lead	Follow RCF algorithm
	Formation of outbreak control team (OCT)	OCT may be convened following risk assessment	
	Testing	Viral swab	As per current guidance
	Initial Actions	Daily Case list	
		Activate Daily surveillance	
		Appropriate IPC precautions in place	Contact and Droplet precautions in the cohorted area/zone. Note must have access to respirator masks (such as FFP2)

	Domain	Action	Comment
During an Outbreak		Resident placement	Single rooms Cohorting or zone allocation
		Respiratory etiquette	
	Infection Control Measures	Hand Hygiene	5 Critical points: <ul style="list-style-type: none"> <li>• Before patient contact</li> <li>• Before an aseptic procedure</li> <li>• After body fluid exposure</li> <li>• After patient contact</li> <li>• After contact with patient surroundings</li> </ul> Hand hygiene after PPE removal
		PPE	Gloves Mask (respirator mask or surgical mask) Aprons / Gowns Face protection (as required based on risk assessment)
		Aerosol Generating Procedure associated with increased risk of infection (AGP)	See HPSC guidance <a href="#">document</a> . Ventilation, closed door, respirator mask (FFP2), gown, eye protection and gloves
	Environmental control measures		Resident environmental cleaning and disinfection Residential care equipment Laundry Eating utensils and crockery Practical measures to increase ventilation to the greatest extent practical consistent with comfort and weather (gentle movement of air rather than strong airflow is the objective)
	Containment Measures Note in most cases these measures can cease 14 days (1 incubation period) after most recent case		New admissions restricted Transfers restricted Restricted communal activities Staffing precautions Visitor access restrictions
Post Outbreak	Declaration of end of outbreak		As advised by Public Health
	Final evaluation	Review of management of outbreaks and lesson learned	Coordination with Public Health and OCT if this was convened

## Appendix B: Details for line listing

1. Outbreak code (on top of line list as title);
2. Name of case;
3. Case ID;
4. Location (unit/section);
5. Date of birth/age;
6. Sex;
7. Status i.e., resident, staff member, volunteer, visitor;
8. Vaccination status of resident, staff member, volunteer, visitor; vaccine protection;
9. Previous Covid-19 infection (last 9 months) [For COVID-19 outbreaks];
10. Date of onset of symptoms;
11. Date of notification of symptoms;
12. Clinical symptoms (outline dependent on case definition) e.g., fever, cough, myalgia, headache, other;
13. Samples taken and dates;
14. Laboratory results including test type e.g., RT-PCR;
15. Date when isolation of resident was started;
16. Date of recovery;
17. Duration of illness;
18. Outcomes: recovery, pneumonia, other, hospitalisation, death;
19. Also include work assignments of staff and last day of work of ill staff member;
20. State if staff worked in other facilities;

**Have separate sheets for both staff and residents**



### Appendix C: Part 2 –Residents ONLY

Name of Facility: ..... Name of Outbreak: ..... Outbreak Code.....

Test Results		Outcome				
ID	Laboratory Test Done Yes/No, If yes, date:	Type of Test and Result	Pneumonia	Hospitalisation (Date)	Death (Date)	Recovered to pre-outbreak health status. Yes/No. If Yes, date:

Key: (Y =Yes, N=No, U=Unknown)





**Appendix C: Part 4 –Staff ONLY\***

Name of Facility: ..... Name of Outbreak: ..... Outbreak Code:.....

ID	Test Results		Outcome				Work exclusion
	Pathology Test Done Yes/No, If yes, date:	Type of Test and Result	Pneumonia	Hospitalisation (Date)	Death (Date)	Recovered to pre-outbreak health status. Yes/No. If Yes, date:	Excluded from work until (Date)

Key: (Y = Yes, N = No, U = Unknown)

## Appendix D: Checklist for outbreak management


	Discussion point	Decision/action to be taken (date completed)	Person responsible
1	Declare an outbreak and convene an OCT following Public Health risk assessment		
2	Agree the chair		
3	Formulate an outbreak code and working case definition		
4	Define the population at risk		
5	Active case finding, request line listing of residents and staff from the RCF		
6	Discuss whether it is a facility-wide outbreak or unit-specific		
7	Confirm how and when communications will take place between the RCF, CIPCN, CHO NH lead, Public Health and the laboratory		
8	Review the control measures (infection prevention and control necessary to prevent the outbreak from spreading). Confirm that the management of the facility is responsible for ensuring that agreed control measures are in place and enforced		
9	Review vaccination status of all residents and staff		
10	Discuss which specimens have been collected. Notify the laboratory of the investigation.		
11	Confirm the type and number of further laboratory specimens to be taken. Clarify which residents and staff should be tested.		
12	Confirm with the laboratory that it will phone or fax results (both positive and negative) directly to the requesting doctor and that this person will notify Public Health. Review the process for discussing laboratory results with the RCF's designated officer.		
13	Liaise with the RCF and laboratory regarding specimen collection and transport		


	<b>Discussion point</b>	<b>Decision/action to be taken (date completed)</b>	<b>Person responsible</b>
14	Identify persons/institutions requiring notification of the outbreak e.g. families of ill or all residents of the facility; health care providers e.g. GPs, physiotherapists etc.; infectious disease consultants, consultant microbiologists, infection prevention & control specialists, Emergency Departments; local hospitals, other RCF, HPSC		
15	Discuss whether a media release is required		
16	Ensure that the incident is promptly reported to HPSC and surveillance details entered onto CIDR		
17	Provide updates on the investigation to the Assistant National Director, ISD-Health Protection when/if required		
18	Discuss communication arrangements with HSE management ± HSE crisis management team		
19	Discuss communication arrangements with local GPs and Emergency Departments		
20	Decide how frequently the OCT should meet and agree criteria to declare outbreak over		
21	Prepare/circulate an incident report/set date for review meeting		

## Appendix E Hand Hygiene poster

### How to hand wash

Wash hands when visibly soiled. Otherwise, use handrub with hand sanitiser.

 Length of time to spend washing: 40-60 seconds




Wet hands with water



Apply enough soap to cover all hand surfaces




Rub hands palm to palm



Right palm over the back of the left hand with interlaced fingers and do same on other hand



Palm to palm with fingers interlaced



Backs of fingers to opposite palm with fingers interlocked



Rotational rubbing of left thumb clasped in right palm and do same on other hand




Rub in a circle with clasped fingers of right hand in left palm do same on other hand



Rinse hands with water



Dry hands thoroughly with a clean towel or single use towel



Use single use towel or piece of tissue to turn off tap




Your hands are now safe



## Appendix F Donning and Doffing PPE

A full range of resources including posters, videos and webinars relating to the safe donning and doffing of PPE is accessible [here](#)

# Coronavirus COVID-19








## Guide to donning and doffing standard Personal Protective Equipment (PPE)

### FOR HEALTH AND SOCIAL CARE SETTINGS

#### DONNING OR PUTTING ON PPE







- Before putting on the PPE, perform hand hygiene.
- Be well hydrated and have taken a toilet break
- Have removed all jewellery including earrings
- Be bare below the elbows
- Have secured your hair back off your face
- Do not bring mobile phones/bleeps into an isolation area

**CLICK HERE FOR FIT CHECK GUIDE!**

- 1** Put on your plastic apron, making sure it is tied securely at the back.  

- 2** Put on your surgical or FFP2 mask. For mask with ties - tie the upper straps on top of head and bring the lower straps up in front of the ears and tie on top of head. For mask with loops - loop straps over the ears. Mould the metal strap over the bridge of the nose and make sure the mask is extended to cover your mouth and chin.  

- 3** Put on your eye protection if there is a risk of splashing.  

- 4** Put on non-sterile nitrile gloves.  

- 5** You are now ready to enter the patient/resident area.  


#### DOFFING OR TAKING OFF PPE

Surgical or FFP2 mask may be used for single session use but gloves and apron must be changed between patients/residents. Gloves and aprons must be changed between patients/residents or between care activities as appropriate.

- 1** Remove gloves, grasp the outside of the cuff of the glove and peel off, holding the glove in the gloved hand, insert the finger underneath and peel off second glove.  

- 2** Perform hand hygiene.  

- 3** Remove eye protection.  

- 4** Snap or unfasten the neck ties and allow to fall forward. Snap waist ties and fold apron in on itself, do not touch the outside as it is contaminated, and put into Healthcare risk waste.  

- 5** Once outside the patient room or cohort area, remove facemask.  

- 6** Perform hand hygiene.  


## Appendix G Transmission based precautions

Table 2: Personal Protective Equipment (PPE) Requirements by Precaution Type

Personal Protective Equipment			
Precaution	Contact	Droplet	Airborne
<b>Gloves</b>	Yes	As per Standard Precautions	As per Standard Precautions
<b>Gown/Apron (impermeable)</b>	When healthcare worker's clothing is in substantial contact with the resident, items in contact with the resident, and their immediate environment	As per Standard precautions	Gown
<b>Surgical Mask</b>	When adequate distance cannot be maintained	Yes	No
<b>Respirator Mask (FFP or equivalent)</b>	Not required	Not required (see text regarding access to respirator mask when caring for people with COVID)	Yes
<b>Goggles/face Shield</b>	Not required	As per standard precautions	Yes

## Appendix H Admissions, transfers and discharges to and from residential care facilities

### **Note regarding testing and restricting movement of People Pre-transfer/Admission to a LTRCF**

The requirement for testing should now apply to all people before transfer or admission including those who are fully vaccinated or who have had COVID-19 in the previous 9 months.

In the context of a LTRCF with a high level of vaccine protection there is no requirement to limit the movement of a resident within the LTRCF after return from an outing or hospital attendance regardless of the duration of the absence unless some significant and unanticipated exposure risk occurred or there is a specific public health or IPC recommendation that requires limitation of movement.

In the absence of a high level of vaccination non-vaccinated residents absent from the LTRCF for more than 12 hours should be advised to limit their contact with other residents. Such non-vaccinated residents should be offered testing on day 5 after their return and if they test not-detected and are asymptomatic they may return to normal activities at that time. For those who are fully vaccinated in such a LTRCF there is no requirement to limit the movement after return from an outing or hospital attendance regardless of the duration of the absence unless some significant and unanticipated exposure risk occurred or there is a specific public health or IPC recommendation that requires limitation of movement.

### **Introduction**

Long-term residential care facilities (LTRCF) are a critical part of health and social care services. LTRCFs should put in place clear processes that facilitate the return of residents from an acute setting and the admission of new residents, where it is clinically safe to do so.

It is recognised that accepting admission or transfer of residents poses a risk of introducing COVID-19, even where processes to manage the risks are in place however it is essential that this risk is balanced against the consequences of restricting access to a facility/service or disproportionately impacting on the wellbeing of residents. The risk of harm from introduction of COVID-19 is greatly reduced when residents and staff who are fully vaccinated and have had booster vaccination.

Given the uptake of vaccination in the population most residents transferring to a LTRCF are likely to be fully vaccinated and have had booster vaccine. If an unvaccinated person is transferring from an acute hospital they should generally have had the first dose of vaccine before transfer. While the vaccine should ideally be administered as long as possible in advance of transfer, there is no requirement to delay transfer of a person who is otherwise ready for discharge to allow time for an immune response to the vaccine. Arrangements to complete the vaccination in the LTRCF are essential.



In all instances, careful attention to standard precautions will assist in minimising risk of infection to residents and staff. Key elements include; hand hygiene, respiratory hygiene and cough etiquette, use of personal protective equipment (PPE), for example wearing disposable gloves when in contact with blood or other body fluids (other than sweat), non-intact skin or mucus membranes and regular environmental cleaning.

It is essential that residents and clients and their significant persons are informed of the issues and risks of decisions related to their care and that their preferences are taken into account in applying this guidance.

### **Background on testing for COVID-19**

**The key point about testing is that interpretation is not straightforward**

- 1. A test result that says not-detected or “negative” does not prove the person is not infectious to others**
- 2. A test result that says a virus is detected does not prove the person is still infectious to others**

Over the course of the COVID-19 pandemic, there has been significant learning about the role of testing for COVID-19 and its role in determining levels of asymptomatic infection and tracking spread of infection, especially in congregated settings, such as LTRCF.

A single test may be reported as not-detected or “negative” in a substantial proportion of people with infection. The test is more likely to miss infection in people with pre-symptomatic or asymptomatic infection. Therefore, a not-detected or “negative” test makes COVID-19 infection less likely, but it does not prove the person is not infected.

Equally, for those who have been infected and infectious with COVID 19, a continued positive test result does not mean they are still infectious to others. Some people have a positive test for weeks after onset of symptoms, but latest evidence shows they are very unlikely to spread infection. **For people with a diagnosis of COVID-19 infection who are in a RCF or are planning to move into a RCF, the period of isolation is 14 days after onset of symptoms with no fever for the last five of this period.**

Note that if a person is detected by testing and subsequently develops symptoms the 14 days is counted from the date of symptom onset (not the sample date); however if they do not develop symptoms the 14 days is counted from the sample date.

The period of 14 days continues to apply in this setting although the infectious period is now 10 days with no fever for the last 5 days for people who do not require hospitalisation for care of COVID-19 and who are not resident in LTRCF. Note that repeat testing at the end of the isolation period is generally not appropriate though exceptions may arise in the context of discussion with Microbiology, Infectious Diseases or Public Health Clinicians.

### **The role of COVID-19 testing in assisting with decision-making regarding transfers to congregated settings**

1. People who plan to live in a LTRCF should be tested for SARS-CoV-2 before they move to the LTRCF. This applies to transfer from a hospital or another LTRCF or to people admitted from the community. This is to help identify most of those who have infection, but it will not detect all of those with infection.
2. Testing, , should be performed within three days of planned admission to the LTRCF if transfer is from a hospital or another LTRCF and should also be performed whenever possible for admissions from the community.
3. Where testing is required and is not performed before admission, it should be carried out within one day of admission.
4. Irrespective of testing, all residents should be assessed before admission to determine if they have clinical symptoms suggestive of COVID-19 or if they are known COVID-19 contacts
5. People who are COVID-19 contacts may transfer to a LTRCF if they have been tested by PCR and SARS-CoV-2 virus was not detected. If they are not-fully vaccinated they should restrict their movement within the LTRCF for 14 days. If they are fully vaccinated they should restrict their movements for 10 days.
6. The requirement for testing in advance of transfer or admission may be waived if appropriate to the needs of an individual person transferring to settings caring for children under the age of 18
7. The requirement for testing in advance of transfer or admission does not apply to people who are returning to supported/assisted living or small group homes (generally less than five residents) following discharge from hospital, where the facility is more reflective of a household setting
8. For patients or residents who decline or are clearly distressed by collection of a nasopharyngeal sample a deep nasal sample (also known as a mid-turbinate swab) is often less uncomfortable. Deep nasals swabs should generally be used for surveillance testing on people who require very frequent testing and for those in whom a nasopharyngeal sample collection is difficult or distressing. An anterior nasal swab is NOT a suitable sample. Some residents may decline testing, or may find the process too distressing and that testing may not be appropriate in every situation (Refer to DOH [Guidance](#) on Ethical Considerations Relating to Long-Term Residential Care Facilities in the context of COVID-19).

## Procedure for Testing of People Pre-transfer/Admission to a LTRCF

1. If a person is being transferred from an acute hospital to a LTRCF, the hospital should arrange for the person to be swabbed in the three days before transfer.
2. If a person is being admitted to the LTRCF from home, where possible, the GP should arrange for the person to be swabbed within the three days before admission. This can be done using Healthlink. If the person cannot travel to the test centre, a home test can be ordered by clicking on the 'no transport available' option as shown on the screenshot below (Figure 1).
3. If a test pre-admission cannot be arranged, including for urgent admissions, the person should be admitted as planned. **If the person is not fully vaccinated**, the person will need to be isolated, with full contact and droplet precautions until the result of the test is available. The facility can arrange swabbing after admission. This can be done by the person's own GP or the GP/Medical Officer who provides medical care for the residents in the facility. If the sample is reported not detected, the precautions that apply are those that apply to a person with sample reported not-detected before admission (see below).

The screenshot shows a Microsoft Remote Desktop window displaying the Healthlink web application. The browser title is 'Socrates (2.7.3.4) - Dr Nuala O'Connor Logged On - ELMWOOD MEDICAL PRACTICE'. The page is titled 'Consultation' and shows a form for a COVID-19 test referral. The form includes the following fields and options:

- Patient's GP:** [Text input field]
- GP Telephone:\*** (preferably mobile): 021 4893255
- Referral Category:\***
  - General Covid-19 Test
  - Healthcare Worker
  - Close Contact of Confirmed Case
  - At Risk Group
- Covid-19 Symptomatic:\***
  - Yes
  - No
- Date of Last Test:\*** (where applicable): [Date picker]
- Transport:\***
  - Transport available
  - No transport available
  - Unable to travel for medical reasons
  - Unable to travel for personal reasons
  - Other (please specify)
- Additional Details:**
  - Hearing Loss
  - Can Read Sign Language
  - Visually Impaired
  - Other (please specify)
- Additional Relevant Information:** [Text input field]

At the bottom of the form, it says 'Login Complete'.

*Figure 1. Snapshot of Health link web page*

## Requirements for placement and restricted movement of the person as part of transfer protocols Planning

1. All LTRCF should review their accommodation to identify areas where new residents can safely restrict their movement. It is understood that the creation of such areas may be

constrained by existing accommodation availability (e.g., rooms already in use by existing residents).

2. Where possible the use of single rooms in LTRCF with significant numbers of multi-occupancy rooms should be prioritised for new transfers and admissions from community or other healthcare facilities (acute hospital or other LTRCF), regardless of the pre-admission COVID-19 test result.
3. For those LTRCF providing a blend of longer-term nursing home and short-term respite or convalescence care, it is advised to consider where the longer and shorter-term residents will be accommodated and where it is feasible, to try and place residents for shorter-term accommodation in an area separate to those for longer-term accommodation.
4. The identification of space for the period of restricted movement needs to be managed carefully with residents, families and others. Existing residents should not be required to move from their room / accommodation in order to facilitate the creation of new areas to facilitate transfers.
5. Careful consideration should also be given to the consequences of closing facilities/rooms within a service for the purpose of having an isolation area should a need arise – the potential harms of such decisions should be balanced against the likely requirement.
6. All transfers or new admissions should have a risk assessment, to ensure sufficient resources are available within the LTRCF to support physical distancing and placement of residents.
7. Any person who is not fully vaccinated and has not been diagnosed with COVID-19 in the previous 9 months will need to restrict their movements for 14 days after transfer regardless of the test result.
8. If the person is fully vaccinated or if the person has had COVID-19 in the past 9 months they do not normally need either testing or to restrict their movement after transfer unless they are a COVID-19 Contact. In that case the period of restricted movement is 10 days.
9. Public Health may recommend that a person who is transferring from a particular congregated healthcare setting (a hospital or RCF) where there is evidence of ongoing transmission of COVID-19 (one or more open outbreaks) is managed as a COVID-19 Contact after transfer based on risk assessment. In this context Public Health may also recommend a repeat test at 5 days to enhance detection of hospital acquired infection.
10. For those who require restricted movement the following applies:
11. Care delivered within the single room of a person on restricted movement can be delivered with Standard Precautions plus surgical mask and the resident may leave their room as per guidance below on transfers.
12. The resident is not required to remain in strict isolation, but should practice restricted movement:
13. The resident may leave their room, but should remain separated from other residents (e.g. to go the garden or for a short walk)

14. A move to a multi-occupancy room (where this is the planned accommodation in the longer term for the resident) will be appropriate after the 14-day period, once the resident is symptom free and there is no evidence of infection in residents within the room it is proposed for the resident to move to.

### **Transfer of people with COVID-19**

1. Any resident transferred to a LTRCF before the 14 days have elapsed since date of onset of symptoms or date of first positive test (if symptom onset undetermined/asymptomatic), must be isolated with transmission-based precautions up to day 14 on return to the LTRCF. Such transfer should not proceed if the receiving LTRCF has no other residents with infectious COVID-19 at the time. Provided the resident has remained afebrile for the last five of the 14 days, the resident is generally no longer infectious to others after day 14 has elapsed. However if the hospital practice requires repeat testing or extended isolation for the patient for a specific reason the same level caution applies in advance of transfer to the LTRCF.
2. In particular, existing residents from a LTRCF who require transfer to hospital from the LTRCF for assessment or care related to COVID-19 acquired in the LTRCF should be allowed to transfer back to that LTRCF following assessment / admission, if clinically fit for discharge and risk assessment with the facility determines there is capacity for them to be cared for there, with appropriate isolation and where that transfer represents the most appropriate place of care for the resident (e.g. ongoing need for palliative care).
3. If the resident in an LTRCF has been diagnosed with COVID-19 while in hospital, it is important to assess if the person was infected in the LTRCF before transfer to the hospital or if this is a hospital-acquired infection. If it is likely that infection was acquired in hospital and there are no other known cases of COVID-19 in the LTRCF, transfer back to the LTRCF should be delayed until the resident is no longer infectious to others.
4. The public health team should be notified immediately where newly-diagnosed COVID-19 is assessed as acquired within a LTRCF.
5. In all instances the discharging hospital should provide the LTRCF with the following information on the arrival of the resident:
  - a. The date and results of COVID-19 tests (including dates of tests reported as not-detected);
  - b. The date of onset of any symptoms of COVID-19;
  - c. Date of last documented fever while in hospital (particularly important where resident is being transferred to RCF within 14 days of COVID-19 diagnosis);
  - d. Details of any follow-up treatment or monitoring require.

### **Residents who become symptomatic during their stay in the LTRCF**

1. Following transfer/admission to a LTRCF, the resident should be evaluated by their doctor if they become symptomatic, including changes in the resident's overall clinical condition and a further viral swab for SARS-CoV-2 sent for testing. **They may also require testing for other viruses in particular influenza virus;**
2. The rationale for this recommendation is that, in the context of a pandemic, there may have been contact between the resident and HCW or other people who may have had COVID-19 infection, but who may have been in the pre-symptomatic incubation period or have had minimal symptoms/been asymptomatic at the time. In that case, there would be an associated risk of unrecognised onward transmission to the resident. They may also have been exposed to other respiratory viruses.

### **Cessation of new admissions to a facility during an outbreak of COVID-19 in a LTRCF**

1. Following the declaration of an outbreak within a LTRCF, admissions of new residents to the facility (i.e. residents not previously living in the LTRCF) should be suspended until at least 1 incubation period (14 days) after the most recent case and Public Health have advised that admissions may resume;
2. Residents normally cared for in the LTRCF who are admitted to hospital while an outbreak is ongoing in the LTRCF may have their discharge to the same LTRCF facilitated if it is deemed to be clinically appropriate and a risk assessment has been carried out which identifies that the resident can be isolated and the facility has capacity to manage their care needs and where that transfer represents the most appropriate place of care for the resident (e.g. ongoing need for palliative care).

### **Transfers from LTRCF to an acute hospital**

1. COVID-19 positive status must not significantly delay transfer to an acute hospital, where it is deemed clinically appropriate. The national ambulance service (NAS) and the local receiving hospital must be informed by the LTRCF, in advance of transfer of any COVID-19 positive or suspected COVID-19 resident AND where there is a suspected or confirmed COVID-19 outbreak in the LTRCF;
2. People with COVID-19 do not require to be hospitalised for the 14 days when they are infectious if they are clinically fit for discharge, if infection was acquired in the LTRCF or if the LTRCF already has cases of COVID-19 and the LTRCF has appropriate facilities and capacity for isolation and can support care;
3. Residents do not require isolation on return to their LTRCF following hospital transfer to facilitate short investigations (e.g., diagnostics, haemodialysis, radiology, outpatient appointment);
4. Residents who are not fully vaccinated and have not had COVID-19 in the previous 9 months and are absent from the LTRCF for more than 12 hours should be advised to limit their

contact with other residents on their return. Such non-vaccine protected residents and those who have not had COVID-19 in the previous 9 months should be offered testing on day 5 after their return and if they test not-detected and are asymptomatic they may return to normal activities at that time;

5. Residents who are fully vaccinated or who have had COVID-19 in the previous 9 months will not need either testing or to restrict their movements on return to their LTRCF from an outing or hospital attendance (as distinct from hospital admission) regardless of the duration of the absence unless some significant and unanticipated exposure risk occurred or there is a specific public health or IPC recommendation that requires limitation of movement.

**Table Transfer/admission of a resident to a LTRCF**

CLINICAL SCENARIO	RECOMMENDED PRECAUTIONS ON ARRIVAL TO LTRCF	PRE-ADMISSION TEST FOR SARS-CoV-2 (COVID-19)	TIMING OF TRANSFER TO LTRCF	DAY OF TRANSFER
<p><b>CONFIRMED COVID-19 &amp; will be still infectious to others</b> on planned date of transfer</p> <p>(less than 14 days since onset/test date)</p>	<p>Transmission-based Precautions until 14 days reached and has been afebrile for last five of those days</p>	<p>Not required, as already confirmed COVID-19</p>	<p><b>LTRCF has other resident(s) with COVID-19:</b> Transfer when fit for discharge to LTRCF AND provided LTRCF can meet care needs</p> <p><b>LTRCF has no other resident with COVID-19</b> - Remain in hospital until no longer infectious to others</p>	<p>Confirm date of onset/first positive test result</p> <p>Confirm date last febrile</p>
<p><b>CONFIRMED COVID-19 in past 9 months &amp; no longer infectious to others</b> (more than 14 days since onset/test date and afebrile for last five of those days)</p>	<p>No requirement for Transmission based Precautions or restricted movement</p>	<p>Test within the 3 days prior to scheduled transfer date</p>	<p>When fit for discharge to LTRCF</p>	<p>Confirm date of onset/first positive test result is more than 14 days ago and was afebrile for last five days of that</p>
<p><b>FULLY VACCINATED ASYMPTOMATIC &amp; NOT A CONTACT</b></p>	<p>No requirement for Transmission based Precautions or restricted movement <b>[may be exceptions based on risk assessment]</b></p>	<p>Test within the 3 days prior to scheduled transfer date</p>	<p>When fit for discharge to LTRCF</p>	<p>Confirm details of vaccination</p>
<p><b>NOT FULLY VACCINATED AND NO PRIOR CONFIRMATION OF COVID-19 IN THE PREVIOUS 9 MONTHS ASYMPTOMATIC &amp; A RECENT CONTACT</b></p>	<p>Restricted-movement until 14 days reached. Single room accommodation with monitoring for symptoms until 14 days reached. Standard Precautions plus surgical face mask.</p>	<p>Test within the 3 days prior to scheduled transfer date <b>[public health may advise an additional test on day 5 post transfer]</b></p>	<p>Test result- not-detected</p> <p>LTRCF can meet care needs</p>	<p>Confirm test result received</p> <p>Ensure no new symptoms</p> <p>LTRCF to repeat test day 5 after most recent exposure</p>



CLINICAL SCENARIO	RECOMMENDED PRECAUTIONS ON ARRIVAL TO LTRCF	PRE-ADMISSION TEST FOR SARS-CoV-2 (COVID-19)	TIMING OF TRANSFER TO LTRCF	DAY OF TRANSFER
<b>FULLY VACCINATED OR COVID-19 IN THE PREVIOUS 9 MONTHS ASYMPTOMATIC &amp; A RECENT CONTACT</b>	Restricted movement (as above) until 10 days after most recent exposure	Test within the 3 days prior to scheduled transfer date. <b>[public health may advise an additional test on day 5 post transfer]</b>	Test result- not-detected  LTRCF can meet care needs	Confirm test result received  Ensure no new symptoms
<b>NOT FULLY VACCINATED AND NO PRIOR CONFIRMATION OF COVID-19 IN THE PREVIOUS 9 MONTHS ASYMPTOMATIC &amp; NOT A CONTACT</b>	Restricted movement (as above) until 14 days after most recent exposure	Test within the 3 days prior to scheduled transfer, <b>[public health may advise an additional test on day 5 post transfer]</b>	Test result- not-detected  LTRCF can meet care needs	Confirm test result received  Ensure no new symptoms
<b>NO PRIOR CONFIRMATION OF COVID-19 or COVID-19 MORE THAN nine months previously &amp; test result available before transfer But Test result is NOT available prior to admission</b>	Transmission-based Precautions until test result is available then follow as per immediately above	Test within one day of admission <b>[public health may advise an additional test on day 5 post transfer]</b>	When fit for discharge to LTRCF	Take sample for COVID-19 test. Ensure no symptoms and not newly identified contact of a COVID-19 case