





Weekly report on COVID-19 deaths reported in Ireland

Report produced by Health Protection Surveillance Centre on 26/01/2022

This report includes data for COVID-19 cases notified on CIDR up to and including midnight on 25/01/2022 who have been notified as COVID-19 deaths

Note: Data were extracted from Computerised Infectious Disease Reporting (CIDR) system on 26/01/2022 and are provisional and subject to ongoing review, validation and update. As a result, figures in this report may differ from previously reported figures.

Table 1: Summary characteristics of COVID-19 deaths notified in Ireland with date of death from 19/01/2022 to 25/01/2022

Characteristics		Number of deaths
Total number of deaths		13
Age	Mean age (Years)	71
	Median age (Years)	72

The number of deaths described in the above table relate only to COVID-19 cases who died within this time period and whose death has been reported to CIDR up to 26/01/2022. It does not include deaths of COVID-19 cases reported to CIDR in the last week but with an earlier date of death. It also does not reflect the final number of deaths occurring for this period as the outcome may not yet have occurred, or is yet to be reported to CIDR.

Table 2: Summary characteristics of COVID-19 deaths in Ireland, cases with a date of notification from 01/03/2020 to 25/01/2022

Characteristics		Number of Deaths	Percentage
Total number of deaths		6,136	
Sex	Female	2,840	46.3
	Male	3,295	53.7
	Unknown	1	0.0
	M:F ratio	1.16	
Case classification*	Possible	178	2.9
	Probable	97	1.6
	Confirmed	5,861	95.5
Healthcare Worker	Yes	21	0.3
	No	5,020	81.8
	Unknown	1,095	17.8
Underlying Conditions	Yes	5,149	83.9
	No	514	8.4
	Unknown	473	7.7
ICU Admission	Yes	784	12.8
	No	5,352	87.2

^{*} Case definition

Table 3: Number and percentage of COVID-19 deaths and mortality rate per 100,000 in Ireland by age group, cases with a date of notification from 01/03/2020 to 25/01/2022

Characteristics		Number of Deaths	Percentage	Mortality rate per 100,000
Age	Median Age (Years)	82		
	Mean Age (Years)	80		
Age group	<25 yrs	8	0.1	0.4
	25-34 yrs	19	0.3	2.9
	35-44 yrs	51	0.8	6.8
	45-54 yrs	157	2.6	25.1
	55-64 yrs	398	6.5	78.2
	65-74 yrs	1,036	16.9	277.4
	75-84 yrs	2,038	33.2	1037.1
	85+ yrs	2,426	39.5	3591.1
	Unknown	3	0.0	
	Total	6,136	100	128.9

Please note that due to the small number of persons who died due to COVID-19 in the age groups 0-25 years this information has been aggregated in order to protect patient confidentiality

Table 4: Number and percentage of COVID-19 deaths and mortality rate per 100,000 in Ireland by county of notification, cases with a date of notification from 01/03/2020 to 25/01/2022

County	Number of deaths	Percentage	Mortality rate per 100,000
Carlow	83	1.4	145.8
Cavan	132	2.2	173.3
Clare	150	2.4	126.2
Cork	532	8.7	98.0
Donegal	235	3.8	147.6
Dublin	2,115	34.5	157.0
Galway	174	2.8	67.4
Kerry	103	1.7	69.7
Kildare	368	6.0	165.4
Kilkenny	98	1.6	98.8
Laois	100	1.6	118.1
Leitrim	30	0.5	93.6
Limerick	294	4.8	150.8
Longford	44	0.7	107.7
Louth	233	3.8	180.8
Mayo	230	3.7	176.2
Meath	194	3.2	99.5
Monaghan	108	1.8	175.9
Offaly	82	1.3	105.2
Roscommon	77	1.3	119.3
Sligo	45	0.7	68.7
Tipperary	131	2.1	82.1
Waterford	139	2.3	119.6
Westmeath	94	1.5	105.9
Wexford	174	2.8	116.2
Wicklow	171	2.8	120.1

Table 5: Place of death for COVID-19 deaths in Ireland, cases with a date of notification from 01/03/2020 to 25/01/2022

Place of death	Number of deaths	Percentage	
Hospital	3,237	52.8	
Residential Institution*	2,063	33.6	
Hospice	62	1.0	
Home	272	4.4	
Other	94	1.5	
Unknown	408	6.6	

^{*}Residential institution includes: community hospital/long stay unit, homeless facility, mental health facility and nursing homes.

Table 6: Summary of COVID-19 deaths linked to outbreaks in Ireland, cases with a date of notification from 01/03/2020 to 25/01/2022

		Number of Deaths	Percentage of total deaths	Percentage of deaths linked to outbreaks
Total number of deaths		6,136	100.0	
Deaths linked to outbreaks	3	3,692	60.2	100.0
Outbreaks by location	Nursing homes	2,271	37.0	61.5
	Hospital	885	14.4	24.0
	Community Hospitals/Long-stay units	153	2.5	4.1
	Residential institutions	112	1.8	3.0
	Other locations*	271	4.4	7.3

^{*} Other locations include community outbreak, extended family, hotel, other, other healthcare service, private house, public house, religious/other ceremony, workplace

A death linked to an outbreak in a particular setting does not of itself indicate that transmission occurred within that setting. A case may be detected as part of an outbreak investigation, and associated with an outbreak, despite exposure and transmission having occurred elsewhere.

Table 7: Number of COVID-19 deaths in Ireland by month of death, cases with a date of notification from from March 2020 to January 2022

Year	Month	Number of deaths *
2020	March	129
2020	April	1,160
2020	May	363
2020	June	67
2020	July	17
2020	August	7
2020	September	39
2020	October	132
2020	November	190
2020	December	195
2021	January	1,419
2021	February	892
2021	March	258
2021	April	102
2021	May	43
2021	June	21
2021	July	22
2021	August	89
2021	September	181
2021	October	226
2021	November	249
2021	December	203
2022	January	101

^{*} Date of death reported for 6,105 of the deaths. Number of deaths for January 2022 is incomplete.




Figure 1: Total number of COVID-19 deaths in Ireland and cumulative number by week of death, cases with a date of notification from 01/03/2020 to 25/01/2022. Date of death reported for 6,105 of deaths.

Acknowledgments

Sincere thanks are extended to all those who are participating in the collection of data and reporting of data used in these reports. This includes the HSE COVID-19 Contact Management Programme (CMP), staff in ICU units, notifying clinicians, laboratory staff, public health doctors, nurses, surveillance scientists, microbiologists and administrative staff.

Report prepared by COVID-19 Epidemiology Team, HPSC

Technical Notes

1. Data Source

Data are based on statutory notifications and were extracted from Computerised Infectious Disease Reporting (CIDR) system at 08:27 on 26/01/2022. Data are provisional and subject to ongoing review, validation and update. As a result, figures in this report may differ from previously published figures.

2. Population Data

Population data were taken from Census 2016. Data were aggregated into the following age groups for the analysis of deaths for the entire pandemic: 0-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years and \geq 85 years. Data were aggregated into the following age groups for the analysis of deaths with date of death in the past 7 days: 0-64 years, 65-74 years, 75-84 years and \geq 85 years.

3. Definition of a COVID-19 death used by HPSC

For surveillance purposes, COVID-19 deaths include deaths in all possible, probable and confirmed COVID-19 cases (as per the COVID-19 case definition) and all should be notified, unless there is a clear alternative cause of death that cannot be related to COVID-19 infection (e.g. trauma). There should be no period of complete recovery* from COVID-19 between the illness and death. All COVID-19 deaths are notified regardless of the setting, including home, community and hospital settings. HPSC reports all deaths among these COVID-19 cases as outlined above and does not just confine the death reporting to those who die within 28 days of a positive test. This is in line with how COVID-19 cases are reported by the majority of European countries and follows WHO guidance for COVID-19 death surveillance.

*Please note that discharge from ICU or hospital is not in itself evidence of recovery. To determine if the case had recovered, it should be based on clinical assessment or alternatively a period or ≥ 3 months must have elapsed since the case was initially diagnosed with COVID-19 and the case must have no evidence of COVID-19 infection prior to death as assessed by a clinician.

Deaths not reported as COVID-19 deaths:

- 1. Persons with COVID-19 may die directly due to accidents. Such deaths are not due to COVID-19 and should not be certified as such. This decision not to certify as COVID-19 death will be based on clinical judgement.
- 2. In some instances, a death due to COVID-19 may not be attributed to another disease (e.g. cancer) and would be counted as a COVID-19 death independently of pre-existing conditions that are suspected of triggering a severe course of COVID-19.

More resources:

COVID-19 interim case definition

Epidemiology of COVID-19 in Ireland Frequently Asked Questions