



Weekly report on COVID-19 deaths reported in Ireland

Report produced by Health Protection Surveillance Centre on 05/01/2022

This report includes data for COVID-19 cases notified on CIDR up to and including midnight on 04/01/2022 who have been notified as COVID-19 deaths

Note: Data were extracted from Computerised Infectious Disease Reporting (CIDR) system on 05/01/2022 and are provisional and subject to ongoing review, validation and update. As a result, figures in this report may differ from previously reported figures.

Notification of deaths to HPSC may be impacted by the extended Christmas and New Year weekends.

Table 1: Summary characteristics of COVID-19 deaths notified in Ireland with date of death from 29/12/2021 to 04/01/2022

Characteristics	Number of deaths	
Total number of deaths	11	
Age	Mean age (Years)	77
	Median age (Years)	75

The number of deaths described in the above table relate only to COVID-19 cases who died within this time period and whose death has been reported to CIDR up to 05/01/2022. It does not include deaths of COVID-19 cases reported to CIDR in the last week but with an earlier date of death. It also does not reflect the final number of deaths occurring for this period as the outcome may not yet have occurred, or is yet to be reported to CIDR.

Table 2: Summary characteristics of COVID-19 deaths in Ireland, cases with a date of notification from 01/03/2020 to 04/01/2022

Characteristics		Number of Deaths	Percentage
Total number of deaths		5,952	
Sex	Female	2,765	46.5
	Male	3,187	53.5
	Unknown	0	0.0
	M:F ratio	1.15	
Case classification*	Possible	176	3.0
	Probable	97	1.6
	Confirmed	5,679	95.4
Healthcare Worker	Yes	20	0.3
	No	4,964	83.4
	Unknown	968	16.3
Underlying Conditions	Yes	5,032	84.5
	No	504	8.5
	Unknown	416	7.0
ICU Admission	Yes	747	12.6
	No	5,205	87.4

* [Case definition](#)

Table 3: Number and percentage of COVID-19 deaths and mortality rate per 100,000 in Ireland by age group, cases with a date of notification from 01/03/2020 to 04/01/2022

Characteristics		Number of Deaths	Percentage	Mortality rate per 100,000
Age	Median Age (Years)	82		
	Mean Age (Years)	80		
Age group	<25 yrs	8	0.1	0.4
	25-34 yrs	18	0.3	2.7
	35-44 yrs	47	0.8	6.3
	45-54 yrs	144	2.4	23.0
	55-64 yrs	378	6.4	74.3
	65-74 yrs	991	16.6	265.3
	75-84 yrs	1,981	33.3	1008.1
	85+ yrs	2,382	40.0	3526.0
	Unknown	3	0.1	
	Total	5,952	100	125.0

Please note that due to the small number of persons who died due to COVID-19 in the age groups 0-25 years this information has been aggregated in order to protect patient confidentiality

Table 4: Number and percentage of COVID-19 deaths and mortality rate per 100,000 in Ireland by county of notification, cases with a date of notification from 01/03/2020 to 04/01/2022

County	Number of deaths	Percentage	Mortality rate per 100,000
Carlow	80	1.3	140.5
Cavan	128	2.2	168.0
Clare	148	2.5	124.6
Cork	512	8.6	94.3
Donegal	229	3.8	143.9
Dublin	2,057	34.6	152.7
Galway	168	2.8	65.1
Kerry	96	1.6	65.0
Kildare	361	6.1	162.2
Kilkenny	93	1.6	93.7
Laois	97	1.6	114.5
Leitrim	30	0.5	93.6
Limerick	283	4.8	145.2
Longford	42	0.7	102.8
Louth	229	3.8	177.7
Mayo	228	3.8	174.7
Meath	188	3.2	96.4
Monaghan	106	1.8	172.7
Offaly	81	1.4	103.9
Roscommon	75	1.3	116.2
Sligo	44	0.7	67.1
Tipperary	125	2.1	78.3
Waterford	126	2.1	108.5
Westmeath	91	1.5	102.5
Wexford	165	2.8	110.2
Wicklow	170	2.9	119.4

Table 5: Place of death for COVID-19 deaths in Ireland, cases with a date of notification from 01/03/2020 to 04/01/2022

Place of death	Number of deaths	Percentage
Hospital	3,112	52.3
Residential Institution*	2,034	34.2
Hospice	60	1.0
Home	259	4.4
Other	91	1.5
Unknown	396	6.7

*Residential institution includes: community hospital/long stay unit, homeless facility, mental health facility and nursing homes.

Table 6: Summary of COVID-19 deaths linked to outbreaks in Ireland, cases with a date of notification from 01/03/2020 to 04/01/2022

		Number of Deaths	Percentage of total deaths	Percentage of deaths linked to outbreaks
Total number of deaths		5,952	100.0	
Deaths linked to outbreaks		3,647	61.3	100.0
Outbreaks by location	Nursing homes	2,245	37.7	61.6
	Hospital	875	14.7	24.0
	Community Hospitals/Long-stay units	146	2.5	4.0
	Residential institutions	111	1.9	3.0
	Other locations*	270	4.5	7.4

* Other locations include community outbreak, extended family, hotel, other, other healthcare service, private house, public house, religious/other ceremony, workplace

A death linked to an outbreak in a particular setting does not of itself indicate that transmission occurred within that setting. A case may be detected as part of an outbreak investigation, and associated with an outbreak, despite exposure and transmission having occurred elsewhere.

Table 7: Number of COVID-19 deaths in Ireland by month of death, cases with a date of notification from March 2020 to January 2022

Year	Month	Number of deaths *
2020	March	129
2020	April	1,160
2020	May	363
2020	June	67
2020	July	17
2020	August	7
2020	September	39
2020	October	132
2020	November	190
2020	December	195
2021	January	1,417
2021	February	892
2021	March	258
2021	April	101
2021	May	43
2021	June	20
2021	July	22
2021	August	87
2021	September	179
2021	October	217
2021	November	230
2021	December	157

* Date of death reported for 5,922 of the deaths.




Figure 1: Total number of COVID-19 deaths in Ireland and cumulative number by week of death, cases with a date of notification from 01/03/2020 to 04/01/2022. Date of death reported for 5,922 of deaths.

Acknowledgments

Sincere thanks are extended to all those who are participating in the collection of data and reporting of data used in these reports. This includes the HSE COVID-19 Contact Management Programme (CMP), staff in ICU units, notifying clinicians, laboratory staff, public health doctors, nurses, surveillance scientists, microbiologists and administrative staff.

Report prepared by COVID-19 Epidemiology Team, HPSC

Technical Notes

1. Data Source

Data are based on statutory notifications and were extracted from Computerised Infectious Disease Reporting (CIDR) system at 08:28 on 05/01/2022. Data are provisional and subject to ongoing review, validation and update. As a result, figures in this report may differ from previously published figures.

2. Population Data

Population data were taken from Census 2016. Data were aggregated into the following age groups for the analysis of deaths for the entire pandemic: 0-24 years, 25-34 years, 35-44 years, 45-54 years, 55-64 years, 65-74 years, 75-84 years and ≥ 85 years. Data were aggregated into the following age groups for the analysis of deaths with date of death in the past 7 days: 0-64 years, 65-74 years, 75-84 years and ≥ 85 years.

3. Definition of a COVID-19 death used by HPSC

For surveillance purposes, COVID-19 deaths include deaths in all possible, probable and confirmed COVID-19 cases (as per the COVID-19 case definition) and all should be notified, unless there is a clear alternative cause of death that cannot be related to COVID-19 infection (e.g. trauma). There should be no period of complete recovery* from COVID-19 between the illness and death. All COVID-19 deaths are notified regardless of the setting, including home, community and hospital settings. HPSC reports all deaths among these COVID-19 cases as outlined above and does not just confine the death reporting to those who die within 28 days of a positive test. This is in line with how COVID-19 cases are reported by the majority of European countries and follows WHO guidance for COVID-19 death surveillance.

*Please note that discharge from ICU or hospital is not in itself evidence of recovery. To determine if the case had recovered, it should be based on clinical assessment or alternatively a period of ≥ 3 months must have elapsed since the case was initially diagnosed with COVID-19 and the case must have no evidence of COVID-19 infection prior to death as assessed by a clinician.

Deaths not reported as COVID-19 deaths:

1. Persons with COVID-19 may die directly due to accidents. Such deaths are not due to COVID-19 and should not be certified as such. This decision not to certify as COVID-19 death will be based on clinical judgement.
2. In some instances, a death due to COVID-19 may not be attributed to another disease (e.g. cancer) and would be counted as a COVID-19 death independently of pre-existing conditions that are suspected of triggering a severe course of COVID-19.

More resources:

[COVID-19 interim case definition](#)

[Epidemiology of COVID-19 in Ireland Frequently Asked Questions](#)