

Influenza in hospitalized children in Ireland in the pandemic and 2010/2011 season: risk factors for Pediatric Intensive Care Unit admission

J. Rebolledo^{1,2}, D. Igoe¹, J. O'Donnell¹, L. Domegan¹, M. Boland³, B. Freyne⁴, A. McNamara⁵, E. Molloy^{4,6}, M. Callaghan⁷, A. Ryan⁸, D. O'Flanagan¹

1. Health Protection Surveillance Centre, Health Service Executive, Dublin, Ireland
2. European Programme for Intervention Epidemiology Training (EPIET), European Centre for Disease Prevention and Control (ECDC), Stockholm, Sweden
3. Department of Public Health, Health Service Executive Dublin, Ireland
4. National Maternity Hospital, Holles Street, Dublin, Ireland
5. Department of Public Health, Health Service Executive Tullamore, Ireland
6. Our Lady's Children's Hospital, Crumlin, Dublin, Ireland
7. Department of Anaesthesia and Intensive Care Medicine, Our Lady's Children's Hospital, Crumlin, Dublin, Ireland
8. Department of Public Health, Health Service Executive Sligo, Ireland

Corresponding author:

Javiera Rebolledo, Health Protection Surveillance Centre
25-27 Middle Gardiner Street,
Dublin 1
Ireland
javierarebolledo@gmail.com

Running head: Risk factors for severe Influenza in children 2009-2011

Summary

Influenza causes significant morbidity and mortality in children. This study's objectives were to describe influenza A(H1N1)pdm09 during the pandemic, to compare it with circulating influenza in 2010/2011, and to identify risk factors for severe influenza defined as requiring admission to a paediatric intensive care unit (PICU). Children hospitalized with influenza during the pandemic were older, and more likely to have received antiviral therapy (AVT) than children hospitalized during the 2010/2011 season. In 2010/2011, only one child admitted to PICU with underlying medical conditions had been vaccinated. The risk of severe illness in the pandemic was higher in females and those with underlying conditions. In 2010/2011, infection with influenza A(H1N1)pdm09 as opposed to other influenza viruses was a significant risk factor for severe disease. An incremental relationship was found between the number of underlying conditions and PICU admission. These findings highlight the importance of improving low vaccination uptake and increasing the use of antivirals in vulnerable children.

Introduction

In April 2009, a novel influenza virus, influenza A(H1N1)pdm09, was identified in Mexico and the United States (US), and spread worldwide to become the first pandemic of the 21st century [1]. The estimated global death toll was more than 284,000, and the cumulative symptomatic attack rate was highest in children aged 0-17 years, ranging from 4-33% [2]. The number of paediatric deaths in children in the US was four times that recorded in the previous five influenza seasons [3]. Children with underlying metabolic and neurological conditions were identified as being at highest risk for severe complications [4].

In Ireland, two waves of influenza activity occurred between April 2009 and May 2011: one wave during the pandemic period (2009/2010) and one post pandemic wave in the 2010/2011 season. The first wave began in July and August 2009, 14 weeks after the first case of influenza A(H1N1)pdm09 was confirmed in Ireland in May 2009. This wave began with relatively low levels of influenza activity, with sentinel GP influenza-like illness (ILI) consultation rates increasing above baseline levels between mid-July and August 2009. During this initial increase in activity, the majority of cases were amongst teenagers associated with outbreaks in residential summer language camps. The Irish school holidays in July and August may have limited further spread among children and thus hospitalisations in children during this time. Influenza activity increased significantly with the return to school in September 2009 and peaked during mid-October 2009. During this pandemic wave the peak age specific ILI rates occurred in children aged 5-9 and 10-14 years in mid-October 2009. The post pandemic wave occurred between December 2010 and February 2011 and was characterized by a short sharp

rise and fall in influenza activity. This wave was associated with higher numbers of critical care admissions and influenza-associated deaths.

In Ireland, during the pandemic period, antiviral therapy was recommended for all children with influenza-like illness (ILI) who were either: less than five years of age, or hospitalized with their illness, or with underlying predisposing medical conditions. This advice continued during the 2010/2011 season, and was actively promoted via guidelines and clinical algorithms provided to General Practitioners (GPs) and Emergency Departments.

Pandemic vaccination became available in early November 2009. Initially children in risk groups defined by the National Immunisation Advisory Committee (NIAC), and those under five years of age were vaccinated. Subsequently all other children were offered vaccination in schools, clinics or general practice settings [5]. During the 2010/2011 influenza season, vaccination was recommended for children in the NIAC risk groups. The risk groups were expanded in the 2010/2011 season to include children with any condition that could compromise respiratory function (e.g. spinal cord injury, seizure disorder, or other neuromuscular disorder) especially those attending schools/day centres for children with intellectual and/or physical disabilities.

International papers have compared A(H1N1)pdm09 in children with previous seasonal influenza viruses, and found that influenza A(H1N1)pdm09 affected an older age group and those with a history of asthma and chronic disease, particularly neurological and metabolic conditions [6-8]. There is however limited literature comparing the A(H1N1)pdm09 pandemic with the subsequent 2010/2011 influenza season [9-11].

The objectives of this paper are to describe influenza A(H1N1)pdm09 in hospitalized children during the pandemic in Ireland and compare it to circulating influenza viruses during the 2010/2011 season; and to identify risk factors for severe influenza in children, defined as admission to a Paediatric Intensive Care Unit (PICU).

Methods

Data collection

In April 2009, enhanced surveillance was undertaken on all cases of confirmed influenza A(H1N1)pdm09 in children 14 years and younger deemed to require admission to hospital for management of their condition. All laboratory confirmed cases of influenza in Ireland were notified to Regional Departments of Public Health, and for each case notified, they established whether the person had been hospitalized. Regional Departments of Public Health or the treating hospital clinician completed the enhanced surveillance form on hospitalized cases. Data collected included demographic details, pre-morbid medical conditions, antiviral therapy, length of hospital stay (LOS), admission to PICU, and outcome [12]. Data were checked by Departments of Public Health for completeness and cases identified with missing or inconsistent data were followed up and the information was corrected. Surveillance and laboratory information was entered into the Irish Computerized Infectious Disease Reporting system (CIDR). For those children admitted to PICU, intensive care staff completed an additional form on admission and discharge from PICU. This included information on medical interventions, complications and clinical outcome while in PICU.

Epidemiological studies and study period

Influenza A(H1N1)pdm09 cases by time, place, and person were described. A retrospective cohort study was conducted to identify risk factors for admission to a Pediatric Intensive Care Unit (PICU).

Two study periods were compared and analysed; the pandemic (week 17 2009 to week 32 2010) and the 2010/2011 influenza season (week 40 2010 to week 20 2011).

Laboratory methods

Ireland's Health Service Executive (HSE) recommended that all children admitted to hospital with ILI be tested for influenza. The majority (84.2%) of specimens were tested at the National Virus Reference Laboratory for influenza A and B using multiplex RT-PCR [13]. Influenza A positive specimens were further sub-typed using multiplex RT-PCR. A small percentage (15.8%) of samples were tested in three other sites, using either in-house real time reverse transcription RT-PCR assays, or Prodesse ProFluTM+ and Prodesse ProFAST+; or an in-house uniplex system for influenza A. A small proportion of influenza A positive specimens were not subtyped (23/665, 3.4%).

Analysis of Data

Confirmed influenza age specific hospitalization rates per 100,000 population were calculated. Population data were taken from the Irish 2011 census.

Descriptive, univariate and multivariable analyses were conducted using SPSS[®] (SPSS Inc) version 16.0 and STATA version 12.0. In the univariate analysis we calculated proportions,

means and medians for each study period. We used Student's t test to analyse differences in normally distributed continuous variables and non-parametric Mann-Whitney and Kruskal-Wallis methods for comparison of skewed data. The X^2 or Fisher's exact test were used to compare categorical variables between groups. For the multivariable analysis we used a Log-linear model with log link function and a Poisson distribution to obtain adjusted risk ratios. We performed two different models: one to assess risk factors for being admitted to PICU during each period separately, and the second to look at the effect of the number of medical conditions on the risk of admission to PICU for both study periods. Age, gender and any variables with p values less than 0.25 in the univariate analysis were added to the model and a backward variable selection procedure was used for both models [14]. Non-typed and H3N2influenza A were excluded from the model and the use of antiviral therapy (AVT) was not included, as we did not have information on timing of use.

Ethical approval

Ethical approval was not required for the collection of these data, as it was part of national surveillance conducted under the Infectious Disease Regulations, 1981 as amended [15].

Results

Age specific hospitalization, PICU and mortality data

Table 1 provides age specific hospitalization, PICU and mortality rates per 100,000 population for each season. Overall, the age specific hospitalisation and PICU rates decreased with

increasing age, in each period. There was a shift in the age specific hospitalization and PICU admission rates to the younger age group (< one year) during the 2010/2011 season.

Descriptive analysis

Table 2 compares the pandemic period and the 2010/2011 influenza season in hospitalized children. Children hospitalized with confirmed influenza during the pandemic period were older and were treated more frequently with AVT compared to those in the 2010/2011 season. There was no significant difference in the likelihood of treatment with AVT by age during the pandemic period. However, during the 2010/2011 period (using the <1 age stratum as the reference), only children aged 10-14 were 1.6 times more likely to receive AVT ($p=0.034$; 95%CI= 1.09-2.35).

There was a significant association between increasing age and the presence of underlying medical conditions in both the pandemic and the 2010/2011 season (Table 3).

Vaccination status was only available for PICU admissions during the 2010/2011 influenza season. Information on vaccine uptake was not available for the hospitalized population in general. In the 2010/2011 season, of the 14 children admitted to ICU, 11 were over six months of age. Of these 11, eight had underlying medical conditions, and therefore were in risk groups for seasonal flu vaccination. Only one (12.5%) of these children had been vaccinated. This child had also not received pandemic vaccination. Six of the 11 children had also been eligible for pandemic vaccination. In the six children aged two years or more at onset of illness during the 2010/2011 season, three (50%) of them had received pandemic vaccine.

PICU clinical course and outcome

In the analysis of all children admitted to PICU in both study periods; 29 (76.3%) underwent mechanical ventilation, 19 (51%) had Acute Respiratory Distress Syndrome (ARDS) and two (5.6%) had acute kidney injury. Complications included primary influenza viral pneumonia in 29 (80.6%), secondary bacterial pneumonia in eight (22.2%) and sepsis/multiorgan failure in four (11.1%). No child required Extracorporeal Membrane Oxygenation (ECMO) and no child developed myocarditis or encephalitis. The median length of stay (LOS) in PICU was seven days, and three (8.6%) of patients admitted to PICU died. There was no significant difference in the clinical course and outcome between children with and without underlying risk factors.

Effect of the number of medical conditions on risk of admission to PICU

Both univariate and multivariable analysis showed that the risk of being admitted to PICU increased with the number of underlying conditions present (Table 4). After adjustment by age and gender, the risk of being admitted to PICU during the pandemic period was 4.8, 10.91 and 73.12 times higher in children that had one, two, and three or more medical conditions, respectively. The increase in risk with the number of underlying conditions present was similar in the 2010/2011 season.

Risk of admission to PICU

Table 5, shows that during the pandemic, the risk of admission to PICU was lower in males and higher in those with respiratory conditions/asthma, heart conditions, immunological or renal

conditions. Some of the numbers affected with these conditions were however quite small. Age was not a risk for PICU admission during the pandemic. In the 2010/2011 influenza season, influenza A(H1N1)pdm09 emerged as a risk factor for admission to PICU, with the risk of PICU admission being 7 times higher with A(H1N1)pdm09 than with other flu strains. The majority of these were influenza B (Table 2). The risk of admission was also raised in those with neurological or metabolic conditions. There was no increased risk associated with age or gender.

Discussion

This is the first detailed comparison of influenza during the pandemic and the 2010/ 2011 season in hospitalized children in Ireland. As there is a national enhanced surveillance system in place, age specific hospitalization, PICU, and mortality rates could be calculated.

In Ireland during the pandemic, children were affected disproportionately, with relative sparing of the older population, similar to findings elsewhere [2]. The 2010/2011 influenza season was shorter but more intense than the pandemic, with a slightly higher overall ILI rate and significant pressure on adult intensive care units [16]. In children the age specific ILI rates were much lower during the 2010/2011 season, but still higher than those recorded in influenza seasons prior to the pandemic. The age specific hospitalization and PICU admission rates were lower for all ages with the exception of children aged less than one year. The lower impact in children during the 2010/2011 season may reflect the effect of interventions such as vaccination, as well as exposure to A(H1N1)pdm09 in the paediatric population. Unfortunately

there are no Irish sero-epidemiological data which could confirm this. In Ireland, pandemic specific vaccination uptake was high in children; 61.5% in children aged 0-4 years and 53.1% in those aged 5-14 years [17].

Our pandemic hospitalization data showed similarities with other countries: the median age (five years) of children hospitalized was similar to that found in large studies in Canada and Korea, but slightly lower than that found in Japan [18-20]. The proportion (34%) with underlying medical conditions was lower than that seen in some countries, 43% in Germany, and 42% of cases in Australia [5,21], but in line with or higher than other large population based studies (32% in the US, 27.5% in Madrid and 22.4% in Korea) [6, 22-23]. The proportion of hospitalized children requiring PICU admission was low at five per cent when compared with that found in Canada and Argentina at 16.8% and 19% respectively, but similar to Australia at 4.6% [6,21,24,28]. It has been noted in previous analyses that Irish data are similar in demographics, admission rates and bed occupancy to those in Australia and New Zealand [25]. However this may also reflect different thresholds for admission to PICU. There were more males than females hospitalized in Ireland during the pandemic, as was found in many other studies [19, 26, 27]. Length of hospital stay was similar during the pandemic and the 2010/2011 season, and also similar to that found in other studies [4, 28].

The presence of medical conditions increased with age, apart from the older age group (10-14 years). This was also found in Germany by Altman in his study of children admitted to PICU, and by Bettinger in Canada, who found that a third of hospitalized children aged less than two years had at least one medical condition versus 72% in older children [18, 29].

Significant differences were observed in the age of those hospitalized between both periods. Post pandemic, hospitalized children were younger. In a sentinel surveillance system for severe influenza in England, they also found that among hospitalized children, there was a shift away from the 5- 14 year age group in the post pandemic season, and an increase in the rate of hospitalization in children aged 0-4 years, with a trend of decreasing hospitalisation rates with increasing age in both periods [9].

Other studies comparing pandemic influenza with previous seasons have found that children admitted with influenza A(H1N1)pdm09 were older than those admitted with seasonal influenza A in the years prior to the pandemic [3,7,8,25,31-33]. Apart from age, no other differences were found in hospitalized and PICU cases during the pandemic and the 2010/2011 season. This was a similar finding to a Canadian study which also found no differences in the proportions requiring PICU or in the mortality rates [28].

Unfortunately, we found an extremely low vaccination uptake among children at high risk in the 2010/2011 season. Only one child in PICU of those eligible for vaccination had been vaccinated. A similar pattern was seen with fatal cases in the US during the 2010/2011 season [33]. Of 74 children who died aged six months or older, and in whom vaccination records were available, 39 had high risk conditions and only one third of these had been vaccinated.

Antivirals were used more frequently during the pandemic than in the 2010/2011 season. This is of concern, as the same recommendations for use were made during both influenza seasons, i.e. to treat all those sick enough to be hospitalized with antivirals. They were promoted strongly to general practitioners, consultants in emergency medicine, and paediatricians. This

has been found elsewhere too. In the US, the use of antivirals in hospitalized children also dropped from 77% to 56% in the post pandemic season [34]. In our study, when use of antivirals was compared by age group, an increase was seen in their use in those aged less than five years of age in the 2010/2011 season versus the pandemic period, whereas in older children, they were less used. This may reflect a welcome heightened awareness of the risk of influenza in younger children, but points to the need to re-emphasize the benefits of antiviral treatment in all hospitalized children. r.

We were able to identify and compare risk factors for admission to PICU with influenza during the pandemic and the 2010/2011 season. During the pandemic, the risk of admission to PICU was higher in female children, and in those with underlying medical conditions. This finding of female predominance is in contrast to previous published studies showing male predominance of severe disease associated with influenza A(H1N1)pdm09 [35-36]. This was not explained by a higher incidence of medical conditions in girls. Age group was not a risk factor for PICU admission. A comparison of pandemic with prior seasonal influenza showed similar findings [7].

During the 2010/2011 season the risk of admission to PICU was higher in those with influenza A(H1N1)pdm09, and in those with underlying medical conditions, but was not higher in females. In a study on seasonal influenza subtypes in children, it was found previously that disease due to influenza A(H3N2) was significantly more severe than that due to influenza A(H1N1) and influenza B [37]. Our findings provide evidence that disease due to influenza A(H1N1)pdm09 is also more severe than that due to influenza B. We also demonstrated a clear

incremental relationship between the number of underlying medical conditions and the risk of PICU admission with influenza.

There were some limitations to this study. Not all hospitalized children with influenza might have been clinically suspected or tested and identified, and so the true numbers may be underrepresented. There were some limitations to the data collected. We were unable to capture data on the timing of administration of antivirals, and so we were unable to assess whether the use of antivirals affected risk of admission to PICU. The data on respiratory conditions and asthma were not sufficiently detailed to allow us to separate them out into discrete categories. The numbers overall in PICU were small, and this limited our analysis.

Conclusions and recommendations

In a large national population based surveillance system, the epidemiology of influenza A(H1N1)pdm09 was described and contrasted with subsequent seasonal influenza. Risk factors for severe illness among hospitalized children with influenza were identified, including the fact that having influenza A(H1N1)pdm09 subtype is a risk factor for PICU admission.

The significantly increased risk of a serious episode of illness in those with underlying conditions, and also the incremental relationship between the number of medical conditions present and the risk of a severe episode present powerful evidence of the importance of targeting vaccination programmes to children with complex underlying medical conditions. These findings should be used to influence and encourage vaccination in these extremely

vulnerable groups. The findings also reinforce the importance of using antivirals in these children.

The use of antiviral therapy in all hospitalized children should also be promoted, with a focus not only on young children, but also those over five years of age, give our findings.

This study demonstrates that the collaborative efforts of clinicians, intensivists, GPs, public health and surveillance staff in surveillance activities can yield significant new information for action and response.

Acknowledgements

We thank Ajay Oza for his work on the database for patients admitted to PICU, and enabling us to merge this dataset with the national Computerised Infectious Disease Reporting system (CIDR). [We thank Sylvanus Fonguh from the Belgian Scientific Institute of Public Health for his advice in statistics.](#) We thank staff in Departments of Public Health, Paediatric Intensive Care Units, General Practitioners in the sentinel GP surveillance scheme, the Irish college of General Practitioners, the National Virus Reference Laboratory, University College Galway, Cork University Hospital, and staff at the Health Protection Surveillance Centre for their contribution to surveillance.

References

1. Dawood FS, *et al.* Emergence of a novel swine-origin influenza A (H1N1) virus in humans. *New England Journal of Medicine* 2009; **360**: 2605-2615.
2. Dawood FS, *et al.* Estimated global mortality associated with the first 12 months of 2009 pandemic influenza A H1N1 virus circulation: a modelling study. *The Lancet Infectious Diseases* 2012; **12**: 687-695.
3. Cox CM, *et al.* 2009 pandemic influenza A (H1N1) deaths among children – United States, 2009-2010. *Clinical Infectious Diseases* 2011; **52** (Suppl. 1):S69-S74.
4. Stein M, *et al.* Hospitalization of children with influenza A(H1N1) virus in Israel during the 2009 outbreak in Israel: a multicenter survey. *Archives of Pediatrics and adolescent Medicine* 2010; **164**: 1015-1022.
5. Health Service Executive Immunisation Website. Pandemic (H1N1) 2009 influenza. It stops with you. Frequently asked questions for health professionals. 2010 Jan <http://www.immunisation.ie/en/HealthcareProfessionals/PandemicCampaign2009/>. Accessed 13th November 2012.
6. Gordon R, *et al.* Pediatric hospitalisations associated with 2009 Pandemic Influenza (H1N1) in Argentina. *New England Journal of Medicine* 2010; **362**: 45-55.
7. Morris SK, *et al.* A retrospective cross-sectional study of risk factors and clinical spectrum of children admitted to hospital with pandemic H1N1 influenza as compared to influenza A. *British Medical Journal Open*. Published online 11 March 2012. doi:10.1136/bmjopen-2011-000310.

8. Engelhard D, *et al.* Increased extent of and risk factors for pandemic (H1N1) 2009 and seasonal influenza among children, Israel. *Emerging Infectious Diseases* 2011; **17**: 1740-1743.
9. Bolotin S, *et al.* A new sentinel surveillance system for severe influenza in England shows a shift in age distribution of hospitalised cases in the post-pandemic period. *Public Library of Science one* Published online 23 January 2012.
doi:10.1371/journal.pone.0030279.
10. Mytton OT, Rutter PD, Donaldson LJ. Influenza A(H1N1)pdm09 in England, 2009 to 2011: a greater burden of severe illness in the year after the pandemic than in the pandemic year. *Eurosurveillance*. Published online 5 April 2012. pii: 20139.
11. Altmann M, *et al.* Unchanged severity of influenza A(H1N1)pdm09 infection in children during first postpandemic season. *Emerging Infectious Diseases*. Published online 9 November 2012. doi: 10.3201/eid1811.120719.
12. Cullen G, *et al.* Surveillance of the first 205 confirmed hospitalised cases of pandemic H1N1 influenza in Ireland, 28 April – 3 October 2009. *Eurosurveillance*. Published online 5 November 2009. pii: 19389.
13. Carr M, *et al.* Development of a real-time RT-PCR for the detection of swine-lineage influenza A (H1N1) virus infections. *Journal of Clinical Virology*. 2009; **45**: 196-199.
14. Hosmer DW, Lemeshow S: Applied Logistic Regression. New York: Wiley; 2000
15. The Irish Statute Book (Government of Ireland). Infectious Diseases Regulations 1981. S.I. No. 390 of 1981. Available at: <http://www.irishstatutebook.ie/1981/en/si/0390.html>

16. Domegan, L *et al.* Epidemiology of influenza in Ireland for the 2010/2011 influenza season. *Epi-insight*. 2011; 12.
<http://ndsc.newsweaver.ie/epiinsight/dobfuin7on4?a=1&p=18519315&t=17517774>.
Accessed 10 July 2013.
17. Cotter S, *et al.* National Summary of Influenza pandemic (H1N1) 2009 vaccination in Ireland – provisional data. *Epi insight*
<http://ndsc.newsweaver.ie/epiinsight/viz7qx9a6lrqkeph6tk9uv>. Accessed 13 November 2012.
18. Bettinger JA, *et al.* Pandemic influenza in Canadian children: a summary of hospitalized pediatric cases. *Vaccine* 2010; **28**: 3180-3184.
19. [Costello BE, *et al.* Pandemic H1N1 Influenza in the Pediatric Emergency Department: A Comparison With Previous Seasonal Influenza Outbreaks. *Annals of Emergency Medicine* 2010; **56** \(6\): 643-648.](#)
20. Kobayashi M, *et al.* Pediatric hospitalizations with influenza A infection during the 2009-2010 pandemic in five hospitals in Japan. *Pediatrics international* 2012; **54**: 613-8.
21. Larcombe P, *et al.* Pandemic (H1N1) 2009: a clinical spectrum in the general paediatric population. *Archives of disease in childhood* 2011; **96**: 96-98.
22. Ko J.H., *et al.* Characteristics of Hospitalized Children with 2009 Pandemic Influenza A (H1N1): A Multicenter Study in Korea. *Journal of Korean Medical Science*. 2012 Apr;27(4):408-15
23. Del Rosal T, *et al.* Pandemic H1N1 influenza-associated hospitalizations in children in Madrid, Spain. *Influenza and other Respiratory viruses*. 2011 Nov;5(6):e544-51.

24. O'Riordan S, *et al.* Risk factors and outcomes among children admitted to hospital with pandemic H1N1 influenza. *Canadian Medical Association journal* 2010; **182**: 39-44.
25. Nicolay N, *et al.* Epidemiology, clinical characteristics and resource implications of pandemic (H1N1) 2009 in intensive care units in Ireland. *Critical care and resuscitation* 2010; **12**: 255-261.
26. Wieching A, *et al.* Clinical characteristics of pediatric hospitalizations associated with 2009 pandemic influenza a(H1N1) in Bavaria, Germany. *BMC research notes* 2012; **5**: 304.
27. Çiftçi E, *et al.* Clinical and epidemiological features of Turkish children with 2009 pandemic influenza A (H1N1) infection: experience from multiple tertiary paediatric centres in Turkey. *Scandinavian journal of infectious diseases* 2011; **43**: 923-929.
28. Tran D, *et al.* Comparison of children hospitalised with seasonal versus pandemic influenza A, 2004-2009. *Pediatrics* 2012; **130**: 397-406.
29. Altmann M, *et al.* Severe cases of pandemic (H1N1) 2009 in children, Germany. *Emerging infectious diseases* 2011; **17**: 186-192.
30. Blumental S, *et al.* Pandemic A/H1N1v influenza 2009 in hospitalized children: a multicenter Belgian survey. *BMC infectious diseases* 2011; **11**: 313.
31. Van Zwol A, *et al.* Clinical features of a Dutch cohort of critically ill children due to the 2009 new influenza A H1N1 pandemic. *Clinical pediatrics* 2011; **50**: 69-72.
32. Bagdure D, *et al.* Hospitalized children with 2009 pandemic influenza A (H1N1): comparison to seasonal influenza and risk factors for admission to the ICU. *Public Library of Science one*. Published online 15 December 2010. doi:10.1371/journal.pone.0015173.

33. Centers for Disease Control and Prevention (CDC). Influenza-associated pediatric deaths-
-United States, September 2010-August 2011. *MMWR Morbidity and mortality weekly report* 2011; 60: 1233-1238.
34. Garg S, *et al.* Reduced influenza antiviral treatment among children and adults
hospitalised with laboratory confirmed influenza infection in the year after the 2009
pandemic. *Clinical Infectious diseases*. Published online 27 April 2012. doi:
10.1093/cid/cis442
35. Xu C, *et al.* (2013) Characteristics of Hospitalized Cases with Influenza A (H1N1)pdm09
Infection during First Winter Season of Post-Pandemic in China. *PLoS ONE* 8(2): e55016.
doi:10.1371/journal.pone.0055016
36. Choi WI, *et al.* Male predominance of pneumonia and hospitalization in pandemic
influenza A (H1N1) 2009 infection. *Indian J Med Res*. 2012 Apr;135(4):534-7.
37. Esposito S, *et al.* Clinical and socioeconomic impact of different types and subtypes of
seasonal influenza viruses in children during influenza seasons 2007/2008 and
2008/2009. *BMC Infectious diseases*. Published online 12 October 2011. doi:
10.1186/1471-2334-11-271.

Table 1: Age specific, hospitalization Paediatric Intensive Care Unit (PICU) admission and mortality rates (per 100,000 population) for laboratory confirmed cases, by pandemic period and 2010/2011 season, Ireland.

Years of age	Hospitalized		In PICU		Influenza deaths	
	Pandemic	2010/2011	Pandemic	2010/2011	Pandemic	2010/2011
<1	93.9	98.1	5.5	8.3	0	0
1-4	58.5	41.2	4.2	2.1	0	0.7
5-9	47.4	16.5	1.6	0.3	0.6	0
10-14	31.7	12.6	1	0.3	0.3	0

Rate calculated using Central Statistics Office Population Census: 2011

Table 2: Demographic characteristics, length of stay in hospital, outcome, and underlying medical conditions in hospitalized children and children in Paediatric Intensive Care Units, by period in Ireland

Total hospitalization (N=761)			
	Pandemic period (n= 482)	2010/2011 influenza season (n=279)	<i>p</i>
	n (%)	n (%)	
Age			
Median (min-max)	5 (0-14)	2 (0-14)	<0.001
Age group			
<1	71 (14.7)	77 (27.6)	<0.001
1-4	163 (33.8)	111 (39.8)	
5-9	152(31.5)	53 (19)	
10-14	96 (19.9)	38 (13.6)	
Gender *			
Male	291 (60.6)	150 (53.8)	0.07
Female	189 (39.4)	129 (46.2)	
Influenza type/subtype			
A	NA	23 (8.2)	<0.001
A (H1N1)pdm09	482 (100)	158 (56.6)	
A (H3)	NA	2 (0.7)	
B	NA	96 (34.4)	
LOS [†]			
Mean	4.15	4.15	0.99
Median (Min-max)	2 (0-79)	3 (1-35)	
Admitted to PICU	24 (5.0)	14 (5.0)	0.98
Outcome [‡]			
Death	3 (0.6)	2 (0.7)	0.88
Presence of any medical condition [§]	154 (33.6)	79 (34.5)	0.82
Specific medical condition			
Respiratory and/or Asthma	73 (18.6)	34 (16.2)	0.46
Heart	18 (4.6)	12 (5.8)	0.53
Metabolic	12 (3.1)	3 (1.4)	0.23
Immunological	24 (6.2)	7 (3.3)	0.14
Hepatic	5 (1.3)	4 (2.0)	0.52
Neurological	33 (6.8)	20 (7.2)	0.87

Renal	5 (1.3)	4 (2.0)	0.52
Presence of medical conditions [§]			
No medical condition	304 (66.4)	150 (65.5)	
1 medical condition	117 (25.5)	65 (28.4)	0.33
2 medical conditions	27 (5.9)	13 (5.7)	
≥ 3 medical conditions	10 (2.2)	1 (0.4)	
On Antiviral treatment 	285 (67.7)	77 (38.9)	<0.001

* n = 480 for total hospitalization in the pandemic period

[†] n = 480 for total hospitalization in the pandemic period and n = 217 for total hospitalization in the 2010/2011 influenza season

[‡] n = 378 for total hospitalization in the pandemic period and n = 277 for total hospitalization in the 2010/2011 influenza season

[§] n = 458 for total hospitalization in the pandemic period and n = 228 for total hospitalization in the 2010/2011 influenza season

^{||} n = 18 for in PICU in the pandemic period and n = 11 for in PICU in the 2010/2011 influenza season

NA: Not Applicable

Table 3: Relationship between a) age and presence of underlying medical conditions, or b) being on antiviral treatment, for all hospitalized children during the pandemic and 2010/2011 season, Ireland

Years of age	Having any underlying medical conditions (N=233)					Received antiviral treatment (N=362)				
	Pandemic Period [*] (n=154)		2010/2011 influenza season [†] (n=79)		<i>p</i>	Pandemic period [‡] (n=285)		2010/2011 influenza season [§] (n=77)		<i>P</i>
	n	%	N	%		n	%	n	%	
<1	14	9.1	16	20.3	0.04	38	13.3	16	20.8	0.06
1-4	48	31.2	29	36.7		101	35.4	32	41.6	
5-9	59	38.2	22	27.8		94	33.0	14	18.2	
10-14	33	21.4	12	15.2		52	18.2	15	19.5	

* the p value for differences of having medical conditions among age groups within the Pandemic period is p=0.02

† the p value for differences of having medical conditions among age groups within the 2010/2011 influenza season is p=0.02

‡ the p value for differences of having received antiviral treatment among age groups within the Pandemic period is p=0.56

§ the p value for differences of having received antiviral treatment among age groups within the 2010/2011 influenza season is p=0.09

Table 4: Univariate and multivariate analysis of the effect of the number of medical conditions on risk of admission to PICU, for hospitalized children during the Pandemic and 2010/2011 season (n=687), Ireland.

Variable (Reference)	Univariate								Multivariable*					
	Pandemic period (n=428)				2010/2011 season (n=279)				Pandemic period			2010/2011 season		
	n	RR	P	95% CI	n	RR	P	95% CI	RR	P	95% CI	RR	P	95% CI
Medical Conditions (No medical condition)														
1 Medical Condition	117	1.20	0.676	0.56-2.82	65	1.40	0.53	0.049_4.02	4.79	0.010	1.45-15.91	2.83	0.093	0.84-9.53
2 medical condition	27	3.19	0.021	1.17_8.68	13	6.65	0.001	2.41_18.35	10.91	0.001	2.83-42.11	20.37	<0.001	7.29-56.82
≥ 3 medical condition	10	26.88	<0.001	15.68-46.07	1	-	0.798	-	73.12	<0.001	26.15-204.48	-	<0.001	-

* Adjusted by age and gender

Table 5: Univariate and multivariate analysis of risk factors for PICU admission for hospitalized children, by period, Ireland.

Variable (Reference)	Univariate								Multivariable					
	Pandemic period (N=428)				2010/2011 season (N=279)				Pandemic period [†]			2010/2011 season ^{††}		
	n	RR	P	95% CI	n	RR	P	95% CI	RR	P	95% CI	RR	P	95% CI
Age (<1 year)														
1-4	163	1.96	0.086	0.90-4.26	111	1.14	0.81	0.40-3.18						
5-9	152	0.57	0.247	0.22-1.50	53	0.33	0.246	0.04-2.45						
10-14	96	0.57	0.351	0.17-1.89	38	0.49	0.468	0.07-3.62						
Gender (Male)	291	0.46	0.051	0.21-1.02	150	0.86	0.772	0.31-2.31	0.48	0.041	0.24-0.97			
Influenza type* A(H1N1)pdm09					158	7.90	0.015	1.05-59.43				7.36	0.052	0.98-55.15
Medical condition														
Respiratory and/or Asthma	73	2.33	0.04	1.03-5.29	34	2.07	0.193	0.69-6.22	3.14	<0.001	1.66-5.96			
Heart	18	4.36	0.003	1.66-11.50	12	4.43	0.010	1.42-13.80	4.08	0.001	1.72-9.66			
Metabolic	12	11.18	<0.001	5.37-23.24	3	5.26	0.064	0.98-28.3				6.30	0.018	1.37-29.03
Immunological	24	5.37	<0.001	2.33-12.36	7	NA	0.472	NA	9.90	<0.001	5.16-19.01			
Liver	5	16.17	<0.001	8.70-30.05	4	NA	0.581	NA						
Neurological	34	13.65	<0.001	6.48-28.77	20	3.8	0.012	1.31 – 11.01				6.07	<0.001	2.39-15.43
Renal	5	3.48	0.18	0.58-21.05	4	NA	0.581	NA	6.40	<0.001	0.02-0.09			

* Results only for the 2010/2011 influenza season, because the pandemic period had only had one type of Influenza

[†] Variables taken out from the Log-linear model for the pandemic period were: age, metabolic, liver and neurological conditions

^{††} Variables taken out from the Log-linear model for the 2010/2011 season were: age, gender, respiratory and/or asthma, heart, metabolic, immunological and liver conditions