

Influenza Weekly Surveillance Report



A REPORT BY THE HEALTH PROTECTION

SURVEILLANCE CENTRE IN COLLABORATION WITH THE IRISH COLLEGE OF GENERAL PRACTITIONERS,
THE NATIONAL VIRUS REFERENCE LABORATORY & THE DEPARTMENTS OF PUBLIC HEALTH

Week 23 2009 (1st-7th June 2009)

Summary

During week 23 2009, seasonal influenza activity decreased in Ireland. Influenza-like illness (ILI) consultation rates decreased and are well below baseline thresholds. Nine non-sentinel specimens tested positive by the NVRL for influenza during week 23 2009: eight specimens, from five cases, were positive for influenza A(H1N1) and one specimen was positive for influenza A. See the link to the separate report on Influenza A(H1N1) infection for Week 23 2009 on page 2.

Background

This is the ninth season of influenza surveillance using computerised sentinel general practices in Ireland. The Health Protection Surveillance Centre (HPSC) is working in collaboration with the Irish College of General Practitioners (ICGP), the National Virus Reference Laboratory (NVRL) and the Departments of Public Health on this sentinel surveillance project. Fifty-nine sentinel general practices have been recruited to report on the number of patients with ILI on a weekly basis. ILI is defined as the sudden onset of symptoms with a temperature

of 38°C or more, with two or more of the following: headache, sore throat, dry cough and myalgia. Sentinel GPs send a combined nasal and throat swab, to the NVRL, on at least five patients per week where a clinical diagnosis of ILI is made during the influenza season. This report includes data on ILI cases reported by sentinel GPs, influenza test results from the NVRL, influenza notifications, registered deaths attributed to influenza, and regional influenza activity reported by the Departments of Public Health.

Results

Clinical Data

Sentinel GPs reported five ILI cases during week 23 2009. This corresponds to an ILI consultation rate of 2.8 per 100,000 population, which is a decrease in comparison to the updated rate of 4.0 per 100,000 population reported during week 22 2009 (figure 1). This rate is well below the baseline threshold of 17.8 per 100,000 population. Forty-five of 59 (76.3%) sentinel general practices reported during week 23 2009, with three reporting ILI cases.




Figure 1: Influenza baseline activity threshold and GP consultation rate for ILI per 100,000 population by week, during the 2000/2001*, 2006/2007, 2007/2008, 2008/2009 and Summer 2009 influenza seasons

*Highest recorded levels of ILI activity since initiation of sentinel surveillance

Results (continued)

During week 23 2009, three ILI cases were reported in the 15-64 year age group (2.4 per 100,000 population) and two in the 65 years and older age group (10.1 per 100,000 population). No ILI cases were reported in the 0-4 year and 5-14 year age groups as shown in figure 2.




Figure 2: Age specific GP consultation rate for ILI per 100,000 population by week during the 2008/2009 and Summer 2009 influenza seasons

Influenza A(H1N1) infection

Please see separate report by HPSC on Influenza A(H1N1) infection for Week 23 2009, available at: <http://www.hpsc.ie/hpsc/A-Z/EmergencyPlanning/AvianPandemicInfluenza/SwineInfluenza/Surveillance%20Reports/>

Oseltamivir Resistance in Europe

During the 2008/2009 influenza season, the NVRL has conducted nucleotide sequencing on 10 influenza A(H1) specimens taken by sentinel GPs in Ireland, nine (90%) of which were resistant to oseltamivir and one was sensitive. During the 2007/2008-influenza season, seven of 63 specimens (11.1%) tested by the NVRL showed resistance to oseltamivir.

Preliminary results from antiviral drug susceptibility testing among seasonal influenza viruses circulating in Europe have revealed that some of the influenza A (H1N1) viruses in circulation this season are resistant to the antiviral drug, oseltamivir. In Europe, all influenza A(H3N2) viruses tested were sensitive to oseltamivir (n=653) and zanamivir (n=612), whereas 100% of the 644 A(H3N2) viruses tested were resistant to M2 inhibitors. Of the influenza A(H1N1) viruses analysed, 100% were sensitive to zanamivir (n=259) and 99.2% were sensitive to M2 inhibitors (n=124) but 98% were resistant to oseltamivir (n=259). One hundred and seventeen influenza B isolates were analysed and all were sensitive to oseltamivir and 113 influenza B isolates were all sensitive to zanamivir.

Latest information on oseltamivir resistance in Europe:

http://ecdc.europa.eu/en/Health_topics/Seasonal%20Influenza/Epidemiological_updates.aspx
http://ecdc.europa.eu/en/Health_Topics/influenza/antivirals.aspx

In the UK, since week 40 2008 all of the 237 influenza A(H3) isolates that have been tested for drug sensitivity have been found to be sensitive to oseltamivir and zanamivir, but resistant to amantadine. Eighty-six influenza A(H1) specimens have been tested for anti-viral drug resistance, 85 (98.8%) of these were resistant to oseltamivir and all were sensitive to zanamivir and amantadine. Fifty-five influenza B specimens have been tested and all were sensitive to oseltamivir and zanamivir.

Latest information on oseltamivir resistance in the UK:

<http://www.hpa.org.uk/web/HPAweb&Page&HPAwebAutoListName/Page/1191942171468>

Virological Data from the NVRL

Five sentinel specimens were tested by the NVRL during week 23 2009, none of which were positive for influenza. The NVRL tested 108 non-sentinel specimens taken during week 23 2009, mainly from hospitalised paediatric cases. Eight non-sentinel specimens, from five cases, tested positive for influenza A(H1N1) and one tested positive for influenza A. No specimens tested positive for respiratory syncytial virus (RSV) during week 23 2009 (table 1).

Please note that non-sentinel specimens include all specimens referred to the NVRL, these specimens are mainly from hospitals and some GPs and may include more than one specimen from each case.

Table 2 shows the number and percentage of sentinel specimens that tested positive for influenza, by type and subtype, for the Summer 2009 influenza season to date. Influenza positive specimens by HSE area and age group (in years), for week 23 2009 and the Summer 2009 season to date are shown in tables 3 and 4, respectively. Figure 3 compares the ILI consultation rates by season and the number of positive influenza specimens tested by the NVRL. Figure 4 compares the number and percentage of non-sentinel RSV positive specimens detected during the 2007/2008 and 2008/2009 influenza seasons and the respective Summer 2008 and 2009 seasons.

Table 1: Number of sentinel and non-sentinel* respiratory specimens and positive results for week 23 2009 and Summer 2009 season to date

Week Number	Specimen Type	Total Specimens	No. Influenza Positive	% Influenza Positive	Influenza A	Influenza B	RSV	% RSV Positive
23 2009	Sentinel	5	0	0.0	0	0	NA	NA
	Non-Sentinel	108	9	8.3	9	0	0	0
	Total	113	9	8.0	9	0	0	0
Summer 2009 season to date	Sentinel	19	0	0	0	0	NA	NA
	Non-Sentinel	290	18	6.2	18	0	5	1.7
	Total	309	18	5.8	18	0	5	1.6

Table 2: Number and percentage of positive sentinel specimens by type and subtype, Summer 2009 season to date

Summer 2009 season to date [†]						
Influenza (all types)		Influenza A (all subtypes)	Influenza B	Influenza A Unsubtyped	Influenza A(H1)	Influenza A(H3)
		(n=0)		(n=0)		
Number positive	0	0	0	0	0	0
% Positive	0.0	0.0	0.0	0.0	0.0	0.0

Table 3: Total number of sentinel and non-sentinel[†] influenza A and B positive specimens by HSE area for week 23 2009 and the Summer 2009 season to date

	Week 23 2009			Summer 2009 season to date		
	Flu A	Flu B	Total	Flu A	Flu B	Total
HSE-ER	2	0	2	4	0	4
HSE-M	0	0	0	2	0	2
HSE-MW	2	0	2	4	0	4
HSE-NE	0	0	0	0	0	0
HSE-NW	0	0	0	0	0	0
HSE-SE	0	0	0	3	0	3
HSE-S	5	0	5	5	0	5
HSE-W	0	0	0	0	0	0
HSE area unknown	0	0	0	0	0	0
Total	9	0	9	18	0	18

* Please note that non-sentinel specimens include all specimens referred to the NVRL, these specimens are mainly from hospitals and some GPs and may include more than one specimen from each case

[†] Number of sentinel specimens tested = 19

Table 4: Total number of sentinel and non-sentinel† influenza A and B positive specimens by age group (in years) for week 23 2009 and the Summer 2009 season to date

	Week 23 2009			Summer 2009 season to date		
	Flu A	Flu B	Total	Flu A	Flu B	Total
0-4 years	0	0	0	0	0	0
5-14 years	0	0	0	0	0	0
15-64 years	5	0	5	14	0	14
65 years and older	0	0	0	0	0	0
Age group unknown	4	0	4	4	0	4
Total	9	0	9	18	0	18




Figure 3: GP ILI consultation rate per 100,000 population and the number of positive influenza specimens detected by the NVRL by week and season, 2000/2001 - 2008/2009, including Summer 2009

†Please note that non-sentinel specimens include all specimens referred to the NVRL, these specimens are mainly from hospitals and some GPs and may include more than one specimen from each case




Figure 4. Number and percentage of non-sentinel RSV positive specimens detected during the 2007/2008 and 2008/2009 influenza seasons (data points from week number 21 onwards represent the summer seasons for 2008 and 2009)

Weekly Influenza Notifications

During week 23 2009 (week 22 2009 of epidemiological calendar), eight influenza A(H1N1) cases were notified to HPSC. Influenza cases, including those with influenza A(H1N1), notified to HPSC during the 2008/2009 and Summer 2009 influenza seasons are shown in figure 5 and compared to GP ILI consultation rates.




Figure 5: Number of notifications of influenza by type and week of notification compared to sentinel GP ILI consultation rates per 100,000 population during the 2008/2009 and the summer 2009 influenza seasons[‡]

Baseline thresholds

A baseline threshold of 17.8 cases per 100,000 population will be used alongside expert opinion to assess influenza activity during the 2008/2009 and Summer 2009 influenza season in Ireland. This baseline was derived from the EISS method using a mathematical algorithm to identify the influenza activity period of the previous eight seasons.

Mortality Data

No deaths attributed to influenza were reported to HPSC by the General Register Office during week 23 2009.

Outbreak Reports

No seasonal influenza/ILI outbreaks were reported to HPSC during week 23 2009. During the 2008/2009 season and the Summer 2009 influenza season to date, six seasonal influenza/ILI outbreaks have been reported to HPSC.

Regional Influenza Activity by HSE-Area

Influenza activity is reported on a weekly basis from the Departments of Public Health. Influenza activity is based on sentinel GP ILI consultation rates, laboratory confirmed influenza cases and ILI/influenza outbreaks. Each Department of Public Health has established at least one sentinel hospital in each HSE area, to report total hospital admissions, total emergency admissions and total respiratory admissions data on a weekly basis. Sentinel primary and secondary schools were also established in each HSE area in close proximity to the sentinel GPs, reporting absenteeism data on a weekly basis. During influenza week 22 2009, no increases in respiratory admissions were reported by sentinel hospitals. No data on absenteeism were received from sentinel schools still open (i.e. primary schools). No activity was reported by HSE-NE, while sporadic influenza activity (based on isolated cases of ILI and/or virological results) was reported by HSE-E, -M, -MW, -NW, -S, -SE and -W during week 22 2009 (figure 6).

[‡] Notification data are provisional and were extracted from [CIDR](#) on the 09/06/2009 at 18.40




Figure 6: Map of influenza activity by HSE area during week 22 2009

Influenza Activity in Northern Ireland

During week 23 2009, 10 ILI cases and one influenza case were reported in Northern Ireland, corresponding to a combined rate of 7.9 per 100,000 population, a decrease compared to the updated rate of 13.8 per 100,000 population reported during week 22 2009. During week 23 2009, two specimens tested positive for influenza A(H1N1). For more details see: <http://www.cdscni.org.uk/>

Influenza Activity in England, Scotland & Wales

The HPA's weekly influenza report recommenced on 28th May 2009.

Clinical rates increased across the UK in week 18 when the influenza A(H1N1) outbreak was declared. Most rates have subsequently declined. All rates are now at low levels, slightly higher than expected for this time of year but below the threshold levels, where they are set (30 and 50 consultations per 100,000 population in England and Wales, and Scotland, respectively). In week 22, the rates for all countries declined: England and Wales, 4.8 per 100,000 population (compared to 7.9 per 100,000 population in week 21); and Scotland, 6 per 100,000 population (compared to 7 per 100,000 population for week 21). The QSurveillance@ UK rate was 4.2 per 100,000 population in week 22 compared to 6.5 per 100,000 population in week 21. Please note that the GP consultation rates for England and Wales, Scotland and QSurveillance@ are for ILI, while for Northern Ireland it is a combined ILI and influenza rate, which in part explains the higher rates observed in Northern Ireland.

The Centre for Infections' Respiratory Virus Unit (RVU) reported 25 positive specimens for week 22 [2 A(H3), 1 B, 23 A(H1N1swl)].

Seasonal influenza surveillance data for the UK can be accessed at the following link:

<http://www.hpa.org.uk/servlet/Satellite?c=Page&childpagename=HPAweb%2FPage%2FHPAwebAutoListName&cid=1153999752025&p=1153999752025&pagename=HPAwebWrapper&searchmode=simple&searchterm=National+Influenza+Report>

Influenza A(H1N1) surveillance data for the UK can be accessed at the following link:

<http://www.hpa.org.uk/webw/HPAweb&Page&HPAwebAutoListName/Page/1240732817665?p=1240732817665>

Influenza Activity in Europe

Seasonal influenza surveillance data for Europe can be accessed at the following link:

<http://www.eiss.org/index.cgiUU>

Influenza A(H1N1) surveillance data for Europe can be accessed at the following link:

<http://ecdc.europa.eu/>

Influenza Activity in Canada

Seasonal influenza surveillance data for Canada can be accessed at the following link:

<http://www.phac-aspc.gc.ca/fluwatch/index.html>

Influenza A(H1N1) surveillance data for Canada can be accessed at the following link:

http://www.phac-aspc.gc.ca/alert-alerte/swine_200904-eng.php?utm_source=phac-goc-departments-eng&utm_medium=banner&utm_content=swineflu-landing-page&utm_campaign=swineflu

Influenza Activity in the United States

Seasonal influenza surveillance data for the United States can be accessed at the following link:

<http://www.cdc.gov/flu/weekly/fluactivity.htm>

Influenza A(H1N1) surveillance data for the United States can be accessed at the following link:

<http://www.cdc.gov/h1n1flu/>

Influenza Activity Worldwide

Global seasonal influenza surveillance data can be accessed at the following link:

<http://gamapserv.who.int/GlobalAtlas/home.asp>

Global influenza A(H1N1) surveillance data can be accessed at the following link:

<http://www.who.int/en/>

Avian Influenza

No further reports of avian influenza activity in Egypt were available on the WHO website (latest information: 2nd June 2009).

As of 2nd June 2009, 433 confirmed human cases and 262 (60.5%) deaths from avian influenza A (H5N1) have been reported to the WHO from Azerbaijan, Bangladesh, Cambodia, China, Djibouti, Egypt, Indonesia, Iraq, Lao Peoples Democratic Republic, Myanmar, Nigeria, Pakistan, Thailand, Turkey and Viet Nam.

Further information on avian influenza is available on the following websites:

WHO http://www.who.int/csr/disease/avian_influenza/en/

HPSC <http://www.ndsc.ie/hpsc/A-Z/Respiratory/AvianInfluenza/>

ECDC <http://ecdc.europa.eu/en/>

Northern Hemisphere Influenza Vaccine for the 2009/2010 Season

For the 2009/2010 influenza season in the Northern Hemisphere, the members of the WHO Collaborating Centres on Influenza have recommended that influenza vaccines for the contain the following strains:

- an A/Brisbane/59/2007 (H1N1)-like virus**
- an A/Brisbane/10/2007 (H3N2)-like virus^{††}
- a B/Brisbane/60/2008-like virus^{‡‡}

**A/Brisbane/59/2007 is a current vaccine virus; A/South Dakota/6/2007 (an A/Brisbane/59/2007-like virus) is a current vaccine virus used in live attenuated vaccines.

^{††}A/Brisbane/10/2007 and A/Uruguay/716/2007 (an A/Brisbane/10/2007-like virus) are current vaccine viruses.

^{‡‡}B/Brisbane/33/2008 is a B/Brisbane/60/2008-like virus.

http://www.who.int/csr/disease/influenza/recommendations2009_10north/en/index.html.

Further information on influenza can be found on the [HPSC website](#)

Acknowledgements

HPSC, ICGP and NVRL wish to thank the sentinel GPs who have participated in the GP sentinel surveillance system and who have contributed towards this report

This report was produced by Stephen Murchan, Sarah Jackson and Dr Joan O'Donnell, HPSC