

# Malaria in Ireland 2005

## Key Points

- In 2005, 44 cases of malaria were notified, an increase of 62% on the number reported in 2004
- The most common species reported was *P. falciparum*, accounting for 75% of cases notified
- The majority of cases became ill after exposure in sub-Saharan Africa
- Visiting family in country of origin was the most common reason reported for travel
- Only one malaria case in Ireland in 2005 reported full compliance with their prescribed course of malaria prophylaxis. The remaining cases either failed to take any malaria prophylaxis prior to exposure or failed to continue prophylaxis for the required time period.
- It is important that travellers to endemic areas are made aware of the need to be properly compliant with their antimalarial medication and anti-mosquito measures, and the potential health consequences of non- or partial compliance.

## Introduction

Malaria is the most important vectorborne disease in the world, with 400 million infections and around 1 million deaths annually. It is endemic in over 100 countries in sub-tropical and tropical areas of Africa, Central and South America, Asia, the Middle East and Oceania, with 90% of deaths occurring in Sub-Saharan Africa, mostly among young children.

Infection is caused by transmission of one of 4 species of *Plasmodium* (*P. falciparum*, *P. ovale*, *P. vivax* or *P. malariae*) through the bite of infected female anopheline mosquitoes. The species vary in their clinical effects, *P. falciparum* causing the most severe form of malaria, and the most deaths. TropNetEurop reported a case fatality of 1.4% for *P. falciparum* infections imported into Europe.<sup>1</sup> Malaria caused by other *Plasmodium* species is less severe. *P. ovale* and *P. vivax* also differ in that they have persistent liver stages, which can resist conventional treatment and can produce relapses up to a year after the initial infection.

Worldwide each year, up to 30,000 travellers fall ill with malaria on their return from visiting countries where the disease is endemic.<sup>2</sup> Pregnant woman, young children and the elderly are particularly at risk. Malaria in pregnancy increases the risk of maternal death, miscarriage, stillbirth and neonatal death. As malaria is a relatively rare disease in Ireland, a high level of suspicion is necessary when travel to endemic areas has occurred in either the recent or distant past.

Increasing numbers of Irish residents are travelling to malarious regions for holiday and business travel. An

Table 1. Malaria notifications, Ireland 2005 by country of exposure

Country of exposure	Number of notifications	% of all cases
Sub-saharan Africa	30	68%
<i>Nigeria</i>	22	50%
<i>Other than Nigeria</i>	8	18%
Asia	3	7%
Not reported	11	25%
<b>Total</b>	<b>44</b>	<b>100%</b>




Figure 1. Number of malaria notifications, Ireland 1982-2005

increasing proportion of the population in Ireland originates from malaria endemic regions and regularly travels home.

Malaria surveillance aims to document the burden of illness in Ireland, to define the characteristics of those most at risk, and to identify those who could benefit most from preventive messages. In addition, the data collected permit monitoring for prophylaxis failures that might indicate emergence of drug resistance. There is potential also to identify cryptic cases; these are cases where the route of transmission is unclear or unusual.

### Materials and Methods

Malaria has been notifiable in Ireland since 1948. The case definition adopted since 2004 is based on the EU case definition.<sup>3</sup> Since 2001, public health physicians have provided enhanced surveillance data, e.g. country of infection, reason for travel and use of chemoprophylaxis, where available to HPSC. Notification and enhanced surveillance data are maintained in the CIDR (Computerised Infectious Disease Reporting) database. The data used in this report are based on information retrieved from the CIDR database on malaria cases in 2005.

### Results

#### Malaria incidence

In 2005, 44 cases of malaria were notified (figure 1). This is an increase of 62% on the number reported in 2004, and equates to a crude annual incidence rate of 1.1 per 100,000 (95% CI 0.79-1.45).

Areas reporting the highest number of cases were the HSE-E (n=12), HSE-NE (n=10) and HSE-S (n=10). There were also 5

cases in the HSE-M, 1 in the HSE-MW, 2 in the HSE-NW, 2 in the HSE-SE and 2 in the HSE-W.

#### Species of Plasmodium

As in previous years, the most common species reported was *P. falciparum*, accounting for 75% of all cases notified (n=33). There were also three *P. vivax*, one *P. ovale*, one *P. malariae* and one mixed *P. vivax/P. ovale* infection. This is similar to the species distribution reported by the UK and in Europe for cases of imported malaria.<sup>4,5</sup>

#### Age and sex distribution

Twenty-one cases were male and 21 were female (unknown/unspecified=2). Cases ranged in age from 1 to 64 years. Females in the 25-34 age group were the most common age-sex group reported, and there were ten notifications (23%) in children under the age of 15 (figure 2).

#### Clinical features

Twenty-one cases were hospitalised, one case was described as a GP patient, and for 22 cases this information was not specified. In 2005 one death in an adult was reported as being due *P. falciparum*.

#### Country of infection

In 2005, there were no cases of airport, congenital, induced or introduced malaria reported. One relapsed case of *P. vivax* was reported. The remaining 43 cases were either reported as being, or were assumed to be, imported. Country of infection was recorded for 33 cases, the majority of whom were exposed in sub-Saharan Africa, with the remainder exposed in Asia (table 1).




Figure 2. Age-sex distribution malaria cases, Ireland 2005.

### Reason for travel

Reason for travel was recorded for 28 cases. The largest subgroup identified in 2005 was people who had travelled to visit family in their country of origin –over half of those for whom the information was available (n=15). New entrants made up a further quarter of cases (n=7), with the remainder reported as holidaymakers (n=1), business travellers (n=1), armed services (n=1), Irish citizen living abroad (n=1), other (n=2) and not specified (n=16).

### Use of chemoprophylaxis

Excluding new entrants (those who had spent their lives to date living in an endemic region would not be expected to be taking chemoprophylaxis), information on malaria prophylaxis was available for 21 of the remaining 37 cases. Of these, 17 (81%) took no prophylaxis, 3 (14%) took prophylaxis but failed to continue for the required period. Only one case reported full compliance with prescribed course of prophylaxis.

### Interval between arrival in Ireland and onset of symptoms

Malarial infections can occur up to several months following exposure. For 19 cases, data were available both for the date of arrival from malarious region and date of onset of illness. The interval between these dates varied between 0 and 181 days (median 7). Eighty-eight per cent of *P. falciparum* infections had an interval of less than 2 weeks although intervals of 6 months and longer can occur<sup>6</sup>. The longest interval recorded here was for a *P. vivax* case. *P. vivax* (and *P. ovale* and *P. malariae*) tend to have longer incubation periods than *P. falciparum*.<sup>6</sup>

### Discussion

In 2005, there were 44 cases of malaria notified in Ireland. Although malaria has been notifiable for many years, it is likely that there has been under-notification and some of the current increase undoubtedly reflects improved reporting consequent to the Infectious Diseases (Amendment) (No 3) Regulations (S.I. 707 of 2003).<sup>7</sup>

The majority of cases in 2005 were associated with travel in Africa (91% of those for whom country of infection information was available), parts of West Africa having some of the most intense malaria transmission in the world. Although in 2005, holidaymakers and business travellers formed only a small proportion of the cases reported, between 2001 and 2004 as many as 36% of malaria cases in Ireland specified holiday or business as their reason for travel.<sup>8</sup> The Health Protection Agency (HPA) in the UK has issued a number of advisories over the last couple of years, after several deaths and cases of severe malaria were reported in holidaymakers returning from The Gambia. Many cases had either failed to take any prophylaxis or had taken inadequate or inappropriate prophylaxis.

Increasing numbers of Irish residents were born or raised in countries endemic for malaria. In 2005, visiting family in country of origin was the most common reason given by cases for travel to an endemic region. In the UK, half of malaria cases occur in minority ethnic groups who have settled in the UK, that were visiting family and relations overseas when they acquired malaria.<sup>6</sup> Individuals in endemic regions build up immunity to malaria that fades rapidly while living in a malaria-free region like Ireland, and may wrongly

assume that they are still immune to the disease. Failure to take appropriate prophylaxis and failure to take sensible protection measure against biting mosquitos are key factors in acquiring malaria. Only one malaria case in Ireland in 2005 reported full compliance with their prescribed course of prophylaxis. The remaining cases either failed to take any malaria prophylaxis prior to exposure or failed to continue prophylaxis for the required time period. It is important that travellers to endemic areas are made aware of the need to be properly compliant with their antimalarial medication and anti-mosquito measures, and the potential health consequences of non- or partial compliance.

The HPA Advisory Committee on Malaria Prevention (ACMP) has published guidelines (and updates) on prevention of malaria.<sup>9</sup> Four steps remain essential to prevent malaria in travellers:

**Awareness:** know about the risk of malaria  
**Bites by mosquitoes:** prevent or avoid  
**Compliance with appropriate chemoprophylaxis.**  
**Diagnose breakthrough malaria swiftly and obtain treatment promptly.**

It is important that travellers to endemic areas are aware that preventive measures are not 100% effective, and that they should seek treatment promptly if they suffer symptoms suggestive of malaria within a year following their return, informing their physician of their travel history.

The guidelines of the HPA Advisory Committee on Malaria

Prevention in travellers have also stated the need for balancing the risk of malaria and the risk of adverse reactions to anti-malarials. This depends upon place to be visited, duration of the visit, degree of exposure, level of drug resistance and the type of traveller.<sup>9</sup>

#### Acknowledgements

We wish to acknowledge the co-operation of microbiologists, medical scientists, public health physicians, SPHMs, surveillance scientists, infection control nurses, PEHOs, and EHOs in providing the information on which this report is based.

#### References

1. Muhlberger et al. 2003 Age as a risk factor for severe manifestations and fatal outcome of falciparum malaria in European patients: observations from TropNetEurop and SIMPID Surveillance Data. *Clin Infect Dis*. 2003 Apr 15;36(8):990-5. Epub 2003 Apr 2
2. WHO 2005 International travel and health. <http://www.who.int/ith/en/>
3. HPSC . 2004. Case definition for Notifiable Diseases. <http://www.ndsc.ie/NotifiableDiseases/CaseDefinitions/>
4. HPA MRL. 2005. Malaria imported into the UK: implications for those advising travellers. *CDR Weekly*. Vol 15(22)
5. TropNet. 2006. TropNetEurop Friends & Observers Sentinel Surveillance Report: Malaria [http://www.tropnet.net/reports\\_friends/pdf\\_reports\\_friends/feb06\\_malaria2005\\_friends.pdf](http://www.tropnet.net/reports_friends/pdf_reports_friends/feb06_malaria2005_friends.pdf)
6. HPA. 2005. Foreign travel-associated illness, England, Wales, and Northern Ireland. Annual Report 2005
7. Infectious Diseases (Amendment) (No 3) Regulations (S.I. 707 of 2003).
8. Garvey, P. P. McKeown and D. O'Flanagan. 2006. Malaria in Ireland 2001-2004. FPHM Summer Scientific Meeting. May 2006. Poster abstract
9. Bradley DJ, Bannister B; Health Protection Agency Advisory Committee on Malaria Prevention for UK Travellers. Guidelines for malaria prevention in travellers from the United Kingdom for 2003. *Comm Dis Public Health*. 2003 Sep;6(3):180-99.